

Quark masses and low energy constants in the continuum from CLQCD ensembles



Yi-Bo Yang










中国科学院大学
University of Chinese Academy of Sciences




ICTP-AP
International Centre
for Theoretical Physics Asia-Pacific
国际理论物理中心-亚太地区

Collaborators: Zhi-Cheng Hu, Bo-Lun Hu, Ji-Hao Wang, Ming Gong,
Liuming Liu, Peng Sun, Wei Sun, Wei Wang, and Dian-Jun Zhao

| | Naive | Staggered/HISQ | Wilson/Clover | Twisted-mass | Overlap/Domain wall |
|--------------------------|---|---|---|--|--|
| Form | $D^{\text{naive}} = \gamma_\mu (\delta_{x,x+\mu} - \delta_{x,x-\mu})$ | $D^{\text{st}} = \gamma_\mu^{\text{st}(x)} (\delta_{x,x+\mu} - \delta_{x,x-\mu})$ | $D^{\text{clv}} = D + aD^2 + ac_{\text{sw}} F_{\mu\nu} \sigma^{\mu\nu}$ | $D^{\text{tm}} = D^{\text{clv}} + i\tau_3 m$ | $D^{\text{ov}} = [1 + \gamma_5 D(-\rho)] / \sqrt{D^\dagger(-\rho)D(-\rho)} / \rho$ |
| Fermion copies | 16 | 4 | 1 | 1 | 1 |
| Chiral symmetry breaking | N/A | $\mathcal{O}(a^4)$ | $\mathcal{O}(\alpha_s/a)$ | $\mathcal{O}(\alpha_s)$ | N/A |
| Cost | 1 | $\sim 1/4$ | ~ 1.1 | ~ 1.1 | $\sim 10-100$ |



CLQCD ensembles







CLQCD choice and informations


- Features:
 - Maximum lattice size $48^3 \times 144$,
 - Clover fermion action with stout smearing,
 - Similar pion mass and volume at different lattice spacing:
- Cost:
 - That of an independent configuration (per 10 traj.'s with $\tau = 1.0$, converted to A100 GPU hours) is shown on the figure;
 - Needs $\sim 1,000$ configurations per ensemble;
 - Currently used 658k A100 hours, equals to 3.3M Chinese Yuan with the market price.
- Working on the Sugon machines to avoid the embargo of A100 GPU.

CLQCD ensembles


Published/accepted works with the CLQCD ensembles




Q.A. Zhang, et.al.,
Chin.Phys.C 46 (2022) 011002



D.J. Zhao, et.al., χ QCD,
Phys.Rev.D 107 (2023) L091501




Q.A. Zhang, et.al.,
Phys.Lett.B 841 (2023) 137941



H. Liu, et.al.,
2207.00183, accepted by SCPMA

CLQCD ensembles




Nucleon mass v.s. pion mass

- At $a = 0.105(3)$ fm, we have

$$m_\pi = 135.5(1.6) \text{ MeV},$$


$$m_N = 890(10) \text{ MeV}.$$

- m_N are $\sim 5\%$ smaller than the physical value, and can reach the physical value after the continuum extrapolation.



Background

Chiral symmetry breaking in the clover fermion



Z.C. Hu, B.L. Hu, J.H. Wang, et. al., CLQCD, 2310.00814

- Due to the additive α_s/a correction, the dimensionless bare quark mass $\tilde{m}_q^b = m_q^b a$ is negative.
- The renormalized quark mass should be defined as $m_q^R = Z_m(m_q^b - m_{\text{crti}})$, where m_{crti} is defined as the m_q^b which vanishes the pion mass.
- One can avoid this difficulty by defining the quark mass through PCAC relation:



$$\langle 0 | \partial_4 A_4 | \text{PS} \rangle = (m_q^{\text{PC}} + m_{\bar{q}}^{\text{PC}}) \langle 0 | P | \text{PS} \rangle$$

T. Ishikawa, et.al., JLQCD, Phys.Rev.D78 (2008) 011502

- And then m_q^{PC} is always positive and can be renormalized as $m_q^R = Z_P/Z_A m_q^{\text{PC}}$.

Renormalization and final results

Renormalized quark mass



- Non-perturbative renormalization to $\overline{\text{MS}}$ 2 GeV eliminates the regularization scale $1/a$ dependence of m_π^2/m_q .
- m_π^2/m_q using the clover fermion also turns out to be consistent with that using the overlap fermion.
- The large uncertainty of the renormalized m_π^2/m_q majorly comes from the missing higher order effect of the perturbative matching

$$\frac{Z_P^{\overline{\text{MS}}}}{Z_P^{\text{MOM}}} = 1 + 0.4244\alpha_s + 1.007\alpha_s^2 + 2.722\alpha_s^3 + 8.263\alpha_s^4 + \mathcal{O}(\alpha_s^5)$$



$$= \frac{1 - 2.611\alpha_s - 0.2813\alpha_s^2 - 0.3349\alpha_s^3}{1 - 3.036\alpha_s} + \mathcal{O}(\alpha_s^5),$$

J.A. Gracey, Eur.Phys.J.C83 (2023) 181

- and can be highly suppressed after the continuum extrapolation.

Chiral symmetry breaking and renormalization

Restore of chiral symmetry in the continuum



- Renormalized quark mass $m_q^R = Z_A/Z_P m_q^{\text{PC}}$ with 317 MeV pion mass at three lattice spacings:
- The intermediate renormalization scheme dependence is 3.1(1.5)%.
- RI/MOM scheme has smaller discretization error.
- Feynman-Hellman theorem can extract $g_{S,\pi}$ as



$$g_{S,\pi}^{\text{FH}} = \frac{1}{2} \frac{\partial m_\pi(m_q)}{\partial m_q} \simeq \frac{Z_P}{Z_A} \frac{m_\pi}{4m_q^{\text{PC}}} + \mathcal{O}(m_q, a^2)$$

which is 4.04(6)(12) for $m_\pi = 317$ MeV in the continuum.

- Renormalized $g_{S,\pi}^{\text{R,ME}} = Z_S \frac{\langle \pi | S | \pi \rangle_{\text{conn}}}{\langle \pi | \pi \rangle}$ based on the direct calculation:
- The intermediate renormalization scheme dependence is 7.6(2.3)% (linear a^2 correction) or 2.0(5.8)% ($a^2 + a^4$ corrections).
- $g_{S,\pi}^{\text{ME}}$ using RI/MOM scheme has smaller discretization error, and agree with $g_{S,\pi}^{\text{R,FH}}$ within 2σ at all the lattice spacings.

Chiral symmetry breaking and renormalization

Global fit of the pion mass







- Present CLQCD prediction of the u-d averaged light quark masses is consistent with the lattice averages within 5% uncertainty.
- Most of the uncertainties come from the non-perturbative renormalization and further improvements are in progress.
- All the finite volume, discretization and sea quark mass effects have been taken into account.

$$m_{\pi, \text{vv}}^2 = \Lambda_\chi^2 2y_v \left\{ 1 + \frac{2}{N_f} [(2y_v - y_s) \ln(2y_v) + (y_v - y_s)] + 2y_v(2\alpha_8 - \alpha_5) + 2y_s N_f(2\alpha_6 - \alpha_4) \right\} \times [1 + c_L^\pi e^{-m_\pi L} + c_s^\pi (m_{\eta_s}^2 - m_{\eta_s, \text{phys}}^2)] \times (1 + d_a^\pi a^2),$$

$$\Lambda_\chi = 4\pi F, \quad y = \frac{\Sigma m}{F^2 \Lambda_\chi^2}$$

Renormalization and final results

Global fit of the low energy constants




$$m_{\pi,vv}^2 = \Lambda_\chi^2 2y_v \left\{ 1 + \frac{2}{N_f} [(2y_v - y_s) \ln(2y_v) + (y_v - y_s)] \right. \\ \left. + 2y_v(2\alpha_8 - \alpha_5) + 2y_s N_f (2\alpha_6 - \alpha_4) \right\} \\ \times [1 + c_L^\pi e^{-m_\pi L} + c_s^\pi (m_{\eta_s}^2 - m_{\eta_s, \text{phys}}^2)] \\ \times (1 + d_a^\pi a^2), \quad \Lambda_\chi = 4\pi F, y = \frac{\Sigma m}{F^2 \Lambda_\chi^2}$$

$$F_{\pi,vv} = F \left(1 - \frac{N_f}{2} (y_v + y_s) \ln(y_v + y_s) + y_v \alpha_5 + y_s N_f \alpha_4 \right) \\ \times [1 + d_L^\pi e^{-m_\pi L} + d_s^\pi (m_{\eta_s}^2 - m_{\eta_s, \text{phys}}^2)] \\ \times (1 + d_a^\pi a^2)$$

- Global fit of all the ensembles to obtain the quark mass dependence of m_π and f_π in the continuum and infinite volume limit, which allows us to extract the χ PT low energy constants.

Chiral symmetry breaking and renormalization

Global fit of the kaon mass



$$m_K^2(m_l^v, m_l^s, m_s^v, m_s^s, a) = (b_s^v m_s^v + b_s^s m_s^s + b_l^v m_l^v + b_l^s m_l^s) \times [1 + c_l^K m_l^v + c_m^K a^2 + c_L^K \exp(-m_K L)],$$

$$m_u^{\text{phys}} + m_d^{\text{phys}} = 2m_l^{\text{phys}}.$$

$$m_K(m_d^{\text{phys}}, m_l^{\text{phys}}, m_s^{\text{phys}}, m_s^{\text{phys}}, 0) = m_{K^0, \text{QCD}},$$

$$m_K(m_u^{\text{phys}}, m_l^{\text{phys}}, m_s^{\text{phys}}, m_s^{\text{phys}}, 0) = m_{K^\pm, \text{QCD}},$$

val sea val sea

P.Zyla et,al, PTEP(2020)083C01 (PDG2020):

- $m_p = 938.27 \text{ MeV} = m_{p, \text{QCD}} + 1.00(16) \text{ MeV} + \dots;$

- $m_n = 939.57 \text{ MeV};$

- $m_\pi^0 = 134.98 \text{ MeV};$


- $m_\pi^+ = 139.57 \text{ MeV} = m_\pi^0 + 4.53(6) \text{ MeV} + \dots;$

- $m_K^0 = 497.61(1) \text{ MeV} = m_{K, \text{QCD}}^0 + 0.17(02) \text{ MeV} + \dots;$

- $m_K^+ = 493.68(2) \text{ MeV} = m_{K, \text{QCD}}^+ + 2.24(15) \text{ MeV} + \dots$

Renormalization and final results

Quark mass of three light flavors




Summary of the lattice methodology.—The lattice setup used for this project is very similar to Ref. [13] and is based on our set of lattice QCD simulations presented in Ref. [6]. It is composed of 47 $N_f = 2 + 1$ QCD ensembles with pion masses down to 120 MeV, 5 lattice spacings down to 0.054 fm, and 16 different volumes up to $(6 \text{ fm})^3$.

BMWc, PRL 117(2016)0820001

Z.C. Hu, B.L. Hu, J.H. Wang, et. al., CLQCD, 2310.00814

Explore heavy flavors





Discretization error

- The heavy quark with mass m_Q will have additional $\mathcal{O}(m_Q^n a^n)$ discretization errors.
- And the one would need $m_b^{\text{bare}} a \sim 9$ to reach the physical bottom quark mass at $a = 0.105 \text{ fm}$.
- The situation can be significantly improved when the lattice spacing becomes smaller.

Explore heavy flavors

Charmonium and Bottomium



H.Y. Du, et. al., CLQCD, in preparation

- At $a = 0.105$ fm, the discretization error can make the 1P-1S fine splitting to be $\sim 10\%$ larger, and the hyperfine splitting to be 50% (charm) and 95% (bottom) smaller.
- The situation can also be significantly improved when the lattice spacing becomes smaller.

Explore heavy flavors

Another choice: anisotropic lattices

2 味轻海夸克


| Size | β | a_s/fm | ξ_g | ξ_f | m_π/MeV | Traj. # |
|-------------------|---------|-----------------|-----------|-----------|--------------------|---------|
| $16^3 \times 128$ | 2.0 | 0.1361(18) | 5.036(10) | 4.978(16) | 416.0(5.7) | 6000 |
| $16^3 \times 128$ | 2.0 | 0.1361(18) | 5.060(11) | 4.956(22) | 359.3(5.0) | 6000 |
| $16^3 \times 128$ | 2.0 | 0.1361(18) | 5.022(10) | 4.956(25) | 304.7(4.6) | 6000 |
| $24^3 \times 192$ | 2.0 | 0.1361(18) | 5.044(5) | 5.043(18) | 305.4(4.2) | 4020 |
| $16^3 \times 128$ | 2.0 | 0.1361(18) | 5.050(10) | 5.017(42) | 245.7(4.6) | 6000 |
| $24^3 \times 192$ | 2.0 | 0.1361(18) | 5.014(8) | 4.966(60) | 237.9(4.8) | 2525 |
| $24^3 \times 192$ | 2.2 | ≈ 0.108 | | | | 调测中 |

| Size | β | a_s/fm | ξ_g | ξ_f | m_π/MeV | Traj. # |
|-------------------|---------|-----------------|---------|---------|--------------------|---------|
| $16^3 \times 128$ | 2.0 | 0.1517 | 4.95 | 5.30 | 349.2 | 70045 |
| $24^3 \times 192$ | 2.0 | 0.1517 | 4.95 | 5.30 | 349.2 | 28000 |

1 味 strange 海夸克

| Size | β | a_s/fm | ξ_g | ξ_f | m_{ps}/MeV | Traj. # |
|-------------------|---------|-----------------|---------|---------|---------------------|---------|
| $16^3 \times 128$ | 2.4 | 0.148 | 5.0 | 5.0 | 686 (strange) | 18000 |

- It is much more efficient to reach small lattice space in the time direction only, with the anisotropic lattices.
- While obtain the renormalized quark mass is more challenging.



Summary

- We chose the clover fermion and Symanzik gauge actions to generate the Lattice QCD ensembles at multiple lattice spacing, pion mass and volume, and figured out the proper renormalization to restore the chiral symmetry at 5% level.
- Current prediction of quark masses and low energy constants agree with the lattice averages within 5%, and more accurate studies are on-going.
- The heavy quark can have huge discretization error at coarse lattice spacings, while further improvement on the fermion action is also on-going.

