

Lattice QCD study of the hidden-charm pentaquark states





Liuming Liu
IMP, CAS



The 7th Symposium on “Symmetries and the emergence of structure in QCD

Shandong, July 19-22,2023



$P_c(4380)$ \rightarrow $P_c(4312)$
 $P_c(4450)$ \rightarrow $P_c(4440)$
 $P_c(4450)$ \rightarrow $P_c(4457)$

R. Aaij et al. (LHCb), Phys. Rev. Lett. 115, 072001 (2015)

R. Aaij et al. (LHCb), Phys. Rev. Lett. 122, 222001 (2019)

Theory interpretations:




- Molecule bound states
- Compact pentaquark states
-

$\Sigma_c^{(*)} \bar{D}^{(*)}$ molecules:

$$\Sigma_c \bar{D}, J^P = \frac{1}{2}^-, P_c(4312)$$

$$\Sigma_c \bar{D}^*, J^P = \left(\frac{1}{2}^-, \frac{3}{2}^- \right), P_c(4440)/P_c(4457)$$

$$\Sigma_c^* \bar{D}, J^P = \frac{3}{2}^-, \quad \Sigma_c^* \bar{D}^*, J^P = \left(\frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^- \right),$$



R. Aaij et al. (LHCb), Phys. Rev. Lett. 115, 072001 (2015)

R. Aaij et al. (LHCb), Phys. Rev. Lett. 122, 222001 (2019)

Theory interpretations:

- Molecule bound states
- Compact pentaquark states
-


$\Sigma_c^{(*)} \bar{D}^{(*)}$ molecules:

$$\Sigma_c \bar{D}, J^P = \left(\frac{1}{2}^-, P_c(4312) \right)$$

$$\Sigma_c \bar{D}^*, J^P = \left(\frac{1}{2}^-, \frac{3}{2}^- \right), P_c(4440)/P_c(4457)$$

$$\Sigma_c^* \bar{D}, J^P = \frac{3}{2}^-, \quad \Sigma_c^* \bar{D}^*, J^P = \left(\frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^- \right),$$

Lüscher's finite volume method: M. Lüscher, Nucl. Phys. B354, 531(1991)



- ◆ General Lüscher's formula for two-body scattering:

$$\det[\mathbf{1} + i\rho \cdot \mathbf{t} \cdot (1 + i\mathbf{M})] = 0$$

Diagonal matrix of
phase-space factors

$$\rho_{ij} = \delta_{ij} \frac{2k_i}{E_{cm}}$$

Infinite-volume
scattering matrix

Finite volume
information

$$M(E_{cm}, L)$$

- ◆ Resonances/bound states are formally defined as poles in scattering amplitudes.



Scattering on lattice



Finite volume spectrum:

- ◆ build large basis of operators $\{\mathcal{O}_1, \mathcal{O}_2, \dots\}$ with desired quantum numbers, construct the matrix of correlation function:

$$C_{ij} = \langle 0 | \mathcal{O}_i \mathcal{O}_j^\dagger | 0 \rangle = \sum_n Z_i^n Z_j^{n*} e^{-E_n t}$$

- ◆ Solve the generalized eigenvalue problem(GEVP):

$$C_{ij} v_j^n(t) = \lambda_n(t) C_{ij}^0 v_j^n(t)$$

- ◆ Eigenvalues: $\lambda_n(t) \sim e^{-E_n t} (1 + e^{-\Delta E t})$

- ◆ Optimal linear combinations of the operators to overlap on the n'th state:

$$\Omega_n = \sum_i v_i^n \mathcal{O}_i$$



Lattice QCD configurations



Lattice spacing	Volume($L^3 \times T$)	M_π (MeV)	# of confs
~0.108fm	$24^3 \times 72$	290	1000
	$32^3 \times 64$	290	1000
	$32^3 \times 64$	220	450
	$48^3 \times 96$	220	200
	$48^3 \times 96$	140	200
~0.080fm	$32^3 \times 96$	300	480
	$48^3 \times 96$	300	200
	$32^3 \times 64$	220	460
	$48^3 \times 96$	220	200
~0.055fm	$48^3 \times 144$	300	200




L. Liu, M. Gong, W. Sun,
P. Sun, W. Wang, Y.B. Yang



Lattice QCD configurations



Lattice spacing	Volume($L^3 \times T$)	M_π (MeV)	# of confs
~0.108fm	$24^3 \times 72$	290	1000
	$32^3 \times 64$	290	1000
	$32^3 \times 64$	220	450
	$48^3 \times 96$	220	200
	$48^3 \times 96$	140	200
~0.080fm	$32^3 \times 96$	300	480
	$48^3 \times 96$	300	200
	$32^3 \times 64$	220	460
	$48^3 \times 96$	220	200
~0.055fm	$48^3 \times 144$	300	200



L. Liu, M. Gong, W. Sun,
P. Sun, W. Wang, Y.B. Yang

$\Sigma_c \bar{D}$ and $\Sigma_c \bar{D}^*$ scattering ($J^P = \frac{1}{2}^-$):

◆ Five operators:


$$\mathcal{O}_1 = \Sigma_c(\mathbf{p}) \bar{D}(-\mathbf{p}) \quad (|\mathbf{p}| = 0)$$

$$\mathcal{O}_2 = \Sigma_c(\mathbf{p}) \bar{D}(-\mathbf{p}) \quad (|\mathbf{p}| = 1)$$

$$\mathcal{O}_3 = \Sigma_c(\mathbf{p}) \bar{D}(-\mathbf{p}) \quad (|\mathbf{p}| = \sqrt{2})$$

$$\mathcal{O}_4 = \Sigma_c(\mathbf{p}) \bar{D}^*(-\mathbf{p}) \quad (|\mathbf{p}| = 0)$$

$$\mathcal{O}_5 = \Sigma_c(\mathbf{p}) \bar{D}^*(-\mathbf{p}) \quad (|\mathbf{p}| = 1)$$



◆ The finite-volume energies lie below the free energies, indicating rather strong attractive interactions.

Scattering amplitude:

$$T \sim \frac{1}{p \cot \delta - ip}$$

Effective range expansion:

$$p \cot \delta(p) = \frac{1}{a_0} + \frac{1}{2} r_0 p^2 + \dots$$

Luscher's formula:



$$p \cot \delta(p) = \frac{2Z_{00}(1; (\frac{pL}{2\pi})^2)}{L\sqrt{\pi}}$$

Bound state pole:

$$p = i|p_B|$$

$$\begin{aligned} \Sigma_c \bar{D} : P_c(4312) ? \\ a_0 = -2.0(3)(5) \text{ fm} \\ E_B = 6(2)(2) \text{ MeV} \end{aligned}$$

$$\begin{aligned} \Sigma_c \bar{D}^* : P_c(4440)/P_c(4457) ? \\ a_0 = -2.3(5)(1) \text{ fm} \\ E_B = 7(3)(1) \text{ MeV} \end{aligned}$$






Coupled channels



Coupled channels: $\eta_c N, J/\psi N, \Lambda_c \bar{D}, \Lambda_c \bar{D}^*, \Sigma_c \bar{D}, \Sigma_c \bar{D}^*$

Non-interacting energy levels:

$$E_{free} = \sqrt{m_1^2 + p_1^2} + \sqrt{m_2^2 + p_2^2}$$





Coupled channels



Coupled channels: $\eta_c N, J/\psi N, \Lambda_c \bar{D}, \Lambda_c \bar{D}^*, \Sigma_c \bar{D}, \Sigma_c \bar{D}^*$

◆ 15 operators for the $L = 32$ ensemble:

$$\mathcal{O}_{1,2,3} = N(\mathbf{p})\eta_c(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2)$$

$$\mathcal{O}_{4,5} = N(\mathbf{p})J/\psi(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1)$$

$$\mathcal{O}_{6,7,8} = \Lambda_c(\mathbf{p})\bar{D}(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2)$$

$$\mathcal{O}_{9,10} = \Lambda_c(\mathbf{p})\bar{D}^*(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1)$$

$$\mathcal{O}_{11,12,13} = \Sigma_c(\mathbf{p})\bar{D}(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2)$$

$$\mathcal{O}_{14,15} = \Sigma_c(\mathbf{p})\bar{D}^*(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1)$$

◆ 23 operators for the $L = 48$ ensemble:

$$\mathcal{O}_{1,2,3,4,5} = N(\mathbf{p})\eta_c(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2,3,4)$$

$$\mathcal{O}_{7,8,9,10} = N(\mathbf{p})J/\psi(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2,3)$$

$$\mathcal{O}_{10,11,12,13,14} = \Lambda_c(\mathbf{p})\bar{D}(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2,3,4)$$

$$\mathcal{O}_{15,16,17,18} = \Lambda_c(\mathbf{p})\bar{D}^*(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2,3)$$



$$\mathcal{O}_{19,20,21} = \Sigma_c(\mathbf{p})\bar{D}(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1,2)$$

$$\mathcal{O}_{22,23} = \Sigma_c(\mathbf{p})\bar{D}^*(-\mathbf{p}) \quad (\mathbf{p}^2 = 0,1)$$

Coupled channels

$L = 32$

$L = 48$






Coupled channels




$N - \eta_c(J/\psi)$ spectrum:





Coupled channels





Summary



- ◆ Single channel analysis indicates bound states in $\Sigma_c \bar{D}$ and $\Sigma_c \bar{D}^*$ channel at $m_\pi \sim 300\text{MeV}$. Coupled channel analysis is ongoing. Pion mass dependence will also be explored.