

enton Search at

The 2023 international workshop on the high energy CPEC Oct 23-27, 2023. Nanjing University



In collaboration with Jin–Min Yang, Yang Zhang & Rui Zhu. Based on PRD. 108 (2023) 7, 075015 [arXiv: 2211.01832]



Pengxuan Zhu(朱鹏轩)

Institute of Theoretical Physics, Chinese Academy of Science Oct-26, 2023



CEPC-2023 Collider Physics



Oct-26, 2023. Nanjing U.

Hard Scattering Level: (Master Formulae)

$$\sigma = \sum_{a,b} \int_0^1 dx_a dx_b \int f_a^{h1}(x_a, \mu_F) f_b^{h2}(x_b, \mu_R) \, d\sigma_{a,b}(\mu_F, \mu_R)$$

- Parton distribution function
- Matrix element
- Initial state radiation \bullet

Parton Level

- Parton shower
- Decay
- Final state radiation

Detector Level

- Object reconstruction
- Object Identification
- Electron, muon, jets, photon, tracks

Event Reconstruction Level

- Event number / cross section: σ
- missing energy: p_{miss} \bullet
- Kinematic variable distributions: $\frac{d\sigma}{d\mathcal{O}}$



Lepton Collider

Advantages

Conservation laws benefits a lot.

Collision system

- O net charge & O lepton number before and after.
 - 1. Suitable to create new particles after e^+e^- annihilation
- Symmetric beams of e^+e^- .
 - 2. CM frame \approx Lab frame;
 - 3. Total $E_{\rm CM}$ fully exploited to reach highest possible physics threshold.
- Well-understood beam properties.
 - 4. PDFs are simple, $E_{\rm CM} = \sqrt{s}$.
 - 5. ISR from electroweak process.

Detector Level

- Low QCD soft background.
 - 1. High object identification & reconstruction efficiency.
- High precision four-momentum vector.
 - 2. Recoil system measurement method.
 - Lorentz invariant kinematic observables. 3.

Oct-26, 2023. Nanjing U.

Higgs mass measurement by recoiled Z



Missing energy and momentum



P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV



Reconstructing events containing two invisible particle

$e^+e^- \rightarrow \mathbf{P}_a\mathbf{P}_b \rightarrow \mathbf{V} \mathbf{I} \mathbf{V} \mathbf{I}$ **Lepton Collider**



- Investigating the invisible decayed \mathbf{P}_{b} can be by recoiled system \mathbf{P}_{a} .
- Two Lorentz invariant observable:
 - Invariant mass: $m_{V_{a1}V_{a2}}$
 - Recoil mass: *m*_{recoil}

Oct-26, 2023. Nanjing U.



• What for semi-invisible decayed topology?

Almost all new physics theories with dark matter candidates I also contain additional particles **P** that are charged not only under newsymmetry but also carry SM charges. At collider, these new particles must be pair produced, and will (cascade) decay into dark matter particle plus one or a group of SM visible particles.



Slepton searches at LHC



- LHC Run-II with 139 fb⁻¹ have already set constraints on di-lepton+ $E_{\rm T}^{\rm miss}$ final states in searches for slepton pair production.
- Compressed spectrum region can be probed by events with ISR via Recursive Jigsaw Reconstruction(RJR) technique.
- Still a gap uncovered for $m_{\tilde{e}}$ at $\mathcal{O}(100)$ GeV.
- CEPC@240 can closely reach to 120 GeV, corresponding to a production cross section around 10 fb.

Oct-26, 2023. Nanjing U.





P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV



Smuon pair production at lepton collider

arXiv: 2104.06421 provide an idea of searching $\tilde{\mu}$ via pair production process at CEPC

$$e^+e^- \rightarrow \tilde{\mu}\tilde{\mu}^{(*)} \rightarrow \mu^+\tilde{\chi}^0_1\mu^-\tilde{\chi}^0_1$$

with one $\tilde{\mu}$ being off-shell, i.e. $m_{\tilde{\mu}} > \sqrt{s/2}$.

• On-shell region: $\sqrt{s/2} - m_{\tilde{\mu}} \gg \Gamma_{\tilde{\mu}}$

narrow width approximation when

$$\sigma(e^+e^- \to \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0) \approx \sigma(e^+e^- \to \tilde{\mu}\tilde{\mu}) \times \left(\mathrm{BR}(\tilde{\mu} \to \mu\tilde{\chi}_1^0)\right)^2$$

• Off-shell region:

$$e^+e^- \to \tilde{\mu}^{\pm} \mu^{\mp} \tilde{\chi}_1^0$$

The cross section of this $2 \rightarrow 3$ process is smaller than that of On-shell region by a factor of $16\pi^2$.

Assuming BR($\tilde{\mu} \rightarrow \mu \tilde{\chi}_1^0$) = 1, for 5.05 ab⁻¹ data, about 800 signal events can be produced of mass point $(m_{\tilde{\mu}}, m_{\tilde{\chi}_1^0}) = (140, 0)$ GeV.

- $(m_{\tilde{\mu}}, m_{\tilde{\chi}_1^0}) = (115, 0)$ GeV: 23.6 fb
- $(m_{\tilde{\mu}}, m_{\tilde{\chi}_1^0}) = (120, 0) \text{ GeV:} 0.41 \text{ fb}$
- $(m_{\tilde{\mu}}, m_{\tilde{\chi}_1^0}) = (140, 0)$ GeV: 0.013 fb

J. Liu, X. P. Wang and K. P. Xie. arXiv: 2104.06421 [hep-ph]

Oct-26, 2023. Nanjing U.



P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV





10⁻³



Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.



Reconstructing events containing two invisible particle

Kinematic reconstruction @ Lepton Collider



Oct-26, 2023. Nanjing U.

 $e^+e^- \rightarrow \mathbf{P}_a\mathbf{P}_b, \mathbf{P} \rightarrow \mathbf{VI}$

What can be reconstructed by detector are:

- $p_{V_z}^{\mu,\text{Lab}}$: four-momentum of the visible particle (group) a in Lab frame;
- $p_{V_{L}}^{\mu,\text{Lab}}$: four-momentum of b;
- $p_{\text{miss}}^{\mu,\text{Lab}} = p_{\text{I}_a}^{\mu} + p_{\text{I}_b}^{\mu}$: Vector sum of four-momentum of two invisible particles

$$p_{\text{miss}}^{\mu,\text{Lab}} = (\sqrt{s}, \vec{0}) - p_{V_a}^{\mu,\text{Lab}} - p_{V_b}^{\mu,\text{Lab}} - p_{\text{R}}^{\mu,\text{Lab}}$$

• $p_{\rm R}^{\mu,{\rm Lab}}$: four-momentum of anything else, including the ISR, FSR, imperfect detector performance, etc.











Reconstructing events containing two invisible particle

Physical Constraints (Assumptions)

1. $p_{P_a}^{\mu} = p_{V_a}^{\mu} + p_{I_a}^{\mu};$ $2.\,p_{\rm P_{\rm b}}^{\,\mu} = p_{V_{\rm b}}^{\,\mu} + p_{\rm I_{\rm b}}^{\,\mu};$ $3. m_{\rm P} = m_{\rm P_a} = m_{\rm P_b};$ 4. $m_{\rm I} = m_{\rm I_a} = m_{\rm I_b};$ 5. $m_{\rm P} > 0;$ 6. $m_{\rm I} \ge 0$.



Oct-26, 2023. Nanjing U.

triangle, and three vectors $\vec{p}_{\rm L}$, $\vec{p}_{\rm L}$ and $\vec{p}_{\rm miss}$ form another invisible triangle.

The mass information are not changed by the Lorentz boost.



Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.

In PP frame,

$$p_{\mathbf{I}_{a}}^{\mu,\mathrm{PP}} = \left\{ \frac{1}{2} \left(-E_{\mathbf{V}_{a}}^{\mathrm{PP}} + E_{\mathbf{V}_{b}}^{\mathrm{PP}} + E_{\mathrm{miss}}^{\mathrm{PP}} \right), \vec{p}_{\mathbf{I}_{a}}^{\mathrm{PP}} \right\},\$$

$$p_{\mathbf{P}_{a}}^{\mu,\mathrm{PP}} = \left\{ \frac{1}{2} \left(E_{\mathbf{V}_{a}}^{\mathrm{PP}} + E_{\mathbf{V}_{b}}^{\mathrm{PP}} + E_{\mathrm{miss}}^{\mathrm{PP}} \right), \vec{p}_{\mathbf{V}_{a}}^{\mathrm{PP}} + \vec{p}_{\mathbf{I}_{a}}^{\mathrm{PP}} \right\},\$$

$$p_{\mathbf{I}_{b}}^{\mu,\mathrm{PP}} = \left\{ \frac{1}{2} \left(E_{\mathbf{V}_{a}}^{\mathrm{PP}} - E_{\mathbf{V}_{b}}^{\mathrm{PP}} + E_{\mathrm{miss}}^{\mathrm{PP}} \right), \vec{p}_{\mathrm{miss}}^{\mathrm{PP}} - \vec{p}_{\mathbf{I}_{a}}^{\mathrm{PP}} \right\},\$$

$$p_{\mathbf{P}_{b}}^{\mu,\mathrm{PP}} = \left\{ \frac{1}{2} \left(E_{\mathbf{V}_{a}}^{\mathrm{PP}} + E_{\mathbf{V}_{b}}^{\mathrm{PP}} + E_{\mathrm{miss}}^{\mathrm{PP}} \right), \vec{p}_{\mathbf{V}_{b}}^{\mathrm{PP}} + \vec{p}_{\mathrm{miss}}^{\mathrm{PP}} - \vec{p}_{\mathbf{I}_{a}}^{\mathrm{PP}} \right\}.$$

•
$$p_{\mathbf{P}_b}^{\mu,\mathrm{PP}} = \left\{ E_{\mathbf{P}_a^{\mathrm{PP}}}, -\vec{p}_{\mathbf{P}_a}^{\mathrm{PP}} \right\}$$
, so $m_{\mathbf{P}_a} = m_{\mathbf{P}_b}$.

- At the event level, the real mass of the particle follows the Breit-Wigner distribution. $m_{\mathbf{P}_a} = m_{\mathbf{P}_b}$ is an assumption which is reasonable in the narrow width.
- The only unknown momentum is $\vec{p}_{\mathbf{I}_a}^{\mathrm{PP}}$, or the location of vertex I



Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.

• $m_{\mathbf{I}_a} \ge 0$: The magnitude of $\vec{p}_{\mathbf{I}_a}^{\text{PP}}$ must be smaller than

$$E_{\mathbf{I}_a} = \left(-E_{\mathbf{V}_a}^{\mathrm{PP}} + E_{\mathbf{V}_b}^{\mathrm{PP}} + E_{\mathrm{miss}}^{\mathrm{PP}} \right) / 2.$$

Therefore, the point I is located in a sphere, where the spherical shell implies $m_{\mathbf{I}_a} = 0.$

- $m_{\mathbf{I}_{h}} \ge 0$: Similar to the constraint $m_{\mathbf{I}_{a}} \ge 0$.
- $m_{\mathbf{I}_a} = m_{\mathbf{I}_b} = m_{\mathbf{I}}$: The point **I** is located in a plane vertical to $\vec{p}_{\text{miss}}^{\text{PP}}$ (the light blue plane).

$$\vec{p}_{\mathbf{I}_a}^{\mathrm{PP}} \cdot \vec{p}_{\mathrm{miss}}^{\mathrm{PP}} - D = 0,$$

$$D = \frac{1}{2} \left(E_{\mathbf{V}_a}^{\mathrm{PP}^2} - E_{\mathbf{V}_b}^{\mathrm{PP}^2} + E_{\mathbf{PP}}^{\mathrm{PP}} (E_{\mathbf{V}_b}^{\mathrm{PP}} - E_{\mathbf{V}_a}^{\mathrm{PP}}) + \left| \vec{p}_{\mathrm{miss}}^{\mathrm{PP}} \right|^2 \right)$$





Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.

What unknown is the location of the pink dot, but it must satisfy the physical assumptions, i.e. it must located in the yellow round plane.

> There are two independent degree to determine the location of the pink dot:

- ϕ : the angle between visible vector plane and invisible plane.
- *R*: the distance between the pink dot and the center of yellow round plane.

The boundary means $m_{\rm I} = 0$ GeV.



Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.



Choose a pink dot location, we can solve a set of (m_P, m_I) :

• For the point in the circle of fixed R (pink circle), the $m_{\rm I}$ is the same.

• For the point in the purple circle (fixed r), the m_P is the same.

So the geometric property of the disk (physical region) implies the kinetic information of the semi-invisibly decayed events, which can be represented as three "reconstructed-mass" variables:

 $m_{\rm RC}^{\rm min}$: the $m_{\rm P}$ by choosing pink dot located in point A.

 $m_{\rm RC}^{\rm max}$: the $m_{\rm P}$ by choosing pink dot located in point B

(or point C, if C is inside the disc).

 $m_{\text{LSP}}^{\text{max}}$: the m_{I} reconstructed by choosing pink dot located in center point O. It is easy to say that:

$$0 < m_{\rm RC}^{\rm min} \le m_{\rm P}^{\rm true} \le m_{\rm RC}^{\rm max} < \sqrt{s} / 2$$

$$0 \le m_{\rm I}^{\rm ture} \le m_{\rm LSP}^{\rm max}$$

P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV

Reconstructing events containing two invisible particle



Oct-26, 2023. Nanjing U.

The problem of "how massive the intermediate particle is" is transformed into a plane geometric problem:

$$\begin{split} &= \sqrt{\frac{E_{\rm PP}^2}{4} - |PC|^2 - \left(|OC| + |OA|\right)^2}, \\ &= \sqrt{\frac{E_{\rm PP}^2}{4} - |PC|^2 - \left(|OC| - |OB|\right)^2}, \text{ if } |OC| > |OB|, \\ &= \sqrt{\frac{E_{\rm PP}^2}{4} - |PC|^2}, \qquad \text{ if } |OC| \le |OB|; \\ &= \sqrt{\frac{E_{\rm PP}^2}{4} - |PC|^2}, \qquad \text{ if } |OC| \le |OB|; \\ &= \sqrt{\frac{E_{\rm PP}^2}{4} - \frac{1}{4\left|\vec{p}_{\rm miss}\right|^2} \left(\left|\vec{p}_{\rm miss}\right|^2 + E_{\rm I_a}^2 - E_{\rm I_b}^2\right)^2}. \\ &= \frac{1}{2\left|\vec{p}_{\rm miss}\right|} \left(-E_{\rm I_a}^2 + E_{\rm I_b}^2 + \left|\vec{p}_{\rm V_a}\right|^2 - \left|\vec{p}_{\rm V_b}\right|^2\right), \\ &= \frac{1}{2\left|\vec{p}_{\rm miss}\right|} \left[-\left|\vec{p}_{\rm miss}\right|^4 - \left(\left|\vec{p}_{\rm V_a}\right|^2 - \left|\vec{p}_{\rm V_b}\right|^2\right)^2 + 2\left|\vec{p}_{\rm miss}\right|^2 \left(\left|\vec{p}_{\rm V_a}\right|^2 + \left|\vec{p}_{\rm V_b}\right|^2\right)\right)^{\frac{1}{2}}, \\ &= \frac{1}{2\left|\vec{p}_{\rm miss}\right|} \left[-E_{\rm I_a}^4 - \left(E_{\rm I_b}^2 - \left|\vec{p}_{\rm miss}\right|^2\right)^2 + 2E_{\rm I_a}^2 \left(E_{\rm I_b}^2 + \left|\vec{p}_{\rm miss}\right|^2\right)^{\frac{1}{2}}. \end{split}$$

P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV

14





Reconstructed mass variables: m_{RC}^{min} vs m_{RC}^{max}





- Both distributions of $m_{\rm RC}^{\rm min}$ and $m_{\rm RC}^{\rm max}$ have the "end-point"s and the peaks at *m***P**.
- The tails of $m_{\rm RC}^{\rm min}$ and $m_{\rm RC}^{\rm max}$ (tail region) are caused by that the muon (bare-level muons used in the calculation) in final state losing energy through photon radiation.
- The distribution of WW, ZZ and $\tau\tau$ events in same final state $\mu^+\mu^- + E_{\text{miss}}$ are much different.
- $\tau\tau$ events' shape are also show the τ mass.

Oct-26, 2023. Nanjing U.

- The calculation is decay topology independent. For example, Z boson in ZZ events is not semi-invisibly decayed.
- A linear relation between $m_{\rm RC}^{\rm min}$ and $m_{\rm RC}^{\rm max}$ of ZZ event.





Reconstructed mass variables: m_{LSP}^{max} (compared to m_{Recoil})





•
$$m_{\text{LSP}}^{\text{max}} \leqslant \frac{1}{2} m_{\text{Recoil}}$$
.

- For WW events, the shape of $m_{\text{LSP}}^{\text{max}}$ is much similar to that of m_{Recoil} .
- For $\tau\tau$ events, $m_{\text{LSP}}^{\text{max}}$ distributed in the small value region. But m_{Recoil} distributed more flat.
- For semi-invisible decay topology event, $m_{\text{LSP}}^{\text{max}}$ is better than m_{Recoil} .

Oct-26, 2023. Nanjing U.



• For invisibly recoiled events, like ZZ events, the recoil mass is more physical.





Reconstructed mass variables: Smuon pair production



- For massless LSP, both $m_{\rm RC}^{\rm min}$ and $m_{\rm RC}^{\rm max}$ have significant peaks at $m_{\tilde{\mu}}$ and also an obvious truncation at $m_{\tilde{\mu}}$.
- For massive LSP, the truncations of $m_{\text{RC}}^{\text{min}}$ and $m_{\text{RC}}^{\text{max}}$ are disappearing, and the range $[m_{\text{RC}}^{\text{min}}, m_{\text{RC}}^{\text{max}}]$ gets bigger with $m_{\tilde{\chi}_1^0}$ increase.
- The end-point of m_{LSP}^{\max} at $m_{\tilde{\chi}_1^0}$ are much obvious.

$$0 < m_{\rm RC}^{\rm min} \le m_{\rm P}^{\rm true} \le m_{\rm RC}^{\rm max} < \sqrt{s} / 2$$
$$0 \le m_{\rm I}^{\rm ture} \le m_{\rm LSP}^{\rm max} \le m_{\rm Recoil} / 2$$

P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV



nt

e

Reconstructed mass variables: Smuon pair production



Oct-26, 2023. Nanjing U.

P.X. Zhu Prospect of off-slepton production at CEPC@240 GeV



Prospect of smuon pair production at CEPC



- Degenerate left-handed and right-handed smuon mass $m_{\tilde{\mu}}$, bino-like LSP $\tilde{\chi}_1^0$.
- The distributions closely related to the $m_{\tilde{\mu}}$ and $m_{\tilde{\chi}_1^0}$
- Three similar signal categories to arXiv:2203.10580



Events 10⁴

10

J. Yuan, H. Cheng and X. Zhuang. arXiv: 2203.10580 [hep-ex]

Oct-26, 2023. Nanjing U.





Prospect of smuon pair production at CEPC

- Preselection Criteria: OS muon pair, $E_{\rm T}^{\rm miss} > 1 \,{\rm GeV}, E_{\mu} > 1$ GeV, $|\eta^{\ell}| < 3, m_{\rm RC}^{\rm max} > 115$ GeV.
- 12 SRs defined due to the distributions are much shape and closed related to the sparticle masses.
- Very good signal noise ratio are reached.
- For $m_{\tilde{\mu}} = 115$ GeV, about 110K smuon events can be generated at CPEC @ 240 GeV, 5 fb^{-1} .
- More than 10K event can be kept in the most sensitive SR.



Oct-26, 2023. Nanjing U.



\mathbf{SR}	SRL-01	SRL-02	SRM-01	SRM-02
		$m_{ m RC}^{ m max}$ >117	$m_{ m RC}^{ m min} > 85$	$m_{ ext{LSP}}^{ ext{max}} \in [50, 2]$
	$m_{ m RC}^{ m min}>\!\!85$	$m_{ m RC}^{ m min}>\!\!95$	$m_{ m LSP}^{ m max} \in [40,60]$	$m_{ m RC}^{ m max}(40)>$
		$E_{\mu^{\pm}} \in [50,70]$	$m_{ m RC}^{ m max}(40) > 110$	$m_{ m RC}^{ m min}(40)>$
BKG	7532 ± 86	900 ± 30	2079 ± 45	970 ± 31
SGN(0)	38900	10900	6360	2270
SGN(40)	23800	1410	22100	10200
SGN(80)	0	0	0	0
$\mathrm{SGN}(110)$	0	0	0	0
SR	SRM-03	SRM-04	SRM-05	SRH-01
	$m_{ m LSP}^{ m max} \in [60, 80]$	$m_{ m LSP}^{ m max} \in [70, 85]$	$m_{ m LSP}^{ m max} \in [80,95]$	$E_{\mu^{\pm}} \in [34, 44]$
	$m_{ m RC}^{ m min}(40) > 95$	$m_{ m RC}^{ m min}(70)>100$	$m_{ m RC}^{ m max}(80) > 105$	$\Delta m^{\max} \in [35]$
BKG	912 ± 30	19941 ± 141	14039 ± 118	6412 ± 80
SGN(0)	59	468	0	49
SGN(40)	2460	4970	359	4340
SGN(80)	0	23900	74100	11600
$\mathrm{SGN}(110)$	0	0	0	0
SR	SRH-02	SRH-03	SRH-04	SRH-05
	$E_{\mu^{\pm}} \in [28, 37]$	$E_{\mu^{\pm}} \in [22, 28]$	$E_{\mu^{\pm}} \in [15, 18]$	$\Lambda mmax \subset [0]$
	$\Delta m^{\max} \in [25, 40]$	$\Delta m^{\rm max} \in [15, 30]$	$\Delta m^{\rm max} \in [0,20]$	$\Delta m \in [0,$
BKG	5913 ± 76	3190 ± 56	786 ± 28	262 ± 12
SGN(0)	0	0	0	0
SGN(40)	5	0	0	0
SGN(80)	29200	11900	0	0
SGN(110)	0	0	0	112000









Prospect of smuon pair production at CEPC

- Preselection Criteria: OS muon pair, $E_{\rm T}^{\rm miss} > 1 \,{\rm GeV}, E_{\mu} > 1$ GeV, $|\eta^{\ell}| < 3, m_{\rm RC}^{\rm max} > 115$ GeV.
- closed related to the sparticle masses.
- generated at CPEC @ 240 GeV, 5 fb^{-1} .



Oct-26, 2023. Nanjing U.



Application in Precision measurement

Measuring the W boson mass in full-leptonic decay channel

- $e^+e^- \to WW \to \ell \ell' \nu \nu$.
- The backgrounds are much more cleaner.
- The distributions of $m_{\rm RC}^{\rm min}$ and $m_{\rm RC}^{\rm max}$ are sensitive to m_W .



Oct-26, 2023. Nanjing U.





- Template-fit using MC data.
- The uncertainty is about 3 MeV at CEPC. The precision is at same order of semi-leptonic decayed channel ($\ell \nu q q'$), please see CEPC CDR.
- The uncertainty may be affect by the beam energy uncertainty.









Summary

- CEPC can provide a precise measurement of the Lorentz four momentum vector of the particles in final states.
- We found a Lorentz invariant way to reconstruct semi-invisibly decayed events at CEPC:



- We get:
 - Decay topology independent & Lorentz invariant mass variables: $m_{\rm RC}^{\rm min}$, $m_{\rm RC}^{\rm max}$ and $m_{\rm LSP}^{\rm max}$; **A**)
 - B) They can get improve the statistical sensitivity of both new physics search & SM measurement.

Oct-26, 2023. Nanjing U.



Transfer the geometry to Put Constraints/Assumption kinematic information $\int \mathcal{L} = 5.05 \text{ ab}^{-1}, \quad \sqrt{s} = 240 \text{ GeV}$ $_{0}^{0}$ 25000 $W^+W^-, W \rightarrow \mu\nu_{\mu}$ [GeV] Events /(1 GeV × 1 GeV) mRC PV 15000 100 20 80 40 -60

 $m_{\rm RC}^{\rm min}$ [GeV]

The 2023 international workshop on the high energy CPEC Oct 23-27, 2023. Nanjing University



In collaboration with Jin–Min Yang, Yang Zhang & Rui Zhu. Based on PRD. 108 (2023) 7, 075015 [arXiv: 2211.01832]



Pengxuan Zhu(朱鹏轩)

Institute of Theoretical Physics, Chinese Academy of Science Oct-26, 2023

