

Elliptic anisotropy measurement of the  $f_0(980)$  in pPb collisions and determination of its quark content by CMS

An Gu  
(for the CMS Collaboration)  
CMS-PAS-HIN-20-002

Purdue University

December 17, 2023



Supported in part by the




U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

# Overview

1. Introduction and Physics Motivation
2. Data Analysis
3.  $v_2$  Results and Systematic Uncertainties
4. Extraction of  $n_q$  for  $f_0(980)$
5. Conclusion

# Introduction and Physics Motivation: Exotic hadrons



Example of exotic hadron: tetra-quark

- ▶ Exotic hadrons: configurations other than the usual  $q\bar{q}$  and  $qqq(\bar{q}\bar{q}\bar{q})$

- ▶  $f_0(980)$ : candidate exotic hadron first observed in  $\pi\pi$  scattering experiments in the 1970's

S.D. Protopopescu, Phys. Rev. D 7 (1973) 1279;  
B. Hyams, Nucl. Phys. B 64(1973) 134;  
G. Grayer, Nucl. Phys. B 75 (1974) 189

- ▶ The configuration of  $f_0(980)$  is still controversial:  
 $q\bar{q}$  meson,  $q\bar{q}q\bar{q}$  tetraquark,  $q\bar{q}g$  hybrid, or  $K\bar{K}$  molecule

D.V. Bugg, Phys. Rept. 397 (2004) 257;  
E. Klempt and A. Zaitsev, Phys. Rept. 454 (2007) 1;  
J.R. Pelaez, Phys. Rept. 658 (2016) 1

# Introduction and Physics Motivation: Elliptic Flow $v_2$ and NCQ scaling


- ▶ Azimuthal anisotropy:

$$\frac{dN}{d\phi} \propto 1 + 2 \sum_{n=1}^{\infty} v_n \cos[n(\phi - \psi_n)], \quad (1)$$

- ▶ Approximate **number of constituent quark (NCQ) scaling** has been observed


$$v_n(p_T)/n_q = v_{n,q}(p_T/n_q) \quad (2)$$

$$v_n(K E_T)/n_q = v_{n,q}(K E_T/n_q) \quad (3)$$



- ▶ **Coalescence** hadronization provides one possible mechanism:  $n_q$  quarks combine into a hadron with  $\sim$  equal momenta.


$$\frac{dN_h}{d\phi} \propto \left( \frac{dN_q}{d\phi} \right)^{n_q} \propto \left[ 1 + \sum 2v_{n,q}(p_T^q) \cos(n[\phi - \psi_n]) \right]^{n_q}$$



- ▶  $v_2$  measurement of  $f_0(980) \rightarrow n_q$


# Data Analysis: Reconstruction of $f_0(980)$

- ▶ Dataset: **pPb collisions** at  $\sqrt{s_{NN}} = 8.16$  TeV in high multiplicity events collected in 2016.
- ▶ Dominant decay channel:  $f_0(980) \rightarrow \pi^+ \pi^-$ .
- ▶ No PID in this analysis; All charged tracks assumed to be pions
- ▶ Mass Spectrum: opposite sign pair  $\pi^+ \pi^-$  subtracted by same sign pair  $\pi^+ \pi^+$ ,  $\pi^- \pi^-$





Large  
Combinatorial  
Background

- ▶ Peak is modeled with **Breit-Wigner function**
- ▶ Residual background: 3rd order polynomial
- ▶ Fitting range:  $0.8 < m_{\pi\pi} < 1.7$  GeV/c<sup>2</sup>





# Data Analysis: $v_2$ Extraction

- ▶ Yield of  $f_0(980)$  extracted for different  $\phi - \psi_2$  ranges
- ▶ Event-plane resolution corrected




# Non-flow subtraction

- ▶ Use published nonflow data of  $K_s^0$  from low-multiplicity subtraction [10.1103/PhysRevLett.121.082301]
- ▶ Assume **relative nonflow/flow** to be as same as that of  $K_s^0$ . Use **absolute nonflow** as a systematic. (infeasible to do  $f_0(980)$  low-multiplicity subtraction)



# $v_2^{\text{sub}}$ results and systematic uncertainties



## ► Systematic uncertainties of $f_0(980) v_2$


- Mix-Event Correction
- Track Selection
- Event-plane Resolution
- Signal Form
- Residual Background Form
- Fit Range
- Nonflow Subtraction

## ► Systematic uncertainties of $f_0(980) n_q$

Source	$n_q$ uncertainty
Statistical	0.16
$f_0(980) v_2$ systematics	0.13
Non-flow effects on $v_2^{\text{sub}}$	0.04
NCQ-scaling fit parameters	0.02
NCQ-scaling functional form	0.04
NCQ-scaling using $p_T/n_q$	0.06




# $n_q$ extraction for $f_0(980)$



► NCQ scaling fit in  $\frac{KE_T}{n_q}$ :

$$\frac{KE_T}{n_q} \left( p_0 + p_1 \frac{KE_T}{n_q} \right) e^{-p_2 \frac{KE_T}{n_q}} .$$

# $n_q$ extraction for $f_0(980)$




- ▶ NCQ scaling fit in  $\frac{KE_T}{n_q}$ :

$$\frac{KE_T}{n_q} \left( p_0 + p_1 \frac{KE_T}{n_q} \right) e^{-p_2 \frac{KE_T}{n_q}}.$$

- ▶ Qualitatively consistent with  $n_q = 2$  for  $f_0(980)$ .

# $n_q$ extraction for $f_0(980)$




- ▶ NCQ scaling fit in  $KE_T/n_q$ :

$$\frac{KE_T}{n_q} \left( p_0 + p_1 \frac{KE_T}{n_q} \right) e^{-p_2 \frac{KE_T}{n_q}} .$$


- ▶ Qualitatively inconsistent with  $n_q = 4$  for  $f_0(980)$ .

# Significance to exclude $n_q = 4$ and $n_q = 3$ hypothesis

- ▶  $\chi^2 = (\vec{y} - \vec{f})^T (C_y + C_f)^{-1} (\vec{y} - \vec{f})$ , with uncertainty covariance matrix.
- ▶ **Measured  $f_0(980)$  data** log-likelihood ratio  $-2 \ln (L_{n_q=4}/L_{n_q=2})$ , i.e.  $\chi^2$  difference
- ▶ Pseudo-experiment assuming  $n_q = 4$  (**yellow peak**):
  - $v_2^{\text{sub}}$  from **NCQ-scaling curve  $\times 4$** ; Smearing with uncertainty.
  - Same calculation of log-likelihood ratio as data  $\rightarrow$  **yellow peak**  $\rightarrow$  get significance
- ▶ Pseudo-experiment assuming  $n_q = 2$  (**green peak**)





NCQ scaling measured up to  $p_T/n_q \approx 3$  GeV/c, so use  $p_T < 10$  GeV/c for  $n_q = 4$  case



NCQ scaling measured up to  $p_T/n_q \approx 3$  GeV/c, so use  $p_T < 8$  GeV/c for  $n_q = 3$  case


# $\chi^2$ scan to extract $n_q$

- ▶  $\chi^2 = (\vec{y} - \vec{f})^T (C_y + C_f)^{-1} (\vec{y} - \vec{f})$ , with uncertainty covariance matrix.



- ▶ NCQ scaling is measured up to  $p_T/n_q \sim 3$  GeV/c, so we use  $f_0(980)$  data within  $p_T < 6$  GeV/c. Extracted  $n_q = 2.4 \pm 0.4$ .
- ▶ Assuming NCQ scaling holds beyond  $p_T/n_q \sim 3$  GeV/c, then  $n_q = 2.10 \pm 0.24$  ( $2.07 \pm 0.22$ ) using data from  $p_T < 8$  ( $10$ ) GeV/c.

# Conclusion




- ▶  $v_2$  of  $f_0(980)$  measured as a function of  $p_T$  up to 10 GeV/c
- ▶ Assuming NCQ scaling,  $n_q$  of  $f_0(980)$  is consistent with 2.
- ▶  $n_q = 4$  (tetra-quark state or  $K\bar{K}$  molecule) excluded with  $7.7\sigma$ .
- ▶  $n_q = 3$  ( $q\bar{q}g$  hybrid) excluded with  $3.5\sigma$ .
- ▶ Our data favor  $q\bar{q}$  normal meson state for  $f_0(980)$ .

# Back-up: Eventplane Reconstruction


- ▶ Event Plane:  $\psi_n = \frac{1}{n} \text{atan2}(\sum_i w_i \sin(n\phi_i), \sum_i w_i \cos(n\phi_i))$ ,  
 $i^{\text{th}}$ -tower of forward hadron calorimeter (HF) ( $3 < |\eta| < 5$ );  $\phi_i$  azimuthal angle,  $w_i$  transverse energy in each tower as weight
- ▶ Event Plane Recentering and Flattening
  - Recentering:  
$$\psi_n = \frac{1}{n} \text{atan2}(\sum_i w_i \sin(n\phi_i) - \langle \sum_i w_i \sin(n\phi_i) \rangle, \sum_i w_i \cos(n\phi_i) - \langle \sum_i w_i \cos(n\phi_i) \rangle)$$
  
<> indicates the average over all events in the same centrality class and vertex locations
  - Flattening:  
$$\psi_n = \psi'_n \left( 1 + \sum_{j=1}^{j_{max}} \frac{2}{jn} (-\langle \sin(jn\psi'_n) \rangle \cos(jn\psi'_n) + \langle \cos(jn\psi'_n) \rangle \sin(jn\psi'_n)) \right)$$
- ▶ HF calorimeter in the Pb-going direction for better resolution ( $3 < \eta < 5$  for pPb beam,  $-5 < \eta < -3$  for Pbp beam)

# Pseudo-experiments to exclude $n_q = 4$ hypothesis


- ▶ **Observed:**  $-2 \ln(L_{n_q=4}/L_{n_q=2})$ : log-likelihood ratio , ( $\chi^2$  difference)
- ▶ Pseudo-experimental data:
  - $f_0(980) v_2^{\text{sub}}$  from NCQ-scaling curve for a given  $n_q$  hypothesis; Smearing with uncertainty.
  - Distribution (yellow peak) fit by a Gaussian
  - Significance extracted from the observed (red)



$0 < p_T < 10 \text{ GeV}/c$



$0 < p_T < 8 \text{ GeV}/c$



$0 < p_T < 6 \text{ GeV}/c$





# Systematic Uncertainties

- ▶ Systematic uncertainties of  $f_0(980) v_2$ 
  - Mix-Event Correction
    - Applying mixevent  $H_{OS,mixEvent}/H_{SS,mixEvent}$  on  $H_{SS,singleEvent}$
  - Track Selection
    - loose and tight track selections
  - Track Efficiency
    - track efficiency correction decreased and increased by 2.4%
  - Event-plane Resolution
    - Error propagation of uncertainties in event-plane resolution
  - Signal Form
    - Breit-Wigner and relativistic Breit-Wigner
  - Residual Background Form
    - 2nd, 3rd (default), 4th, 5th-order polynomial
  - Fit Range
    - Vary  $0.02 GeV/c^2$  on each side
  - Nonflow Subtraction
    - $K_s^0 \Delta v_2/v_2$  (default),  $K_s^0 \Delta v_2$

- ▶ Systematic uncertainties of  $f_0(980) n_q$ 
  - Statistical: error propagated from  $f_0(980) v_2$  statistical uncertainties
  - $f_0(980) v_2$  systematics: error propagated from  $f_0(980) v_2$  systematic (nonflow uncertainty not included)
  - Non-flow effects on  $v_2^{\text{sub}}$ : error propagated from  $f_0(980) v_2$  nonflow systematic
  - NCQ-scaling fit parameters:  $n_q$  uncertainty due to fit parameter uncertainties
  - NCQ-scaling functional form: standard deviation of  $n_q$  with different fit function form
  - NCQ-scaling using  $p_T/n_q$ :  $n_q$  difference from using  $v_2^{\text{sub}}/n_q$  vs.  $KE_T/n_q$

# NCQ scaling of $v_2$



- ▶ NCQ scaling fit in  $KE_T/n_q$ :  $f(KE_T/n_q) = KE_T/n_q (p_0 + p_1 KE_T/n_q) e^{-p_2 KE_T/n_q}$ .
- ▶ Qualitatively consistent with  $n_q = 2$  for  $f_0(980)$ .