

The spatial resolution of AC-coupled LGAD developed by IHEP

Monday, 25 September 2023 15:45 (15 minutes)

AC-coupled Low- Gain Avalanche Detectors (AC-LGAD) are designed as detectors with 100% fill factor for high precision 4D-tracking, which have been studied and researched by many institutes including BNL, FBK et al. Institute of High Energy Physics (IHEP) has also done many researches on AC-LGAD. First IHEP AC-LGAD sensors with a pitch of 2000 μm and AC pad of 1000 μm show time resolution better than 20ps, and spatial resolution better than 16 μm . Testing results show that as the N⁺ doping dose decreases from 10 P to 0.2 P, the spatial resolution is reduced from 35 μm to 16 μm . Details of the results will be discussed. Second version of IHEP AC-LGAD be fabricated, lower than 0.2P n⁺ layer dose be used for improve the spatial resolution and sensors with different pad-pitch structures also be fabricated. Testing results show that as decreasing the n⁺ layer dose from 0.2P to 0.02P, the spatial resolution of AC-LGAD can be lower than 8 μm . Strips with different pad-pitch structures be fabricated and studied, results show that pad-pitch structures will also affect the spatial resolution. Testing results of IHEP AC-LGAD v2 sensors tested by using laser system will also be shown.

Primary author: LI, Mengzhao (高能所)

Presenter: LI, Mengzhao (高能所)

Session Classification: 粒子物理 2 组