



Contribution ID: 242 Contribution code: AAN-1

Type: Oral Presentation

On some unsettled half-lives of AMS radionuclides

Tuesday, 22 October 2024 09:20 (20 minutes)

Half-lives ought to be accurate, and preferably precise as well. In the recently published review on half-lives of long-lived radionuclides (Heinitz et al., 2022), several cases were mentioned where multiple half-life measurements on a specific radionuclide were incompatible with each other within the reported uncertainties. We call these “unsettled” half-lives. There are also cases where only very old half-life measurements (50 to 60 years ago) exist, which may need confirmation.

The direct way to determine a long half-life follows from the radioactive decay law: $dN/dt = -\lambda \cdot N$, where N is the number of radionuclides, dN/dt is its decay rate (activity), and λ the decay constant related to the half-life via $\lambda = \ln 2/t_{1/2}$. Both N and dN/dt need to be measured accurately and independently to obtain the half-life.

The half-life of ^{10}Be can be considered as a good example of how early (trivial) mistakes were corrected and eventually an accurate and precise half-life value of $(1.387 \pm 0.012) \times 10^6$ y was established from two independent measurements (Korschinek et al., 2010, Chmeleff et al., 2010).

In this contribution, we want to discuss unsettled half-lives of some radionuclides where AMS was partly involved in the half-life measurement itself, and which are of interest for applications through AMS measurements. Among others, these comprise the radionuclides ^{32}Si , ^{39}Ar , ^{53}Mn , ^{59}Ni , ^{79}Se , ^{135}Cs , and ^{146}Sm .

We will discuss some ongoing and planned half-life measurements on these radionuclides, which hopefully will lead to a firmly accepted value. The number of radionuclides in the sample whose activity needs to be measured is a crucial input for a half-life determination. Different methods to measure radionuclide concentrations (e.g. AMS, ICP-MS) will be mentioned. In particular, a critical assessment of the measurement of absolute isotope ratios with AMS will be presented. In some cases, geophysical half-life measurements can also be combined with physical measurements to confirm or refute half-life values.

S. Heinitz, I. Kajan, and D. Schumann, How accurate are half-life data of long-lived radionuclides? *Radiochim. Acta* 110/6-9 (2022) 589-608.

G. Korschinek et al., A new value for the half-life of ^{10}Be by heavy-ion elastic recoil detection and liquid scintillation counting. *Nucl. Instr. Meth. Phys. Res. B* 268 (2010) 187–191.

J. Chmeleff et al., Determination of the half-life of ^{10}Be by multicollector ICP-MS and liquid scintillation counting. *Nucl. Instr. Meth. Phys. Res. B* 268 (2010) 192–199.

Student Submission

No

Primary authors: KUTSCHERA, Walter (University of Vienna, Faculty of Physics, Isotope Physics, A-1090 Vienna, Austria); MARTSCHINI, Martin (University of Vienna, Faculty of Physics, Isotope Physics, A-1090 Vienna, Austria); STEIER, Peter (University of Vienna, Faculty of Physics, Isotope Physics, A-1090 Vienna, Austria); WIESER, Alexander (University of Vienna, Faculty of Physics, Isotope Physics, A-1090 Vienna, Austria); PAUL, Michael (Racah Institute of Physics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel); SCHUMANN, Dorothea (Isotope and Target Chemistry, Paul Scherrer Institute (PSI), CH-5232 Villigen, Switzerland); VOCKENHUBER, Christof (Laboratory of Ion Beam Physics, ETH Zuerich, CH-8083, Switzerland); LACHNER, Jo-

hannes (Helmholtz-Zentrum Dresden-Rossendorf, Dresden, Germany); WALLNER, Anton (Helmholtz-Zentrum Dresden-Rossendorf, Dresden, Germany)

Presenter: KUTSCHERA, Walter (University of Vienna, Faculty of Physics, Isotope Physics, A-1090 Vienna, Austria)

Session Classification: Applications in Astrophysics and Nuclear Sciences

Track Classification: Applications in Astrophysics and Nuclear Sciences