

Temporal Changes of Iodine-129 in the Canada Basin Over the Past Decade

Yuanzhi Qi^{1*}, Takeyasu Yamagata¹, Hiroyuki Matsuzaki¹, Hisao Nagai¹, Yuichiro Kumamoto², Qiuyu Yang¹, Xinru Xu¹

1. MALT (Micro Analysis Laboratory, Tandem accelerator), The University Museum, The University of Tokyo

2. Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology

*Correspondence to: Yuanzhi Qi; E-mail: yuanzhiqi@um.u-tokyo.ac.jp

The Arctic, a region highly sensitive to global climate change, has experienced a temperature increase exceeding four times the global average rate since the onset of the industrial era. Concurrently, warmer and saltier Atlantic water is encroaching further northward into the Arctic Ocean, driving the phenomenon known as "atlantification." These shifts have led to significant Arctic sea ice loss and alterations in seawater thermohaline structure, potentially impacting Arctic ocean currents. Understanding the response mechanisms of these currents to climate change is crucial for comprehending and addressing the broader impacts of climate change. Notably, substantial quantities of iodine-129 (^{129}I) have been discharged directly into the North Atlantic from nuclear fuel reprocessing plants (NFRPs). Enriched with NFRP-derived ^{129}I , Atlantic Waters penetrate the Arctic region via the Fram Strait and the Barents Sea, rendering ^{129}I an ideal tracer for tracking Arctic currents. While numerous studies have utilized ^{129}I to investigate ocean currents in the Eurasian Basin of the Arctic Ocean, research in the Canada Basin, characterized by equally intricate current structures, remains scarce. Here, we present a decade-long analysis of temporal changes in ^{129}I concentrations at two stations (NAP and CB, Fig. 1) within the Canada Basin spanning from 2013 to 2023. Our findings reveal a diminishing influence of Atlantic water on the surface of the Canada Basin over time. In the early 2010s, ^{129}I concentrations in the intermediate layer of both stations remained consistent, suggesting a common source. However, by the late 2010s, the concentration of ^{129}I in the intermediate layer of the NAP station significantly surpassed that of the CB station, indicative of a shift in ocean currents, likely influenced by changes in the Arctic Oscillation and potentially linked to atlantification.




Fig. 1. Map of the Arctic Ocean showing the sampling locations over past decade.