

Contribution ID: 231 Contribution code: NIM-6

Type: Oral Presentation

Counting Ar-42 atoms at the 10⁻²¹ isotopic-abundance level

Friday, 25 October 2024 11:20 (20 minutes)

Atom Trap Trace Analysis (ATTA) is a technique that utilizes laser to selective capture and detect individual atoms of the desired isotope. This technique loads atoms into a magneto-optical trap (MOT) through laser cooling and trapping, realizing high sensitivity and detection efficiency of the desired isotope.

Due to the use of multi-photon repetitive excitation in the experiment, the ATTA method has extremely high isotope selectivity and has been successfully applied to the detection of Ar-39 with an abundance level of 10⁻17.

In recent years, we have significantly improved the counting rate and detection efficiency of the Ar isotope ATTA device, and developed a strong flow mass spectrometry isotope pre-concentration method. These advancements have enabled us to successfully detect Ar-42 (with a half-life of 33 years) atoms in the air with an isotopic-abundance level of only 10⁻²¹ using single atom counting method.

Previously, people used dark matter detectors detecting the decay of Ar-42 from tons of liquid argon to calculate the natural abundance of Ar-42. The sample consumption in this work is only 1-2L of gaseous argon, and the final measured natural abundance of Ar-42 is $(5.4\pm0.8)\times10^{-21}$, which is twice more accurate than previous measurements. This work demonstrates the lowest detection limit that can be achieved by all current ultra-sensitive isotope detection methods.

Student Submission

Yes

Primary authors: WAN, Zhaofeng; 梁, 佳炜 (中国科学技术大学); 杨, 国民 (中国科学技术大学); 卢, 征 天 (中国科学技术大学); 贾, 泽华 (近代物理研究所); 孙, 良亭 (近代物理研究所); 蒋, 蔚 (中国科学技术大学)

Presenter: WAN, Zhaofeng

Session Classification: New Isotopes Methodologies

Track Classification: New Isotopes Methodologies