

Contribution ID: 186 Contribution code: ACI-4

Type: Oral Presentation

A first minimum in-situ 10Be age for the Tswaing Meteor Crater, South Africa.

Wednesday, 23 October 2024 10:40 (20 minutes)

The well-preserved Tswaing Meteor Crater–aka Pretoria Saltpan –with a diameter of 1130 m is located ca. 40 km NNW of Tshwane (Pretoria). It has been a subject of several scientific investigations. The crater is formed on a ca. 2 Ga Nebo Granite, the acid phase, of the Bushveld Complex (Lebowa Granite Suite). The interior structure floor is not far removed from the local ground water level and hence the floor is partially filled by a highly saline lake (Tswaing means the place of salt). The elevation of the rim over the present structure floor is ~120 m, while it rises about 60 m above the surrounding plain. The rim, with minimal seasonal brush, has a fair number of granitoid boulders. These are thought to have been uprooted during the cratering event from the Nebo granite basement bedrock. The basement rock is some 200 m below the current floor. The drill cores from this lake give a unique insight into past climate and flora of the region (Partridge et al. 1993, Kristen et al. 2007, Metwally et al. 2014, Scott 2016).

The meteorite impact origin of Tswaing was largely inferred from morphological observations and was later supported by structural and experimental fission track measurements (Reimold et al., 1992). The currently accepted age of 220±50 ka the is based on fission track dating (Storzer et al. 1993). However, this is an average over wildly varying measuements. While a similar result age was inferred by the core drilling on the crater floor that revealed an infilling consisting of 90 m of organic lacustrine sediments, this is based on extrapolation from the average infilling rate of 50 cm/ka of the radiocarbon dated section of the core (Partridge et al, 1993). This rate also has evident significant variance over the first 20 m of the core.

The boulders from the ejecta of the impact around the rim of the crater present an excellent opportunity to independently verify the age of the cratering event through the exposure ages of these boulders by cosmogenic radionuclides using accelerator mass spectrometry. To this end we collected 15 samples from around rim, pre-treated them, and processed selection of 10 at the Helmholtz-Zentrum Dresden-Rossendorf for AMS analysis. Here we will present first results from the AMS measurement of the samples, a contextual interpretation, and the implications for the actual age of the Tswaing impact crater.

References:

Kristen, I., et al. "Hydrological changes in southern Africa over the last 200 Ka as recorded in lake sediments from the Tswaing impact crater." South African Journal of Geology 110.2-3 (2007): 311-326.

Metwally, A. A., et al. "Holocene palynology and palaeoenvironments in the Savanna Biome at Tswaing Crater, central South Africa." Palaeogeography, Palaeoclimatology, Palaeoecology 402 (2014): 125-135.

Partridge, T. C., et al. "The Pretoria Saltpan: a 200,000 year southern African lacustrine sequence." Palaeogeography, Palaeoclimatology, Palaeoecology 101.3-4 (1993): 317-337.

Reimold, Wolf Uwe, et al. "Pretoria Saltpan crater: Impact origin confirmed." Geology 20.12 (1992): 1079-1082. Scott, L. "Fluctuations of vegetation and climate over the last 75 000 years in the Savanna Biome, South Africa: Tswaing Crater and Wonderkrater pollen sequences reviewed." Quaternary Science Reviews 145 (2016): 117-133.

Storzer, Dieter, Christian Koeberl, and Wolf Uwe Reimold. "The age of the Pretoria Saltpan impact crater, South Africa." In Lunar and Planetary Inst., Twenty-Fourth Lunar and Planetary Science Conference. Part 3: NZ p 1365-1366 (SEE N94-20636 05-91). Vol. 24. 1993.

Student Submission

Primary authors: MBELE, Vela (iThemba LABS, South Africa); Ms KHOSA, RR (TAMS Laboratory, NR-F-iThemba LABS, Private Bag 11, WITS, 2050 and Department of Geological Sciences, University of Cape Town, Rondebosch, South Africa); STUEBNER, Konstanze (Helmholtz-Zentrum Dresden-Rossendorf); WINKLER, Stephan (Helmholtz-Zentrum Dresden-Rossendorf)

Presenter: MBELE, Vela (iThemba LABS, South Africa)

Session Classification: Applications of Cosmogenic Isotopes

Track Classification: Applications of Cosmogenic Isotopes