



Nuclear Lattice Effective Field Theory

– Introduction and Perspectives –

Ulf-G. Meißner, Univ. Bonn & FZ Jülich


supported by DFG, SFB/TR-110

by CAS, PIFI

by DFG, SFB 1639

by ERC, EXOTIC

by NRW-FAIR



中國科学院
CHINESE ACADEMY OF SCIENCES

⟨NUMERIQS⟩



Contents


- Very brief Introduction
- Chiral EFT on a lattice
- The minimal nuclear interaction → Bing-Nan Lu's talk
 - Foundations
 - Applications
 - Extension to hyper-nuclei
 - EoS of neutron matter & neutron stars
- Chiral interactions at N3LO → Dean Lee's talk
 - Foundations
 - Applications to nuclear structure
 - Applications to scattering
- Summary & outlook

Very brief Introduction

Our goal: Ab initio nuclear structure & reactions

- Nuclear structure:


- ★ limits of stability
- ★ 3-nucleon forces
- ★ alpha-clustering
- ★ EoS & neutron stars
- ⋮
- ⋮




- Nuclear reactions, nuclear astrophysics:

- ★ alpha-particle scattering
- ★ triple-alpha reaction
- ★ alpha-capture on carbon
- ⋮
- ⋮

de Boer et al, Rev. Mod. Phys. **89** (2017) 035007




Chiral EFT on a lattice



T. Lähde & UGM


Nuclear Lattice Effective Field Theory - An Introduction

Springer Lecture Notes in Physics **957** (2019) 1 - 396



More on EFTs

- Much more details on EFTs in light quark physics:



<https://www.cambridge.org/de/academic/subjects/physics/theoretical-physics-and-mathematical-physics/effective-field-theories>

Nuclear lattice effective field theory (NLEFT)

7

Frank, Brockmann (1992), Koonin, Müller, Seki, van Kolck (2000) , Lee, Schäfer (2004), . . .
Borasoy, Krebs, Lee, UGM, Nucl. Phys. **A768** (2006) 179; Borasoy, Epelbaum, Krebs, Lee, UGM, Eur. Phys. J. **A31** (2007) 105

- new method to tackle the nuclear many-body problem


- discretize space-time $V = L_s \times L_s \times L_s \times L_t$:
nucleons are point-like particles on the sites

- discretized chiral potential w/ pion exchanges
and contact interactions + Coulomb

→ see Epelbaum, Hammer, UGM, Rev. Mod. Phys. **81** (2009) 1773

- typical lattice parameters

$$p_{\max} = \frac{\pi}{a} \simeq 315 - 630 \text{ MeV [UV cutoff]}$$



- strong suppression of sign oscillations due to approximate Wigner SU(4) symmetry

E. Wigner, Phys. Rev. **51** (1937) 106; T. Mehen et al., Phys. Rev. Lett. **83** (1999) 931; J. W. Chen et al., Phys. Rev. Lett. **93** (2004) 242302

- physics independent of the lattice spacing for $a = 1 \dots 2 \text{ fm}$

Alarcon, Du, Klein, Lähde, Lee, Li, Lu, Luu, UGM, EPJA **53** (2017) 83; Klein, Elhatisari, Lähde, Lee, UGM, EPJA **54** (2018) 121

Transfer matrix method

- Correlation–function for A nucleons: $Z_A(\tau) = \langle \Psi_A | \exp(-\tau H) | \Psi_A \rangle$

with Ψ_A a Slater determinant for A free nucleons
[or a more sophisticated (correlated) initial/final state]

- Transient energy

$$E_A(\tau) = -\frac{d}{d\tau} \ln Z_A(\tau)$$

→ ground state: $E_A^0 = \lim_{\tau \rightarrow \infty} E_A(\tau)$


- Exp. value of any normal–ordered operator \mathcal{O}

$$Z_A^\mathcal{O} = \langle \Psi_A | \exp(-\tau H/2) \mathcal{O} \exp(-\tau H/2) | \Psi_A \rangle$$

$$\lim_{\tau \rightarrow \infty} \frac{Z_A^\mathcal{O}(\tau)}{Z_A(\tau)} = \langle \Psi_A | \mathcal{O} | \Psi_A \rangle$$




- Excited states: $Z_A(\tau) \rightarrow Z_A^{ij}(\tau)$, diagonalize, e.g. $0_1^+, 0_2^+, 0_3^+, \dots$ in ^{12}C

Euclidean time



Configurations

9




- ⇒ all *possible* configurations are sampled
- ⇒ preparation of *all possible* initial/final states
- ⇒ *clustering* emerges *naturally*

Auxiliary field method

10

- Represent interactions by auxiliary fields (Gaussian quadrature):


$$\exp \left[-\frac{C}{2} (N^\dagger N)^2 \right] = \sqrt{\frac{1}{2\pi}} \int ds \exp \left[-\frac{s^2}{2} + \sqrt{C} s (N^\dagger N) \right]$$



Comparison to lattice QCD

11

LQCD (quarks & gluons)	NLEFT (nucleons & pions)
relativistic fermions	non-relativistic fermions
renormalizable th'y	EFT
continuum limit	no continuum limit
(un)physical masses	physical masses
Coulomb - difficult	Coulomb - easy
high T/small ρ	small T/nuclear densities
sign problem severe	sign problem moderate



- For nuclear physics, NLEFT is the far better methodology!

Computational equipment

- Present = JUWELS (modular system) + FRONTIER + ...



The minimal nuclear interaction: Foundations

A minimal nuclear interaction

- Basic problem: Straightforward application of chiral EFT forces leads to problems when one goes beyond light nuclei (e.g. the radius problem)
- Main idea: Construct a minimal nuclear interactions that reproduces the ground state properties of light nuclei, medium-mass nuclei, and neutron matter simultaneously with no more than a few percent error in the energies and charge radii
- This can be achieved by making use of Wigner's SU(4) spin-isospin symmetry
Wigner, Phys. Rev. **C 51** (1937) 106
- If the nuclear Hamiltonian does not depend on spin and isospin, then it is obviously invariant under SU(4) transformations [really $U(4) = U(1) \times SU(4)$]:

$$\mathbf{N} \rightarrow U\mathbf{N}, \quad U \in SU(4), \quad \mathbf{N} = \begin{pmatrix} p \\ n \end{pmatrix}$$

$$\mathbf{N} \rightarrow \mathbf{N} + \delta\mathbf{N}, \quad \delta\mathbf{N} = i\epsilon_{\mu\nu}\sigma^\mu\tau^\nu \mathbf{N}, \quad \sigma^\mu = (1, \boldsymbol{\sigma}_i), \quad \tau^\mu = (1, \boldsymbol{\tau}_i)$$

Remarks on Wigner's SU(4) symmetry

Essential elements for nuclear binding

16

Lu, Li, Elhatisari, Epelbaum, Lee, UGM, Phys. Lett. B 797 (2019) 134863 [arXiv:1812.10928]

- Highly SU(4) symmetric LO action without pions, only **four** parameters




$$H_{\text{SU}(4)} = H_{\text{free}} + \frac{1}{2!} C_2 \sum_n \tilde{\rho}(n)^2 + \frac{1}{3!} C_3 \sum_n \tilde{\rho}(n)^3$$

$$\tilde{\rho}(n) = \sum_i \tilde{a}_i^\dagger(n) \tilde{a}_i(n) + s_L \sum_{|n'-n|=1} \sum_i \tilde{a}_i^\dagger(n') \tilde{a}_i(n')$$

$$\tilde{a}_i(n) = a_i(n) + s_{NL} \sum_{|n'-n|=1} a_i(n')$$

s_L controls the locality of the interactions, s_{NL} the non-locality of the smearing

→ describes binding energies, radii, charge densities and the EoS of neutron matter



The minimal nuclear interaction: Applications

Wigner's SU(4) symmetry and the carbon spectrum

18

- Study of the spectrum (and other properties) of ^{12}C

↪ spin-orbit splittings are known to be weak

Hayes, Navratil, Vary, Phys. Rev. Lett. **91** (2003) 012502 Johnson, Phys. Rev. C **91** (2015) 034313

↪ start with cluster and shell-model configurations

→ next slide

- Fit the four parameters:

C_2, C_3 – ground state energies of ^4He and ^{12}C


s_L – radius of ^{12}C around 2.4 fm

s_{NL} – best overall description
of the transition rates

- Calculation of em transitions


requires coupled-channel approach

e.g. 0^+ and 2^+ states




Configurations


- Cluster and shell model configurations




— isoscele right triangle



— “bent-arm” shape




— linear diagonal chain




— acute isoscele triangle

Gaussian wave packets
 $w = 1.7 - 2.1 \text{ fm}$




— ground state $|0\rangle$

$1d_{3/2}$
 $2s_{1/2}$
 $1d_{5/2}$
 $1s_{1/2}$



— $2p\text{-}2h$ state, $J_z = 0$

$1d_{3/2}$
 $2s_{1/2}$
 $1d_{5/2}$
 $1s_{1/2}$





— $1p\text{-}1h$ state, $J_z^{(1)} = J_z^{(2)} = 1$

Transient energies

20


- Transient energies from cluster and shell-model configurations



Spectrum of ^{12}C

Shen, Elhatisari, Lähde, Lee, Lu, UGM, Nature Commun. 14 (2023) 2777

- Improved description when 3NFs are included, amazingly good



→ solidifies earlier NLEFT statements about the structure of the 0_2^+ and 2_2^+ states

Electromagnetic properties

Shen, Elhatisari, Lähde, Lee, Lu, UGM, Nature Commun. **14** (2023) 2777

- Radii (be aware of excited states), quadrupole moments & transition rates



	NLEFT	FMD	α cluster	BEC	RXMC	Exp.
$r_c(0_1^+)$ [fm]	2.53(1)	2.53	2.54	2.53	2.65	2.47(2)
$r(0_2^+)$ [fm]	3.45(2)	3.38	3.71	3.83	4.00	—
$r(0_3^+)$ [fm]	3.47(1)	4.62	4.75	—	4.80	—
$r(2_1^+)$ [fm]	2.42(1)	2.50	2.37	2.38	—	—
$r(2_2^+)$ [fm]	3.30(1)	4.43	4.43	—	—	—

	NLEFT	FMD	α cluster	NCSM	Exp.
$Q(2_1^+)$ [$e \text{ fm}^2$]	6.8(3)	—	—	6.3(3)	8.1(2.3)
$Q(2_2^+)$ [$e \text{ fm}^2$]	−35(1)	—	—	—	—
$M(E0, 0_1^+ \rightarrow 0_2^+)$ [$e \text{ fm}^2$]	4.8(3)	6.5	6.5	—	5.4(2)
$M(E0, 0_1^+ \rightarrow 0_3^+)$ [$e \text{ fm}^2$]	0.4(3)	—	—	—	—
$M(E0, 0_2^+ \rightarrow 0_3^+)$ [$e \text{ fm}^2$]	7.4(4)	—	—	—	—
$B(E2, 2_1^+ \rightarrow 0_1^+)$ [$e^2 \text{ fm}^4$]	11.4(1)	8.7	9.2	8.7(9)	7.9(4)
$B(E2, 2_1^+ \rightarrow 0_2^+)$ [$e^2 \text{ fm}^4$]	2.5(2)	3.8	0.8	—	2.6(4)

Electromagnetic properties cont'd

Shen, Elhatisari, Lähde, Lee, Lu, UGM, Nature Commun. **14** (2023) 2777

- Form factors and transition ffs [essentially parameter-free]:



Sick, McCarthy, Nucl. Phys. A 150 (1970) 631

Strehl, Z. Phys. 234 (1970) 416



Crannell et al., Nucl. Phys. A 758 (2005) 399

Chernykh et al., Phys. Rev. Lett. 105 (2010) 022501

Emergence of geometry

24


- Use the pinhole algorithm to measure the distribution of α -clusters/matter:



- equilateral & obtuse triangles \rightarrow 2^+ states are excitations of the 0^+ states

Emergence of duality




- ^{12}C spectrum shows a cluster/shell-model duality



- dashed triangles: strong 1p-1h admixture in the wave function

Sanity check

- Repeat the calculations w/ the time-honored N2LO chiral interaction
 - ↪ better NN phase shifts than the SU(4) interaction
 - ↪ but calculations are much more difficult (sign problem)




- spectrum as before (good agreement w/ data)
- density distributions as before (more noisy, stronger sign problem)

The ^4He form factor puzzle


- Recent Mainz measurements of $F_{M0}(0_2^+ \rightarrow 0_1^+)$ appear to be in stark disagreement with *ab initio* nuclear theory Kegel et al., Phys. Rev. Lett. **130** (2023) 152502

- Monopole transition ff



[calculations from 2013]

- low-momentum expansion



⇒ A low-energy puzzle for nuclear forces?

Ab initio calculation of the ${}^4\text{He}$ transition form factor 28

UGM, Shen, Elhatisari, Lee, Phys. Rev. Lett. **132** (2024) 062501 [2309.01558 [nucl-th]]

- Use the essential elements action, **all parameters fixed!**
- Calculate the transition ff and its low-energy expansion from the transition density

$$\rho_{\text{tr}}(r) = \langle 0_1^+ | \hat{\rho}(\vec{r}) | 0_2^+ \rangle$$

$$F(q) = \frac{4\pi}{Z} \int_0^\infty \rho_{\text{tr}}(r) j_0(qr) r^2 dr = \frac{1}{Z} \sum_{\lambda=1}^{\infty} \frac{(-1)^\lambda}{(2\lambda + 1)!} q^{2\lambda} \langle r^{2\lambda} \rangle_{\text{tr}}$$

$$\frac{Z|F(q^2)|}{q^2} = \frac{1}{6} \langle r^2 \rangle_{\text{tr}} \left[1 - \frac{q^2}{20} \mathcal{R}_{\text{tr}}^2 + \mathcal{O}(q^4) \right]$$

$$\mathcal{R}_{\text{tr}}^2 = \langle r^4 \rangle_{\text{tr}} / \langle r^2 \rangle_{\text{tr}}$$

- The first excited state sits in the continuum & close to the 3H - p threshold
 - ↪ use large volumes $L = 10, 11, 12$ or $L = 13.2$ fm, 14.5 fm, 15.7 fm
 - ↪ the lattice spacing is fixed to $a = 1.32$ fm, corresponding $\Lambda = \pi/a = 465$ MeV

The first excited state

- 3 coupled channels with 0^+ q.n's \rightarrow accelerates convergence as $L_t \rightarrow \infty$
- Shell-model wave functions (4 nucleons in $1s_{1/2}$, twice 3 in $1s_{1/2}$ and 1 in $2s_{1/2}$)

L [fm]	$E(0_1^+)$ [MeV]	$E(0_2^+)$ [MeV]	ΔE [MeV]
13.2	-28.32(3)	-8.37(14)	0.28(14)
14.5	-28.30(3)	-8.02(14)	0.42(14)
15.7	-28.30(3)	-7.96(9)	0.40(9)

\hookrightarrow statistical and large- L_t errors

\hookrightarrow agreement w/ experiment: $E(0_1^+) = 28.3$ MeV, $\Delta E = 0.4$ MeV

$\hookrightarrow \Delta E$ consistent w/ no-core Gamov shell model (no 3NFs)


Michel, Nazarewicz, Ploszajczak, Phys. Rev. Lett. **131** (2023) 242502

\hookrightarrow consistent w/ the Efimov tetramer analysis $\Delta E = 0.38(2)$ MeV


von Stecher, D'Incao, Greene, Nat. Phys. **5** (2009) 417; Hammer, Platter, EPJA **32** (2007) 113

The transition form factor

- Transition charge density



- Transition form factor



→ agrees with the reconstructed one
from Kamimura

PTEP 2023 (2023) 071D01


→ very small central depletion (no zero)

→ excellent description of the data
→ Coulomb required plus smaller
uncertainty (improved signal)
→ 3NFs important!

The transition form factor II

31

- Small momentum expansion



	$\langle r^2 \rangle_{\text{tr}}$ [fm 2]	\mathcal{R}_{tr} [fm]
Experiment	1.53 ± 0.05	4.56 ± 0.15
Th (AV8' + centr. 3N)*	1.36 ± 0.01	4.01 ± 0.05
Th (AV18 + UIX)	1.54 ± 0.01	3.77 ± 0.08
Th (NLEFT)	1.49 ± 0.01	4.00 ± 0.04

*Hiyama, Gibson, Kamimura, PRC **70** (2004) 031001

- ↪ Also consistent description of the low-energy data
- ↪ No puzzle to the nuclear forces!
- ↪ Can be improved using N3LO action + wave function matching

Elhatisari *et al.*, 2210.17488 [nucl-th]

The minimal nuclear interaction: Extension to hyper-nuclei

The minimal interaction with strangeness I

33

Tong, Elhatisari, UGM, in progress

- Baryon-baryon interaction (consider nucleons and Λ 's plus non-local smearing):

$$V_{\Lambda N} = \textcolor{red}{c_{N\Lambda}} \sum_{\vec{n}} \tilde{\rho}(\vec{n}) \tilde{\xi}(\vec{n}) + \textcolor{red}{c_{\Lambda\Lambda}} \frac{1}{2} \sum_{\vec{n}} \left[\tilde{\xi}(\vec{n}) \right]^2$$

$$\tilde{\rho}(\vec{n}) = \sum_{i,j=0,1} \tilde{a}_{i,j}^\dagger(\vec{n}) \tilde{a}_{i,j}(\vec{n}) + s_L \sum_{|\vec{n}-\vec{n}'|^2=1} \sum_{i,j=0,1} \tilde{a}_{i,j}^\dagger(\vec{n}') \tilde{a}_{i,j}(\vec{n}')$$

$$\tilde{\xi}(\vec{n}) = \sum_{i=0,1} \tilde{b}_i^\dagger(\vec{n}) \tilde{b}_i(\vec{n}) + s_L \sum_{|\vec{n}-\vec{n}'|^2=1} \sum_{i=0,1} \tilde{b}_i^\dagger(\vec{n}') \tilde{b}_i(\vec{n}')$$

- Three-baryon forces (consider nucleons and Λ 's, no non-local smearing):

Peschauer, Kaiser, Haidenbauer, UGM, Weise, Phys. Rev. C 93 (2016) 014001

$$V_{NN\Lambda} = \textcolor{red}{c_{NN\Lambda}} \frac{1}{2} \sum_{\vec{n}} [\rho(\vec{n})]^2 \xi(\vec{n}) , \quad V_{N\Lambda\Lambda} = \textcolor{red}{c_{N\Lambda\Lambda}} \frac{1}{2} \sum_{\vec{n}} \rho(\vec{n}) [\xi(\vec{n})]^2$$

→ must determine 4 LECs! [smearing parameters from the nucleon sector]


→ first time that the $\Lambda\Lambda N$ three-body force is included

The minimal interaction with strangeness II

34

Tong, Elhatisari, UGM, in progress

- Two-body LECs from scattering data (ΛN)
& chiral EFT phase shift ($\Lambda\Lambda$)



- Three-body LECs from the separation energies of Λ and $\Lambda\Lambda$ hyper-nuclei:

$$B_\Lambda(A, Z) = E(A-1, Z) - E(\Lambda, Z)$$

$$B_{\Lambda\Lambda}(A, Z) = E(A-2, Z) - E(\Lambda\Lambda, Z)$$

Nucleus	NLEFT [MeV]	Exp. [MeV]
$^5_\Lambda\text{He}$	3.10(9)	3.10(3)
$^9_\Lambda\text{Be}$	6.64(13)	6.61(7)
$^{13}_\Lambda\text{C}$	11.71(14)	11.80(16)
$^6_{\Lambda\Lambda}\text{He}$	6.96(9)	6.91(16)
$^{10}_{\Lambda\Lambda}\text{Be}$	14.35(13)	14.70(40)

→ this defines our EoS of hyper-nuclear matter called **HMN(I)**

The minimal nuclear interaction: EoS & neutron star properties



Pure neutron matter

36

Tong, Elhatisari, UGM, in progress

- Input: S-wave phase shifts (2N) & symmetric nuclear matter (3N)
- Note: extension of the minimal interaction (leading SU(4) breaking)

⇒ Output: Pure neutron matter (PNM) EoS



– comparable to the renowned APR EoS

Akmal, Pandharipande, Ravenhall, Phys. Rev. C **58** (1998) 1804

– less stiff than the recent AFDMC one

Gandolfi et al., Eur. Phys. J. A **50** (2014) 10

→ work out consequences for neutron stars based on this PNM EoS


Neutron star properties

37


Tong, Elhatisari, UGM, in progress

- Now solve the TOV equations for the PNM and HNM(I) EoSs:

- EoS (PNM and HNM(I))



- Mass-radius relation



- Maximum neutron star mass: $M_{\max} = 2.18(1) M_\odot$ for PNM
 $M_{\max} = 1.54(2) M_\odot$ for HNM(I) \rightarrow need repulsion


EoS of hyper-neutron matter

38


Tong, Elhatisari, UGM, in progress

- Not surprisingly, we need more repulsion [as in the pure neutron matter case]
 - this will move the threshold of $\mu_\Lambda = \mu_n$ up
 - take M_{\max} as data point: $M_{\max} = 1.9M_\odot$ for HNM(II)
 $M_{\max} = 2.1M_\odot$ for HNM(III)

• EoS & speed of sound



• Mass-radius relation



Finite temperature physics

39

- Just two teasers for finite T calculations

→ talks by Bing-Nan Lu and Dean Lee

PHYSICAL REVIEW LETTERS 125, 192502 (2020)

Ab Initio Nuclear Thermodynamics

Bing-Nan Lu^a, Ning Li^a, Serdar Elhatisari^b, Dean Lee^c, Joaquín E. Drut^d, Timo A. Lähde^e, Evgeny Epelbaum^f, and Ulf-G. Meißner^{g,h,7}

^aFacility for Rare Isotope Beams and Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA

^bFaculty of Engineering, Karamanoğlu Mehmetbey University, Karaman 70100, Turkey

^cDepartment of Physics and Astronomy, University of North Carolina, Chapel Hill, North Carolina 27599-3255, USA

^dInstitute for Advanced Simulation, Institut für Kernphysik, and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany

^eRuhr-Universität Bochum, Fakultät für Physik und Astronomie, Institut für Theoretische Physik II, D-44780 Bochum, Germany

^fHelmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany

^gTbilisi State University, 0186 Tbilisi, Georgia

(Received 11 April 2020; revised 6 August 2020; accepted 29 September 2020; published 3 November 2020)

We propose a new Monte Carlo method called the pinhole trace algorithm for *ab initio* calculations of the thermodynamics of nuclear systems. For typical simulations of interest, the computational speedup relative to conventional grand-canonical ensemble calculations can be as large as a factor of one thousand. Using a leading-order effective interaction that reproduces the properties of many atomic nuclei and neutron matter to a few percent accuracy, we determine the location of the critical point and the liquid-vapor coexistence line for symmetric nuclear matter with equal numbers of protons and neutrons. We also present the first *ab initio* study of the density and temperature dependence of nuclear clustering.

- new pinhole trace algorithm
 - liquid-vapor phase transition
 - location of the critical point

Phys. Lett. B 850 (2024) 138463

Contents lists available at ScienceDirect
Physics Letters B
journal homepage: www.elsevier.com/locate/physletb

 ELSEVIER

Check for updates

Letter

Ab initio study of nuclear clustering in hot dilute nuclear matter

Zhengxue Ren^{a,b,*}, Serdar Elhatisari^{c,b}, Timo A. Lähde^{a,d}, Dean Lee^c, Ulf-G. Meißner^{b,a,f}

^a Institut für Kernphysik, Institute for Advanced Simulation and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany
^b Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany
^c Faculty of Natural Sciences and Engineering, Gaziantep Islam Science and Technology University, Gaziantep 27010, Turkey
^d Center for Advanced Simulation and Analytics (CASAS), Forschungszentrum Jülich, D-52425 Jülich, Germany
^e Facility for Rare Isotope Beams and Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
^f Tbilisi State University, 0186 Tbilisi, Georgia

ARTICLE INFO

Editor: A. Schwenk

ABSTRACT

We present a systematic *ab initio* study of clustering in hot dilute nuclear matter using nuclear lattice effective field theory with an SU(4)-symmetric interaction. We introduce a method called light-cluster distillation to determine the abundances of dimers, trimers, and alpha clusters as a function of density and temperature. Our lattice results are compared with an ideal gas model composed of free nucleons and clusters. Excellent agreement is found at very low density, while deviations from ideal gas abundances appear at increasing density due to cluster-nucleon and cluster-cluster interactions. In addition to determining the composition of hot dilute nuclear matter as a function of density and temperature, the lattice calculations also serve as benchmarks for virial expansion calculations, statistical models, and transport models of fragmentation and clustering in nucleus-nucleus collisions.

- new light cluster distillation method
 - abundances of dimers, trimers, tetramers
 - benchmark for virial calculations

Chiral Interactions at N3LO: Foundations

Towards precision calculations of heavy nuclei

- Groundbreaking work (Hoyle state, α - α scattering, ...) done at N2LO
 - precision limited, need to go to N3LO

- Two step procedure:

- 1) Further improve the LO action

- minimize the sign oscillations
- minimize the higher-body forces
- essentially done ✓ → as just discussed

- 2) Work out the corrections to N3LO


- first on the level of the NN interaction ✓
- new important technique: **wave function matching** ✓
- second for the spectra/radii/... of nuclei (first results) ✓
- third for nuclear reactions/astrophysics (first results) ✓

NN interaction at N3LO

42

Li et al., Phys. Rev. C **98** (2018) 044002; Phys. Rev. C **99** (2019) 064001

- np phase shifts including uncertainties for $a = 1.32$ fm (cf. Nijmegen PWA)




uncertainty estimates à la Epelbaum, Krebs, UGM,
Eur. Phys. J. A **51** (2015) 53

Wave function matching I

43

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]]

- Graphical representation of w.f. matching



- W.F. matching is a “Hamiltonian translator”: eigenenergies from H_1 but w.f. from $H_2 = U^\dagger H_1 U$

Wave function matching II

44

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]]

- \mathbf{H}_{soft} has tolerable sign oscillations, good for many-body observables
- \mathbf{H}_χ has severe sign oscillations, derived from the underlying theory
→ can we find a unitary trafo, that creates a chiral \mathbf{H}_χ that is pert. th'y friendly?

$$\mathbf{H}'_\chi = \mathbf{U}^\dagger \mathbf{H}_\chi \mathbf{U}$$

- Let $|\psi_{\text{soft}}^0\rangle$ be the lowest eigenstate of \mathbf{H}_{soft}
- Let $|\psi_\chi^0\rangle$ be the lowest eigenstate of \mathbf{H}_χ
- Let $|\phi_{\text{soft}}\rangle$ be the projected and normalized lowest eigenstate of \mathbf{H}_{soft}

$$|\phi_{\text{soft}}\rangle = \mathcal{P} |\psi_{\text{soft}}^0\rangle / ||\psi_{\text{soft}}^0\rangle||$$

- Let $|\phi_\chi\rangle$ be the projected and normalized lowest eigenstate of \mathbf{H}_χ

$$|\phi_\chi\rangle = \mathcal{P} |\psi_\chi^0\rangle / ||\psi_\chi^0\rangle||$$

$$\hookrightarrow U_{R',R} = \theta(r - R)\delta_{R',R} + \theta(R' - r)\theta(R - r)|\phi_\chi^\perp\rangle\langle\phi_{\text{soft}}^\perp|$$

Chiral Interactions at N3LO: Applications to nuclear structure

Wave function matching for light nuclei

46

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]], L. Bovermann, PhD thesis

- W.F. matching for the light nuclei


Nucleus	B_{LO} [MeV]	$B_{\text{N}3\text{LO}}$ [MeV]	Exp. [MeV]
$E_{\chi,d}$	1.79	2.21	2.22
$\langle \psi_{\text{soft}}^0 H_{\chi,d} \psi_{\text{soft}}^0 \rangle$	0.45	0.62	
$\langle \psi_{\text{soft}}^0 H'_{\chi,d} \psi_{\text{soft}}^0 \rangle$	1.65	2.01	
$\langle \psi_{\text{soft}}^0 H_{\chi,t} \psi_{\text{soft}}^0 \rangle$	5.96(8)	5.91(9)	8.48
$\langle \psi_{\text{soft}}^0 H'_{\chi,t} \psi_{\text{soft}}^0 \rangle$	7.97(8)	8.72(9)	
$\langle \psi_{\text{soft}}^0 H_{\chi,\alpha} \psi_{\text{soft}}^0 \rangle$	24.61(4)	23.84(14)	28.30
$\langle \psi_{\text{soft}}^0 H'_{\chi,\alpha} \psi_{\text{soft}}^0 \rangle$	27.74(4)	29.21(14)	

- reasonable accuracy for the light nuclei

- Tjon-band recovered with H'_{χ}

Platter, Hammer, UGM, Phys. Lett. B **607** (2005) 254

→ now let us go to larger nuclei....




Nuclei at N3LO

47

- Binding energies of nuclei for $a = 1.32$ fm: Determining the 3NF LECs

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]]




→ excellent starting point for precision studies

Prediction: Charge radii at N3LO

48

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]]

- Charge radii ($a = 1.32$ fm, statistical errors can be reduced)




Prediction: Neutron & nuclear matter at N3LO

49

Elhatisari et al., acc. for publication in ... [arXiv:2210.17488 [nucl-th]]

- EoS of pure neutron matter & nuclear matter ($a = 1.32 \text{ fm}$)




→ can be improved using twisted b.c.'s

Prediction: Isotope chains of carbon & oxygen

50

NLEFT collaboration, in progress

- Towards the neutron drip-line in carbon and oxygen:




→ 3NFs of utmost importance for the n-rich isotopes!

Prediction: Be isotopes

51

Shen, ..., NLEFT collaboration, in progress

- Systematic study of the Be isotopes & their em transitions:



→ SU(4) works amazingly well, but clear deviations
→ N3LO works pretty well, few small deviations

Prediction: Triton β -decay at N3LO

52

Elhatisari, Hildenbrand, UGM, in preparation

- Master formula: $(1 + \delta_R) t_{1/2} f_V = \frac{K/G_V^2}{\langle F \rangle^2 + \frac{f_A}{f_V} g_A^2 \langle GT \rangle^2}$

- Experiment: $\langle F \rangle = \sum_{n=1}^3 \langle {}^3\text{He} || \tau_{n,+} || {}^3\text{H} \rangle = 0.9998$ [theory!]

$$\langle GT \rangle = \sum_{n=1}^3 \langle {}^3\text{He} || \sigma_n \tau_{n,+} || {}^3\text{H} \rangle = 1.6474(23)$$



- NLEFT:

$$\langle F \rangle_{\text{N3LO}} = 0.9992(16)$$

$$\langle GT \rangle_{\text{N3LO}} = 1.661(35)$$

- No Coulomb at LO

- Larger L underway...



Chiral Interactions at N3LO: Applications to scattering

Scattering: Methods I

54

- The time-honored Lüscher approach:

Lüscher, Commun. Math. Phys. **105** (1986) 153; Nucl. Phys. B **354** (1991) 531

Phase shifts from the volume dependence of the energy levels

→ works in many cases, problems w/ partial-wave mixing and cluster-cluster scattering

- Spherical wall technique:

impose spherical b.c.'s on the lattice


Carlson et al., Nucl. Phys. A **424** (1984) 47; Borasoy et al., Eur. Phys. J. A **34** (2007) 185

→ not too small lattices, partial-wave mixing under control

- Improved spherical wall method:

Lu, Lähde, Lee, UGM, Phys. Lett. B **760** (2016) 309

- perform angular momentum projection
 - impose an auxiliary potential behind R_{wall}
- much improved precision




Scattering: Methods II

- Adiabatic projection method :

Rupak, Lee, Phys. Rev. Lett. **111** (2013) 032502; Pine, Lee, Rupak, Eur. Phys. J. A **49** (2013) 151;
Elhatisari et al., Eur. Phys. J. A **52** (2016) 174;
 - Construct a low-energy effective theory for clusters
 - Use initial states parameterized by the relative separation between clusters
 - project them in Euclidean time with the chiral EFT Hamiltonian \mathbf{H}
- $$|\vec{R}\rangle = \sum_{\vec{r}} |\vec{r} + \vec{R}\rangle \otimes \vec{r}$$
- $$|\vec{R}\rangle_\tau = \exp(-\mathbf{H}\tau)|\vec{R}\rangle$$
- “dressed cluster states” (polarization, deformation, Pauli)
- Adiabatic Hamiltonian (requires norm matrices)

$$[H_\tau]_{\vec{R}\vec{R}'} = \tau \langle \vec{R} | H | \vec{R}' \rangle_\tau$$




Scattering: Neutron-deuteron scattering at N3LO

56

Elhatisari, Hildenbrand, UGM, in progress

- Use Lüscher's method to calculate spin doublet n - d scattering





→ shows good convergence

Scattering: Neutron-alpha scattering at N3LO

57

Elhatisari, Hildenbrand, UGM, in progress

- Use Lüscher's method to calculate $n\text{-}\alpha$ scattering




- R-matrix results from G. Hale, [private communication](#)
 - ↪ Some fine-tuning of three-body forces for $^2P_{1/2}$ needed

Scattering: Alpha-carbon scattering at N3LO


58

Elhatisari, Hildenbrand, UGM, ... NLEFT, in progress

- Use the APM, first step for the holy grail of nuclear astrophysics
→ different Euclidean times & different initial states



Plaga et al., Nucl. Phys. A 465 (1987) 291

 $\psi_A \sim {}^{16}\text{O}, \psi_B \sim {}^{12}\text{C} + {}^4\text{He}$

Summary & outlook

- Nuclear lattice simulations: a new quantum many-body approach
 - based on the successful continuum nuclear chiral EFT
 - a number of highly visible results already obtained
- Recent developments
 - highly improved LO action based on SU(4)
 - ↪ a number of interesting application (^{12}C , $^4\text{He}, \dots$)
 - ↪ towards the neutron matter EoS at high densities
 - more in the talk by Bing-Nan Lu
 - NN interaction at N3LO w/ wave function matching
 - ↪ first promising results for nuclear structure and scattering
 - ↪ hyper-nuclei are under investigation
 - more in the talk by Dean Lee

SPARES

The hidden spin-isospin exchange symmetry

Nucleon-nucleon interaction in large- N_C

Kaplan, Savage, Phys. Lett. **365B** (1996) 244; Kaplan, Manohar, Phys. Rev. **C 56** (1997) 96

- Performing the large- N_C analysis:

$$V_{\text{large}-N_c}^{\text{2N}} = V_C + W_S \vec{\sigma}_1 \cdot \vec{\sigma}_2 \vec{\tau}_1 \cdot \vec{\tau}_2 + W_T S_{12} \vec{\tau}_1 \cdot \vec{\tau}_2 + \dots$$

- Leading terms are $\sim N_C$
- First corrections are $1/N_C^2$ suppressed, fairly strong even for $N_C = 3$
- Velocity-dependent corrections can be incorporated
- Based on spin-isospin exchange symmetry of the nucleon w.f. $d_\uparrow \leftrightarrow u_\downarrow$ or on the nucleon level $n_\uparrow \leftrightarrow p_\downarrow$
- Constraints on 3NFs: Phillips, Schat, PRC **88** (2013) 034002; Epelbaum et al., EPJA **51** (2015) 26

Hidden spin-isospin symmetry: Basic ideas

63

Lee, Bogner, Brown, Elhatisari, Epelbaum, Hergert, Hjorth-Jensen, Krebs, Li, Lu, UGM, Phys. Rev. Lett. **127** (2021) 062501 [2010.09420 [nucl-th]]

- $V_{\text{large}-N_c}^{2N}$ is not renormalization group invariant: $\frac{dV_\mu(p, p')}{d\mu} \neq 0$
 \simeq implicit setting of a preferred renormalization/resolution scale
- How does this happen?
 - **high energies:** corrections to the nucleon w.f. are $\sim v^2$
 - these high-energy modes must be $\mathcal{O}(1/N_C^2)$ in our low-energy EFT
 - momentum resolution scale $\Lambda \sim m_N/N_C \sim \mathcal{O}(1)$
 - consistent with the cutoff in a Δ less th'y $\sim \sqrt{2m_N(m_\Delta - m_N)}$
 - **low energies:** the resolution scale must be large enough,
 - so that orbital angular momentum and spin are fully resolved
 - as nucleon size is independent of N_C , so should be Λ ✓
- as will be shown, the optimal scale (where corrections are $\sim 1/N_C^2$) is:

$$\Lambda_{\text{large}-N_c} \simeq 500 \text{ MeV}$$

Nucleon-nucleon phase shifts – lattice

Lee, Bogner, Brown, Elhatisari, Epelbaum, Hergert, Hjorth-Jensen, Krebs, Li, Lu, UGM,
 Phys. Rev. Lett. **127** (2021) 062501 [2010.09420 [nucl-th]]


- Use N3LO action (w/ TPE absorbed in contact interactions) at $a = 1.32 \text{ fm}$

$$\hookrightarrow \Lambda = \pi/a = 470 \text{ MeV}$$

- compare $S = 0, T = 1$ w/ $S = 1, T = 0$
- S-waves: switch off the tensor force in 3S_1
- D-waves: average the spin-triplet channel
- NLEFT low-energy constants



ch., order	LEC (l.u.)	ch., order	LEC (l.u.)
$^1S_0, Q^0$	1.45(5)	$^3S_1, Q^0$	1.56(3)
$^1S_0, Q^2$	-0.47(3)	$^3S_1, Q^2$	-0.53(1)
$^1S_0, Q^4$	0.13(1)	$^3S_1, Q^4$	0.12(1)
$^1D_2, Q^4$	-0.088(1)	$^3D_{\text{all}}, Q^4$	-0.070(2)

⇒ works pretty well



Nucleon-nucleon phase shifts – continuum

- Consider various (chiral) continuum potentials → also works ✓



- IDAHO N3LO
- — — IDAHO N4LO ($\Lambda = 500$ MeV)
- — • CD-Bonn
- Bochum N4⁺LO ($\Lambda = 400 - 550$ MeV)
- • • Nijmegen PWA

- Entem, Machleidt, PRC **68** (2003) 041001
- Entem, Machleidt, Nosyk PRC **96** (2017) 024004
- Machleidt, PRC **63** (2001) 024001
- Reinert, Krebs, Epelbaum, EPJA **54** (2018) 86
- Wiringa, Stoks, Schiavilla, PRC **51** (1995) 38

Two-nucleon matrix elements

- Consider the ME between any two-nucleon states A and B . Both have total spin S and total isospin T . Then (for isospin-inv. H):

$$M(S, T) = \frac{1}{2S+1} \sum_{S_z=-S}^S \langle A; S, S_z; T, T_z | H | B; S, S_z; T, T_z \rangle$$

- Spin-isospin exchange symmetry: $M(S, T) = M(T, S)$
- Ex: ${}^{30}\text{P}$ has 1 proton + 1 neutron in the $1s_{1/2}$ orbitals (minimal shell model)
 → if spin-isospin exchange symmetry were exact, the $S = 0, T = 1$ & $S = 1, T = 0$ states should be degenerate
- Data: The 1^+ g.s. is 0.677 MeV below the 0^+ excited state ($E_{g.s.} \simeq 220$ MeV)
 → fairly good agreement, consistent w/ $1/N_C^2$ corrections
 → explanation: interactions of the np pair with the ${}^{28}\text{Si}$ core are suppressing spatial correlations of the 1^+ w.f. caused by the tensor interaction

Two-nucleon matrix elements in the s-d shell

- Test the spin-isospin exchange symmetry for general two-body MEs 1s-0d shell
- Use the spin-tensor analysis developed by Kirson, Brown et al.
Kirson, PLB **47** (1973) 110; Brown et al., JPhysG **11** (1985) 1191; Ann. Phys. **182** (1988) 191
- Seven two-body MEs for $(S, T) = (1, 0)$ and $(S, T) = (0, 1)$

ME	L_1	L_2	L_3	L_4	L_{12}	L_{34}
1	2	2	2	2	0	0
2	2	2	2	2	2	2
3	2	2	2	2	4	4
4	2	2	2	0	2	2
5	2	2	0	0	0	0
6	2	0	2	0	2	2
7	0	0	0	0	0	0

L_1, L_2 : orbital angular momenta of the outgoing orbitals of A

L_{12} : total angular momentum of state A

L_3, L_4 : orbital angular momenta of the outgoing orbitals of B

L_{34} : total angular momentum of state B









ME 7 corresponds to the $1s_{1/2}$ orbitals discussed before

set $L_Z = (L_{12})_Z = (L_{34})_Z$, average over L_Z

→ Work out $M(S, T)$ for various forces at $\Lambda = 2.0, 2.5, 3.0, 3.5 \text{ fm}^{-1}$

Two-nucleon matrix elements in the s-d shell

- Results for the AV18 and N3LO chiral potentials



Two-nucleon matrix elements: Conclusions

- As anticipated:
 - The optimal resolution scale is obviously $\Lambda \sim 500$ MeV
 - For $\Lambda < \Lambda_{\text{large-}N_c}$, the $(S, T) = (1, 0)$ channel is more attractive
 - For $\Lambda > \Lambda_{\text{large-}N_c}$, the $(S, T) = (0, 1)$ channel is more attractive
 - These results do not depend on the type of interaction,
while AV18 is local, chiral N3LO has some non-locality
(and similar for more modern interactions like chiral N4⁺LO)

↪ consistent with the results for NN scattering

⇒ Validates Weinberg's power counting! ✓

Three-nucleon forces

70

- Leading central three-nucleon force at the optimal resolution scale:

$$\begin{aligned} V_{\text{large}-N_c}^{\text{3N}} &= V_C^{\text{3N}} + [(\vec{\sigma}_1 \times \vec{\sigma}_2) \cdot \vec{\sigma}_3] [(\vec{\tau}_1 \times \vec{\tau}_2) \cdot \vec{\tau}_3] W_{123}^{\text{3N}} \\ &+ \vec{\sigma}_1 \cdot \vec{\sigma}_2 \vec{\tau}_1 \cdot \vec{\tau}_2 W_{12}^{\text{3N}} + \vec{\sigma}_2 \cdot \vec{\sigma}_3 \vec{\tau}_2 \cdot \vec{\tau}_3 W_{23}^{\text{3N}} \\ &+ \vec{\sigma}_3 \cdot \vec{\sigma}_1 \vec{\tau}_3 \cdot \vec{\tau}_1 W_{31}^{\text{3N}} + \dots, \end{aligned}$$

- Subleading central 3N interactions are of size $1/N_C$, of type

$$\vec{\sigma}_1 \cdot \vec{\sigma}_2 [(\vec{\tau}_1 \times \vec{\tau}_2) \cdot \vec{\tau}_3], \quad [(\vec{\sigma}_1 \times \vec{\sigma}_2) \cdot \vec{\sigma}_3] \vec{\tau}_1 \cdot \vec{\tau}_2$$

⇒ helps in constraining the many short-range three-nucleon interactions that appear at higher orders in chiral EFT

- The spin-isospin exchange symmetry of the leading interactions also severely limits the isospin-dependent contributions of the 3N interactions to the nuclear EoS
- ⇒ relevant for calculations of the nuclear symmetry energy and its density dependence in dense nuclear matter

Ab Initio Nuclear Thermodynamics


B. N. Lu, N. Li, S. Elhatisari, D. Lee, J. Drut, T. Lähde, E. Epelbaum, UGM,
Phys. Rev. Lett. **125** (2020) 192502 [arXiv:1912.05105]

Phase diagram of strongly interacting matter

72

- Sketch of the phase diagram of strongly interacting matter

Fig. courtesy B.-N. Lu




Pinhole trace algorithm (PTA)

- The pinhole states span the whole A-body Hilbert space
- The canonical partition function can be expressed using pinholes:

$$Z_A = \text{Tr}_A [\exp(-\beta H)] , \quad \beta = 1/T$$

$$= \sum_{n_1, \dots, n_A} \int \mathcal{D}s \mathcal{D}\pi \langle n_1, \dots, n_A | \exp[-\beta H(s, \pi)] | n_1, \dots, n_A \rangle$$

- allows to study: liquid-gas phase transition → this talk
thermodynamics of finite nuclei
thermal dissociation of hot nuclei
cluster yields of dissociating nuclei




New paradigm for nuclear thermodynamics

- The PTA allows for simulations with fixed neutron & proton numbers at non-zero T
 ↳ thousands to millions times faster than existing codes using the grand-canonical ensemble ($t_{\text{CPU}} \sim VN^2$ vs. $t_{\text{CPU}} \sim V^3N^2$)
- Only a mild sign problem → pinholes are dynamically driven to form pairs
- Typical simulation parameters:
 up to $N = 144$ nucleons in volumes $L^3 = 4^3, 5^3, 6^3$
 ↳ densities from $0.008 \text{ fm}^{-3} \dots 0.20 \text{ fm}^{-3}$
 $a = 1.32 \text{ fm} \rightarrow \Lambda = \pi/a = 470 \text{ MeV}$, $a_t \simeq 0.1 \text{ fm}$
 consider $T = 10 \dots 20 \text{ MeV}$
- use twisted bc's, average over twist angles → acceleration to the td limit
- very favorable scaling for generating config's: $\Delta t \sim N^2 L^3$


Chemical potential

- Calculated from the free energy: $\mu = (F(N+1) - F(N-1))/2$



Equation of state

- Calculated by integrating: $dP = \rho d\mu$
- Critical point: $T_c = 15.8(1.6)$ MeV, $P_c = 0.26(3)$ MeV/fm 3 , $\rho_c = 0.089(18)$ fm $^{-3}$




Experiment: $T_c = 15.0(3)$ MeV, $P_c = 0.31(7)$ MeV/fm 3 , $\rho_c = 0.06(2)$ fm $^{-3}$

Vapor-liquid phase transition

77

- Vapor-liquid phase transition in a finite volume V & $T < T_c$
- the most probable configuration for different nucleon number A
- the free energy
- chemical potential $\mu = \partial F / \partial A$




CENTER-of-MASS PROBLEM

- AFQMC calculations involve states that are superpositions of many different center-of-mass (com) positions

$$Z_A(\tau) = \langle \Psi_A(\tau) | \Psi_A(\tau) \rangle$$

$$|\Psi_A(\tau)\rangle = \exp(-H\tau/2)|\Psi_A\rangle$$



- but: translational invariance requires summation over all transitions

$$Z_A(\tau) = \sum_{i_{\text{com}}, j_{\text{com}}} \langle \Psi_A(\tau, i_{\text{com}}) | \Psi_A(\tau, j_{\text{com}}) \rangle, \quad \text{com} = \text{mod}((i_{\text{com}} - j_{\text{com}}), L)$$

i_{com} (j_{com}) = position of the center-of-mass in the final (initial) state

- density distributions of nucleons can not be computed directly, only moments
- need to overcome this deficiency

PINHOLE ALGORITHM

- Solution to the CM-problem:
track the individual nucleons using the *pinhole algorithm*

- Insert a screen with pinholes with spin & isospin labels that allows nucleons with corresponding spin & isospin to pass = insertion of the A-body density op.:

$$\rho_{i_1, j_1, \dots, i_A, j_A}(\mathbf{n}_1, \dots, \mathbf{n}_A) \\ = : \rho_{i_1, j_1}(\mathbf{n}_1) \cdots \rho_{i_A, j_A}(\mathbf{n}_A) :$$

- MC sampling of the amplitude:

$$A_{i_1, j_1, \dots, i_A, j_A}(\mathbf{n}_1, \dots, \mathbf{n}_A, L_t) \\ = \langle \Psi_A(\tau/2) | \rho_{i_1, j_1, \dots, i_A, j_A}(\mathbf{n}_1, \dots, \mathbf{n}_A) | \Psi_A(\tau/2) \rangle$$

- Allows to measure proton and neutron distributions
- Resolution scale $\sim a/A$ as cm position \mathbf{r}_{cm} is an integer n_{cm} times a/A

