

The 23rd International Conference on Few-Body Problems in Physics

$K\bar{K}^*$ and $D\bar{D}^*$ resonance states in the Non-Hermitian Quantum Mechanics

Bao-Xi Sun

Beijing University of Technology

2024.09.25

1. Content

- 1. The Schrodinger equation in a rectangular potential well.
- 2. The Schrodinger equation in a Yukawa potential.
- 3. The $K\bar{K}^*$ interaction: f1(1285) and f1(1420) .
- 4. The $D\bar{D}^*$ interaction: X(3872) and Zc(3900) and X(3940)
- 5. Summary

1. Rectangular potential-well

The rectangular potential-well is defined by

$$V(x) = \begin{cases} 0 & x \notin (0, L), \\ -V_0 & x \in (0, L), \end{cases}$$

with the depth of the well $V_0 > 0$.

The Schrodinger equation can be written as:

$$\begin{cases} \varphi'' + k_0^2 \varphi = 0, & k_0 = \sqrt{2ME} / \hbar & x \notin (0, L), \\ \varphi'' + k^2 \varphi = 0, & k = \sqrt{2M(E + V_0)} / \hbar & x \in (0, L), \end{cases}$$

1. Rectangular potential-well

The general solution is

$$\begin{cases} \varphi = Ce^{ik_0x} + C'e^{-ik_0x} & (x < 0), \\ \varphi = Ae^{ikx} + Be^{-ikx} & (0 < x < L), \\ \varphi = De^{ik_0x} + D'e^{-ik_0x} & (x > L), \end{cases}$$

Incoming wave condition: $C' = D = 0$

Outgoing wave condition: $C = D' = 0$

The transcendental equation is

$$2 \cot k2L = \frac{\pm i(k^2 + k_0^2)}{kk_0}.$$

Positive(negative) sign represents the outgoing(incoming) wave.

1. Rectangular potential-well

The mechanism of transition from a bound state to a resonance state becomes evident when two variables are introduced.

$$\alpha = \sqrt{1 + \frac{E}{V_0}}, \quad \gamma = \sqrt{\frac{2MV_0L^2}{\hbar^2}}, \quad V_0 > 0.$$


Bound states:

$$\gamma\alpha = (n-1)\pi + 2\cos^{-1}\alpha, \quad n = 1, 2, \dots$$

Virtual states:

$$\gamma\alpha = (n-1)\pi + 2\cos^{-1}\sqrt{1-\alpha^2}, \quad n = 1, 2, \dots$$

1. Rectangular potential-well



2. One-Pion Exchange potential

The one-pion-exchange potential is:

$$V(x) = -g^2 \frac{e^{-mr}}{d},$$

where the distance in the denominator has been replaced with the range of force, $d = 1/m$.

Supposing the radial wave function $R(r) = \frac{u(r)}{r}$, the radial Schrodinger equation becomes

$$-\frac{\hbar^2}{2\mu} \frac{d^2 u(r)}{dr^2} + V(r)u(r) = Eu(r),$$

Sun et al., CTP, 76, 105301, 2024

2. One-Pion Exchange potential

With the variable substitution:

$$x = \alpha e^{-\beta r}, \quad u(r) = J(x), \quad \text{with } \alpha = 2g\sqrt{2\mu d}, \quad \beta = \frac{1}{2d}, \quad \rho^2 = -8d^2\mu E,$$

The Schrodinger equation becomes the Bessel eq.

$$\frac{d^2 J(x)}{dx^2} + \frac{1}{x} \frac{dJ(x)}{dx} + \left(1 - \frac{\rho^2}{x^2}\right) J(x) = 0,$$

$$r \rightarrow 0, \quad u(r) \rightarrow 0, \quad J_\rho(x) = J_\rho(\alpha) = 0,$$

If only one bound state exists, and the binding energy is given, the value of $\rho^2 = -8d^2\mu E$ can be determined, and the coupling $g^2 = \frac{\alpha^2}{8\mu d}$ is obtained with the first zero point of $J_\rho(\alpha)$.

2. One-Pion Exchange potential

The Hankel functions are also solutions of the Bessel function. In a scattering process, the general solution of the Bessel equation is $u(r) = DH_{\rho'}^{(1)}(x) + D'H_{\rho'}^{(2)}(x)$,

Incoming wave condition: $r \rightarrow 0$, $u(r) \rightarrow 0$, $H_{\rho'}^{(1)}(x) = H_{\rho'}^{(1)}(\alpha) = 0$,

Outgoing wave condition: $r \rightarrow 0$, $u(r) \rightarrow 0$, $H_{\rho'}^{(2)}(x) = H_{\rho'}^{(2)}(\alpha) = 0$,

With the same zero point (coupling constant), the order of $H_{\rho'}^{(2)}(\alpha)$ is determined, which is related to the energy of the resonance state,

$$E = (M - M_{Threshold}) - i \frac{\Gamma}{2} = -\frac{\rho'^2}{8d^2 \mu}.$$

3. $f_1(1285)$ or $f_1(1420)$?

1. [Roca, Oset, Singh, PRD, 72, 014002, 2005](#)

$f_1(1285)$ is a $K\bar{K}^*$ bound state.

2. [Wan, Zhao, Sun, 1808.08358\[hep-ph\]](#)

$f_1(1420)$ is a $K\bar{K}^*$ resonance state, no other pole is detected.


3. [Debastiani, Aceti, Liang, Oset, PRD, 95, 034015, 2017](#)

$f_1(1420)$ is related to a $K^*\bar{K}K$ triangle singularity.

3. KK^* interaction

In the unitary coupled-channel approach, the vector meson transfer is dominant, while the pion exchange is neglected.

Wan, Zhao, Sun, 1808.08358[hep-ph]





Actually, the OPEP is critical in the generation of resonance states.

3. KK^* interaction

The $f_1(1285)$ particle is treated as a $K\bar{K}^*$ bound state with a binding energy of 105MeV, and the coupling constant is determined as $g=1.682$.


Energy	Name	$I^G(J^{PC})$	Mass	Width
1417-i18	$f_1(1420)$	$0^+(1^{++})$	1426.3 ± 0.9	54.5 ± 2.6



4. DD^* interaction

In the unitary coupled-channel approach, the vector meson transfer is dominant, while the pion exchange is neglected.


Sun, Wan, Zhao, *CPC*, 42, 053105, (2018)



Actually, the OPEP is critical in the generation of resonance states.


4. DD^* interaction

The X(3872) particle is assumed to be a $D\bar{D}^*$ bound state with a zero binding energy since it lies at the $D^0\bar{D}^{*0}$ threshold. Therefore, the coupling constant is determined as $g=0.323$ according to the first zero point $\alpha=2.405$ of $J_0(x)$.



4. DD^* interaction

If the $X(3872)$ particle is a $D\bar{D}^*$ bound state, $Z_c(3900)$ would be a $D\bar{D}^*$ resonance state. All states are isospin degenerate.



$D\bar{D}^*$	Energy	Name	$I^G(J^{PC})$	Mass	Width
1	3885-i1	$Z_c(3900)$	$1^+(1^{+-})$	3887.1	28.4
2	4029-i108	$X(3940)$	$?^?(?^{??})$	3942	37
3	4328-i191	$\chi_{c1}(4274)$	$0^+(1^{++})$	4286	51
4	4772-i267	$\chi_{c1}(4685)$	$0^+(1^{++})$	4684	126

4. DD* interaction

The resonance state at 4029-i108MeV might correspond to the X(3940) particle, which is predicted as a partner state of X(3872) with $J^{PC} = 1^{++}$.

Y. S. Kalashnikova, PRD, 72, 034010, 2005

P. G. Ortega, PRD, 81, 054023, 2010

Zhou and Xiao, PRD, 96, 054031, 2017

Q. Deng et al., 2312.10296[hep-ph].


H. Li et al., 2402.14541[heplat].

F. Giacosa et al., IJMPA, 34, 1950173, 2019.

G. J. Wang et al., 2306.12406[hep-ph].

4. DD^* interaction

If the $X(3872)$ particle is a D^+D^{*-} / D^-D^{*+} bound state, the binding energy is about 8.11 MeV, then the coupling constant $g = 0.6520$. Two resonance states are detected.



5. Summary

B. X. Sun et al., Commun. Theor. Phys., 76, 105301, 2023

1. The interactions of hadrons are studied in the non-Hermitian quantum mechanics. By solving the Schrodinger equation, some resonance states are obtained when the outgoing wave condition is considered.

2. OPEP is critical in the formation of $K\bar{K}^*$ and $D\bar{D}^*$ bound and resonanced states.