

Effective range expansion with the left-hand cut

Meng-Lin Du (UESTC)

University of Electronic Science and Technology of China


In collaboration with A. Fillin, V. Baru, X.-K. Dong, E. Epelbaum, F.-K. Guo, C. Hanhart, A. Nefediev, J. Nieves, Q. Wang, and B. Wu

Based on PRL 131,131903 (2023), and arXiv:2408.09375[hep-ph]

Sep. 26, 2024 @ Beijing, China

The 23rd International Conference on Few-Body Problems in Physics (FB23)

Doubly charmed tetraquark (Tcc)




Breit-Wigner fit

LHCb, Nature Phys. 18, (2022) 751

Parameter	Value
N	117 ± 16
δm_{BW}	$-273 \pm 61 \text{ keV}$
Γ_{BW}	$410 \pm 165 \text{ keV}$

☞ $\Re \sim 400 \text{ keV}$.



Unitarized and analytical

LHCb, Nature Commun. 13 (2022), 3351

$$\delta m = m_{T_{cc}^+} - m_{D^{*+}} - m_{D^0}$$

$$\delta m_{\text{pole}} = -360 \pm 40_{-0}^{+4} \text{ keV}$$


$$\Gamma_{\text{pole}} = 48 \pm 2_{-14}^{+0} \text{ keV}$$

☞ $I = 0$: isoscalar

↪ $D^+ D^0 \pi^+, D^+ D^+$ ✗


☞ T_{cc}^+ resides near $D^* D$ thresholds

↪ approximate 90% of $D^0 D^0 \pi^+$ events contain a D^{*+} .





LHCb, Nature Commun. 13 (2022)

The three-body cut



Three-body cuts



LO Chiral Lagrangian (g determined from $D^* \rightarrow D\pi$)

$$\mathcal{L} = \frac{1}{4} g \text{Tr} (\vec{\sigma} \cdot \vec{u}_{ab} H_b H_a^\dagger)$$


$$U_\alpha(M, p) = P_\alpha - \sum_\beta \int \frac{d^3 \vec{q}}{(2\pi)^3} V_{\alpha\beta}(M, p, q) G_\beta(M, q) U_\beta(M, q)$$

$$\hookrightarrow G_\alpha(M, p) = \frac{1}{m_\alpha^* + m_\alpha + \frac{p^2}{2\mu_\alpha} - M - \frac{i}{2} \Gamma_\alpha(M, p)}$$

Doubly Charm Tetraquark on the Lattice



Padmanath *et al*, PRL129,032002(2022)

	m_D (MeV)	m_{D^*} (MeV)	M_{av} (MeV)	$a_{l=0}^{(J=1)}$ (fm)	$r_{l=0}^{(J=1)}$ (fm)	$\delta m_{T_{cc}}$ (MeV)	T_{cc}
Lattice ($m_\pi \simeq 280$ MeV, $m_c^{(h)}$)	1927(1)	2049(2)	3103(3)	1.04(29)	$0.96^{(+0.18)}_{(-0.20)}$	$-9.9^{+3.6}_{-7.2}$	Virtual bound st.
Lattice ($m_\pi \simeq 280$ MeV, $m_c^{(l)}$)	1762(1)	1898(2)	2820(3)	0.86(0.22)	$0.92^{(+0.17)}_{(-0.19)}$	$-15.0^{(+4.6)}_{(-9.3)}$	Virtual bound st.
Experiment [2,41]	1864.85(5)	2010.26(5)	3068.6(1)	-7.15(51)	$[-11.9(16.9), 0]$	-0.36(4)	Bound st.




$$t = \frac{E_{cm}}{2} \frac{1}{p \cot \delta - ip},$$

$$p \cot \delta = \frac{1}{a_0} + \frac{1}{2} r_0 p^2,$$



The three-body cut vs. left-hand cut





three-body cut

$$E > M_D + M_D + M_\pi$$




left-hand cut

$$\int_{-1}^1 d \cos \theta G_\pi(E, p, p)$$



$$G_\pi^{-1}(E, k, k') \xrightarrow[\text{on shell: } k=k'=p]{\cos \theta = \pm 1} E_{D^*}(p^2) - E_D(p^2) - \omega_\pi(4p^2/0) = 0$$

The left-hand cut



$$m_\pi = 280 \text{ MeV}$$

☞ two-body branch point:

$$E = M_D + M_{D^*}$$

$$\Rightarrow p_{\text{rhc}2}^2 = 0$$

☞ three-body branch point:

$$E = M_D + M_D + m_\pi$$

$$\Rightarrow \left(\frac{p_{\text{rhc}3}}{E_{DD^*}} \right)^2 = +0.019$$

☞ left-hand cut branch point:


$$\Rightarrow \left(\frac{p_{\text{lhc}}^{1\pi}}{E_{DD^*}} \right)^2 = -0.001$$

$$\left(\frac{\tilde{p}_{\text{lhc}}^{1\pi}}{E_{DD^*}} \right)^2 = -0.190$$

Phase shift with the left-hand cut: LSE

$$M_D = 1927 \text{ MeV}, M_{D^*} = 2049 \text{ MeV}, m_\pi = 280 \text{ MeV}$$

Du et al., PRL 131,131903 (2023)



Related recent works on FV w/ LHC...

Plane-wave basis to treat long-range interactions

Meng and Epelbaum, JHEP (2021)

Meng et al., PRD (2024)

Generalization of the Lüscher + K-matrix

Hansen and Raposo, JHEP (2024)

Three-body framework (automatically includes LHC)

Dawid et al., PRD (2023)

Hansen et al., PRD (2024)

Modify the Lüscher formula via “modified effective range expansion”


Bubna et al., JHEP (2024)

The N/D method

$$T(s) = \frac{N(s)}{D(s)}$$


$$\text{Im}D = \text{Im}\frac{N}{T} = N\text{Im}\frac{1}{T} = \begin{cases} -N\rho, & s > s_{\text{thr}} \\ 0, & s < s_{\text{thr}} \end{cases}$$

$$\text{Im}N = \begin{cases} \text{Im}TD, & s < s_{\text{lhc}} \\ 0, & s > s_{\text{lhc}} \end{cases}$$



$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + \sum_{m=0}^{n-1} a_m s^m - \frac{(s - s_0)^n}{\pi} \int_{s_{\text{thr}}}^{\infty} ds' \frac{\rho(s') N(s')}{(s' - s)(s' - s_0)^n},$$

$$N(s) = \sum_{m=0}^{n-\ell-1} b_m s^m + \frac{(s - s_0)^{n-\ell}}{\pi} \int_{-\infty}^{s_{\text{left}}} ds' \frac{\text{Im}T(s') D(s')}{(s' - s_0)^{n-\ell}(s' - s)}.$$

$N=1$ 

$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + P(s) + G(s)$$


$$T(s) = \frac{1}{D(s)}$$

The N/D method

$$T(s) = \frac{N(s)}{D(s)}$$


$$\text{Im}D = \text{Im}\frac{N}{T} = N\text{Im}\frac{1}{T} = \begin{cases} -N\rho, & s > s_{\text{thr}} \\ 0, & s < s_{\text{thr}} \end{cases}$$

$$\text{Im}N = \begin{cases} \text{Im}TD, & s < s_{\text{lhc}} \\ 0, & s > s_{\text{lhc}} \end{cases}$$



$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + \sum_{m=0}^{n-1} a_m s^m - \frac{(s - s_0)^n}{\pi} \int_{s_{\text{thr}}}^{\infty} ds' \frac{\rho(s') N(s')}{(s' - s)(s' - s_0)^n},$$

$$N(s) = \sum_{m=0}^{n-\ell-1} b_m s^m + \frac{(s - s_0)^{n-\ell}}{\pi} \int_{-\infty}^{s_{\text{left}}} ds' \frac{\text{Im}T(s') D(s')}{(s' - s_0)^{n-\ell}(s' - s)}.$$




$$\frac{1}{T_\ell^{\text{II}}} = \frac{1}{T_\ell} + 2i\rho \implies T^{\text{II}} = \frac{1}{\frac{D}{N} + 2i\rho} = \frac{N}{D + 2i\rho N}$$


Along the lhc, $i\rho$ and D is real, N has imaginary part.

$$D + 2i\rho N \neq 0$$

The left-hand cut



$$\text{Im } f(k^2) = c \text{Im } L(k^2) = -\frac{c}{4k^2} \pi, \quad \text{for } k^2 < k_{\text{thc}}^2$$



Solving LSE could be time-consuming.



For a t -channel exchange at low-energies, an S -wave amplitude reads

$$L_t(s) = \frac{1}{2} \int \frac{1}{t - m_5^2} d\cos\theta = -\frac{s}{\lambda(s, m_1^2, m_2^2)} \log \left(\frac{s - 2(m_1^2 + m_2^2) + m_5^2 + \frac{(m_1^2 - m_2^2)^2}{s}}{m_5^2} \right),$$

with m_5 the mass of changed particle. Likewise, the u -channel exchanged S -wave amplitude reads

$$L_u(s) = \frac{1}{2} \int \frac{1}{u - m_5^2} d\cos\theta = -\frac{s}{\lambda(s, m_1^2, m_2^2)} \left(\log(s + m_5^2 - 2(m_1 + m_2)^2) - \log(m_5^2 - \frac{(m_1^2 - m_2^2)^2}{s}) \right).$$

The left-hand cut: nonrelativistic



Exchanged-particle: relativistic

$$\eta = |m_1 - m_2|/(m_1 + m_2)$$

$$\mu_{\text{ex}}^2 = m_{\text{ex}}^2 - (m_1 - m_2)^2$$

$$\mu_+^2 = 4\mu\mu_{\text{ex}}^2/(m_1 + m_2)$$

$$L_t(k^2) \equiv \frac{1}{2} \int_{-1}^{+1} \frac{d \cos \theta}{t - m_{\text{ex}}^2} = -\frac{1}{4k^2} \log \frac{m_{\text{ex}}^2/4 + k^2}{m_{\text{ex}}^2/4},$$

$$L_u(k^2) \equiv \frac{1}{2} \int_{-1}^{+1} \frac{d \cos \theta}{u - m_{\text{ex}}^2} \approx -\frac{1}{4k^2} \log \frac{\mu_+^2/4 + k^2}{\mu_+^2/4 + \eta^2 k^2},$$

$$\frac{1}{2} \int_{-1}^{+1} \frac{(\mathbf{p}_1 - \mathbf{p}_3)^2}{t - m_{\text{ex}}^2} d \cos \theta = -\frac{m_{\text{ex}}^2}{2} \int_{-1}^{+1} \frac{d \cos \theta}{t - m_{\text{ex}}^2} - 1$$

$$\frac{1}{2} \int_{-1}^{+1} \frac{(\mathbf{p}_1 - \mathbf{p}_4)^2}{u - m_{\text{ex}}^2} d \cos \theta \approx -\frac{\mu_{\text{ex}}^2}{2} \int_{-1}^{+1} \frac{d \cos \theta}{u - m_{\text{ex}}^2} - 1$$

$\mathcal{F}_\ell/2$

$$f(k^2) = \frac{n(k^2)}{d(k^2)}$$

$$\begin{aligned} \text{Im } d(k^2) &= -k n(k^2), & \text{for } k^2 > 0, \\ \text{Im } n(k^2) &= d(k^2) \text{Im } f(k^2), & \text{for } k^2 < k_{\text{h.c.}}^2. \end{aligned}$$

The N/D method: nonrelativistic

$$n(k^2) = n_m(k^2) + \frac{(k^2)^m}{\pi} \int_{-\infty}^{k_{\text{lhc}}^2} \frac{d(k'^2) \text{Im} f(k'^2)}{(k'^2 - k^2)(k'^2)^m} dk'^2$$

$\propto \text{Im} L$

No singularity along lhc

$$n(k^2) = n'_m(k^2) + \frac{P(k^2)}{\pi} \int_{-\infty}^{k_{\text{lhc}}^2} \frac{\text{Im} f(k'^2)}{k'^2 - k^2} dk'^2 = n'_m(k^2) + P(k^2) \tilde{g} L(k^2)$$

$$n(k^2) = \tilde{n}(k^2) + \tilde{g}(L(k^2) - L_0)$$

$$L_0 = L(k^2 = 0) = -1/\mu_{\text{ex}}^2$$

$$\begin{aligned} d(k^2) &= \tilde{d}(k^2) - ik(\tilde{n}(k^2) - \tilde{g}L_0) - \frac{\tilde{g}}{\pi} \int_0^\infty \frac{k' L(k'^2)}{k'^2 - k^2} dk'^2 \\ &= \tilde{d}(k^2) - ik n(k^2) - \tilde{g} d^{\text{R}}(k^2) \end{aligned}$$

$$d_u^{\text{R}}(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$

It is worth stressing that $d(k^2)$ is free of lhc, as the lhc associated with $n(k^2)$ below the threshold is counterbalanced by $d^{\text{R}}(k^2)$, which is crucial to ensure that $f(k^2)$ exhibits the correct lhc behavior. Along the rhc, both $n(k^2)$ and $d^{\text{R}}(k^2)$ are real such that $\text{Im} d(k^2) = -k n(k^2)$.

Effective range expansion with the left-hand cut

$$\frac{1}{f(k^2)} = \frac{\tilde{d}(k^2) - \tilde{g}d^R(k^2)}{\tilde{n}(k^2) + \tilde{g}(L(k^2) - L_0)} - ik$$

$$d_u^R(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$

$$L(k^2) = -\frac{1}{4k^2} \log \frac{\mu_+^2/4 + k^2}{\mu_+^2/4 + \eta^2 k^2}$$

$$\frac{1}{f_{[m,n]}(k^2)} = \frac{\sum_{i=0}^n \tilde{d}_i k^{2i} - \tilde{g}d^R(k^2)}{1 + \sum_{j=1}^m \tilde{n}_j k^{2j} + \tilde{g}(L(k^2) - L_0)} - ik$$

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g}d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

$$\tilde{g} \rightarrow 0$$

$$\frac{1}{f(k^2)} = \frac{1}{a} + \frac{1}{2}rk^2 - ik$$

Scattering length

$$a = f(k^2 = 0) = \left[\tilde{d}_0 + \frac{\tilde{g}}{\mu_+} (1 - \eta) \right]^{-1}$$


Effective range

$$r = \left. \frac{d^2(1/f + ik)}{dk^2} \right|_{k=0} = 2\tilde{d}_1 - \frac{8\tilde{g}}{3\mu_+^3} (1 - \eta^3) - \frac{4\tilde{g}}{\mu_+^4 a_u} (1 - \eta^4)$$

Example: Tcc on the Lattice [3 parameters]


$f_{[0,1]}$

Du et al., 2408.09375 [hep-ph]




$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g} d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

Couplings to the exchanged-particle



$$g_P = -\frac{2\pi\tilde{g}}{\mu d^{0,\text{lhc}} \mathcal{F}_\ell}$$

$$g_{D^* D \pi}^2 / (4F^2) = 9.2 \text{ GeV}^{-2}$$



$$-\frac{2\pi}{\mu} \text{Im} f = \text{Im} T = \text{Im} V_{\text{OPE}}(k^2) = g_P \frac{-\pi}{4k^2} \mathcal{F}_\ell, \quad \text{for } k^2 < k_{\text{lhc}}^2$$

$$\text{Im} n(k^2) = -\tilde{g} \frac{\pi}{4k^2}, \quad \text{for } k^2 < k_{\text{lhc}}^2$$

$$d_u^{0,\text{lhc}} = \tilde{d}_0 - \frac{\tilde{d}_1 \mu_+^2}{4} + \frac{\mu_+}{2} \left(1 + \frac{\tilde{g}}{\mu_{\text{ex}}^2} \right) + \frac{\tilde{g} \log[2/(1+\eta)]}{\mu_+}$$

The amplitude zero

At leading order, i.e., $\tilde{n}(k^2) = 1$,

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g} d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

For a general u -channel exchange,

$$1 + \tilde{g} \left[L_u(k_{u,\text{zero}}^2) + \frac{1}{\mu_{\text{ex}}^2} \right] = 0,$$

for the case $|\Delta| \ll m_{\text{th}}$ such that $\eta \ll 1$, \Rightarrow the t -channel exchange

$$k_{t,\text{zero}}^2 = -\frac{m_{\text{ex}}^2}{4} \left[1 + \frac{1}{y} W(-e^{-y} y) \right]$$

where $y \equiv 1 + m_{\text{ex}}^2/\tilde{g}$ and W is the Lambert W function.

$$y = 1 + \frac{1 + \frac{4}{3} a_t m_{\text{ex}} (1 - \log 4) - \frac{4\pi a_t m_{\text{ex}}^2}{\mu g_P \mathcal{F}_\ell}}{2 + a_t m_{\text{ex}} (1 - m_{\text{ex}} r_t/4)}.$$

Summary

- The three-body cut: one-pion exchange + self-energy of D^*
- ★ Unphysical pion masses on the Lattice
 $M_D = 1927 \text{ MeV}$, $M_{D^*} = 2049 \text{ MeV}$, $m_\pi = 280 \text{ MeV}$
 - ↪ the three-body cut above the two-body cut ($\sqrt{s_{\text{thc}}} = 3968 \text{ MeV}$)
 - ↪ The traditional ERE valid only in a very limited range
 - ↪ An accurate extraction of the pole requires the OPE implemented
- ★ The ERE with the left-hand cut

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g} d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

- ↪ correct behavior of the left-hand cut
- ↪ can be used to extract the couplings of the exchanged particle to the scattering particles
- ↪ amplitude zeros caused by the interplay between the short- and long-range interactions

Thank you very much for your attention!

Thank you very much for your attention!

Without $d^R(k^2)$

$$d(k^2) = \tilde{d}(k^2) - ik(\tilde{n}(k^2) - \tilde{g}L_0) - \frac{\tilde{g}}{\pi} \int_0^\infty \frac{k' L(k'^2)}{k'^2 - k^2} dk'^2$$

$$= \tilde{d}(k^2) - ik n(k^2) - \cancel{\tilde{g} d^R(k^2)}$$

$$d_u^R(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$

lhc

$$f^{-1} = \frac{\frac{1}{a} + \frac{1}{2}rk^2}{1 + \tilde{g}(L(k^2) - L_0)} - ik$$

$$d(k^2) = \frac{1}{a} + \frac{1}{2}rk^2 - ik - ik\tilde{g}(L(k^2) - L_0)$$

