






Recent results of Belle II

殷俊昊
南开大学


SuperKEKB

$$e^- \xrightarrow{7 \text{ GeV}} (\star) \xleftarrow{4 \text{ GeV}} e^+$$

Belle II



$$\begin{aligned} \sigma_y^* &= 940 \text{ nm} \\ \beta_y^* &= 5.9 \text{ mm} \\ \sigma_x^* &= 147/170 \text{ } \mu\text{m} \end{aligned}$$



$$\begin{aligned} \sigma_y^* &= 48/62 \text{ nm} \\ \beta_y^* &= 0.27/0.3 \text{ mm} \\ \sigma_x^* &= 10.1/10.7 \text{ } \mu\text{m} \end{aligned}$$

$$\mathcal{L}_{\text{II}}^{\text{peak}} \approx 30 \times \mathcal{L}_{\text{I}}^{\text{peak}}$$


$$\int^{\text{goal}} \mathcal{L}_{\text{II}} dt = 50 \text{ ab}^{-1} \approx 50 \int \mathcal{L}_{\text{I}} dt$$

Belle II RUN-I (2019-2022)

luminosity: $4.7 \times 10^{34} / \text{cm}^2 / \text{s}$! $> 2 \text{ fb}^{-1}$ per day !


June, 2022

$L_{\text{peak}} = 4.653 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ @ 22:58:08 06/08		06/07 23:59:36 - 06/08 23:59:36, 2022 JST			
HER I_{peak}	1127 mA	n_b	2249	β_x^* / β_y^*	60 / 1 mm
int. L/day	1253 / 1681 pb^{-1}	LER I_{peak}	1405 mA	n_b	2249 β_x^* / β_y^* 80 / 1 mm



record of KEKB/Belle
 $2 \times 10^{34} / \text{cm}^2 / \text{s}$; currents $> 1 \text{ A}$
 record of PEP-II/BaBar
 $1 \times 10^{34} / \text{cm}^2 / \text{s}$; currents $> 2 \text{ A}$

**$5 \times 10^{34} / \text{cm}^2 / \text{s}$
 $\sim 1 \text{ BaBar}/\text{year}$**




**squeezing further β_y^* ($\rightarrow 0.6 \text{ mm}$)
 doubling (or more) the currents
 $\Rightarrow L > 10^{35} / \text{cm}^2 / \text{s}$ after LS 1**

- $(364 \pm 2) \text{ fb}^{-1} \Upsilon(4S)$ on resonance data
- $(42.6 \pm 0.3) \text{ fb}^{-1} \Upsilon(4S)$ off resonance data
- $(19.7 \pm 0.1) \text{ fb}^{-1} \Upsilon(10753)$ scan data



Today's highlight

$\Upsilon(10753)$ study



CKM matrix element measurement

B rare decays


τ physics

$|V_{ub}|$ using $B^0 \rightarrow \pi^- l^+ \nu_l$, $B^+ \rightarrow \rho^0 l^+ \nu_l$

Simultaneous fit in 2D grid of M_{bc} and ΔE for each bin of q^2 :

13 bins for $B^0 \rightarrow \pi^- l^+ \nu_l$ mode, 10 bins for $B^+ \rightarrow \rho^0 l^+ \nu_l$ mode.

Preliminary
NEW!!



$$M_{bc} = \sqrt{E_{\text{beam}}^{*2} - |\vec{p}_B^*|^2} = \sqrt{\left(\frac{\sqrt{s}}{2}\right)^2 - |\vec{p}_B^*|^2}$$




$$\Delta E = E_B^* - E_{\text{beam}}^* = E_B^* - \frac{\sqrt{s}}{2}$$

$$\mathcal{B}(B^0 \rightarrow \pi^- l^+ \nu_l) = (1.516 \pm 0.042 \pm 0.059) \times 10^{-4}$$

$$\mathcal{B}(B^+ \rightarrow \rho^0 l^+ \nu_l) = (1.625 \pm 0.079 \pm 0.180) \times 10^{-4}$$

stat syst

Consistent with world average.



Extracted $|V_{ub}|$ with lattice QCD and/or light-cone sum rules (LCSR) constraints of form factors

$$|V_{ub}|_{B \rightarrow \pi \ell \nu} = (3.93 \pm 0.09(\text{stat}) \pm 0.13(\text{syst}) \pm 0.19(\text{theo})) \times 10^{-3}, (\text{LQCD})$$

$$= (3.73 \pm 0.07(\text{stat}) \pm 0.07(\text{syst}) \pm 0.16(\text{theo})) \times 10^{-3}, (\text{LQCD+LCSR})$$

$$|V_{ub}|_{B \rightarrow \rho \ell \nu} = (3.19 \pm 0.12(\text{stat}) \pm 0.18(\text{syst}) \pm 0.26(\text{theo})) \times 10^{-3}, (\text{LCSR})$$

Preliminary

NEW!!




$\sin 2\phi_1$ measurement

- Sensitive to BSM physics
- Fit Δt to extract S_{CP} and C_{CP} :


$$f_{CP}^{\text{true}} = \frac{1}{4\tau_B^0} e^{-|\Delta t|/\tau_B^0} (1 + q[S_{CP}\sin(\Delta m\Delta t) - C_{CP}\cos(\Delta m\Delta t)])$$

- SM expectation: $S_{CP} = \sin 2\phi_1$, and $C_{CP} = 0$
- Deviation from $\sin 2\phi_1$ would suggest BSM physics
- Sensitive to BSM physics in

- $b \rightarrow sq\bar{q}$
- $b \rightarrow s\gamma$



channel	S_meas	C_meas	
$B^0 \rightarrow K_S^0 J/\psi$	$0.724 \pm 0.035 \pm 0.014$	$-0.035 \pm 0.026 \pm 0.012$	preliminary
$B^0 \rightarrow K_S^0 \pi^0 \gamma$	$0.04^{+0.45}_{-0.44} \pm 0.10$	$-0.06 \pm 0.25 \pm 0.07$	preliminary
$B^0 \rightarrow \eta' K_S^0$	$0.67 \pm 0.10 \pm 0.04$	$-0.19 \pm 0.08 \pm 0.03$	preliminary
$B^0 \rightarrow \pi^0 K_S^0$	$0.75^{+0.20}_{-0.23} \pm 0.04$	$-0.04^{+0.14}_{-0.15} \pm 0.05$	PRL 131, 111803 (2023)
$B^0 \rightarrow \phi K_S^0$	$0.54 \pm 0.26^{+0.06}_{-0.08}$	$-0.31 \pm 0.20 \pm 0.05$	PRD 108, 072012 (2023)



- Consistent with world average and SM expectation.
- $B^0 \rightarrow \eta' K_S^0$ provides the most sensitive results up to date.
- Smaller data size but equivalent uncertainties, sometimes better.



Combined measurement of ϕ_3 with Belle & Belle II

Preliminary

- Four different methods using 17 different final states
- Inputs on D decays dynamics from other experiments
 - r_D (amplitude ratio), δ_D (strong-phase difference), κ_D (coherence factor), etc.

B decay	D decay	Method	Data set (Belle + Belle II) [fb ⁻¹]
$B^+ \rightarrow Dh^+$	$D \rightarrow K_S^0 h^- h^+$	BPGGSZ	711 + 128 [JHEP 02 063 (2022)]
$B^+ \rightarrow Dh^+$	$D \rightarrow K_S^0 \pi^- \pi^+ \pi^0$	BPGGSZ	711 + 0 [JHEP 10 178 (2019)]
$B^+ \rightarrow Dh^+$	$D \rightarrow K_S^0 \pi^0, K^- K^+$	GLW	711 + 189 [arxiv:2308.05048]
$B^+ \rightarrow Dh^+$	$D \rightarrow K^+ \pi^-, K^+ \pi^- \pi^0$	ADS	711 + 0 [PRL 106 231803 (2011)]
$B^+ \rightarrow Dh^+$	$D \rightarrow K_S^0 K^- \pi^+$	GLS	711 + 362 [JHEP 09 (2023) 146]
$B^+ \rightarrow D^* K^+$	$D \rightarrow K_S^0 \pi^- \pi^+$	BPGGSZ	605 + 0 [PRD 81 112002 (2010)]
$B^+ \rightarrow D^* K^+$	$D \rightarrow K_S^0 \pi^0, K_S^0 \phi, K_S^0 \omega, K^- K^+, \pi^- \pi^+$	GLW	210+0 [PRD 73 051106 (2006)]


Parameters	$\phi_3(^{\circ})$	r_B^{DK}	$\delta_B^{DK}(^{\circ})$	$r_B^{D\pi}$	$\delta_B^{D\pi}(^{\circ})$	$r_B^{D^*K}$	$\delta_B^{D^*K}(^{\circ})$
PLUGIN method							
Best fit value	78.6	0.117	138.4	0.0165	347.0	0.234	341
68.3% interval	[71.4, 85.4]	[0.105, 0.130]	[129.1, 146.5]	[0.0109, 0.0220]	[337.4, 355.7]	[0.165, 0.303]	[327, 355]
95.5% interval	[63, 92]	[0.092, 0.141]	[118, 154]	[0.006, 0.027]	[322, 366]	[0.10, 0.37]	[307, 369]




$\phi_3 = (78.6 \pm 7.3)^{\circ}$, consistent with WA, $\phi_3 = (66.2_{-3.6}^{+3.2})^{\circ}$, within 2σ

Measurement of branching fraction of $B^+ \rightarrow K^+ \nu \bar{\nu}$

[arXiv:2311.14647](https://arxiv.org/abs/2311.14647)






- The $B \rightarrow K \nu \nu$ process is known with high accuracy in the SM:
 - $\mathcal{B}(B \rightarrow K^+ \nu \nu) = (5.6 \pm 0.4) \times 10^{-6}$ (arXiv:2207.13371)
 - *We use 4.97×10^{-6} as a reference, after removal of $B \rightarrow \tau(K\bar{\nu})\nu$*
- Extensions beyond SM may lead to significant rate increase
- Very challenging experimentally, not yet observed
 - Low branching fraction, high background contributions
 - 3-body kinematics, no good kinematic variable to fit



Analysis strategy

[arXiv:2311.14647](https://arxiv.org/abs/2311.14647)



- Two analyses:
 - More sensitive inclusive tagging (ITA)
 - Conventional hadronic tagging (HTA)
- Kinematic properties to suppress background with MVA
- Use classifier output as (one of) the fit variable(s), use simulation for signal and background templates
- Use multiple control channels to validate simulation with data

FIT



arXiv:2311.14647

- Extract signal from maximum likelihood fit
- Inclusive tag: in bins of q_{rec}^2 and $\eta(\text{BDT}_2)$
- Hadronic tag: in bins of $\eta(\text{BDT}_h)$
- Signal is extracted in terms of signal strength μ

signal relative to SM expectation

- Inclusive tag: $\mu = 5.4 \pm 1.0(\text{stat}) \pm 1.1(\text{syst})$
- Hadronic tag: $\mu = 2.2_{-1.7}^{+1.8}(\text{stat})_{-1.1}^{+1.6}(\text{syst})$
- Combined: $\mu = 4.6 \pm 1.0(\text{stat}) \pm 0.9(\text{syst})$

ITA and HTA results are consistent at 1.2σ level



Inclusive tag: $\mathcal{B} = 2.7 \pm 0.5(\text{stat}) \pm 0.5(\text{syst})$

Hadronic tag: $\mathcal{B} = 1.1^{+0.9}_{-0.8}(\text{stat})^{+0.8}_{-0.5}(\text{syst})$

Combined: $\mathcal{B} = 2.3 \pm 0.5(\text{stat})^{+0.5}_{-0.4}(\text{syst})$

For the inclusive tag, significance of the result

- wrt null hypothesis is 3.5σ

- wrt SM is 2.9σ

For the hadronic tag, significance of the result

- wrt null hypothesis is 1.1σ



- wrt SM is 0.6σ

For the combination, significance of the result

- wrt null hypothesis is 3.5σ


- wrt SM is 2.7σ

First evidence of the $B^+ \rightarrow K^+ \nu \bar{\nu}$ decay!



Search for $B \rightarrow \gamma\gamma$

Belle + Belle II, Preliminary




Very rare in SM with $\mathcal{B}(B \rightarrow \gamma\gamma) = (1.4_{-1.8}^{+1.4}) \times 10^{-8}$ (JHEP12, 169(2020))

Previous searches	Limits
L3 (73 pb ⁻¹)	$< 3.9 \times 10^{-5}$
Belle (104 fb ⁻¹)	$< 6.2 \times 10^{-7}$
BaBar (426 fb ⁻¹)	$< 3.2 \times 10^{-7}$

Big challenge due to large background.

- Simultaneous 3D fit to M_{bc} , ΔE , and C'_{BDT} .
- Combined signal yield: $11.0_{-5.48}^{+6.48}$ with significance of 2.5σ



	$\mathcal{B}(B^0 \rightarrow \gamma\gamma)$	$\mathcal{B}(B^0 \rightarrow \gamma\gamma)$ (at 90% CL)
Belle	$(5.4_{-2.6}^{+3.3} \pm 0.5) \times 10^{-8}$	$< 9.9 \times 10^{-8}$
Belle II	$(1.7_{-2.4}^{+3.7} \pm 0.3) \times 10^{-8}$	$< 7.4 \times 10^{-8}$
Combined	$(3.7_{-1.8}^{+2.2} \pm 0.7) \times 10^{-8}$	$< 6.4 \times 10^{-8}$

x 5 improvement over previous best UL.

CP and isospin asymmetries in $B \rightarrow \rho\gamma$

Combine Belle+Belle II **preliminary**


- Clean environment to search for BSM physics
- SM expect CP-average isospin asymmetry as $\bar{A}_I^{\text{SM}} = (5.2 \pm 2.8) \%$

$$A_I = (A_I^{\bar{0}^-} + A_I^{0^+})/2, \text{ with } A_I^{\bar{0}^-} = \frac{c_\rho^2 \Gamma(\bar{B}^0 \rightarrow \rho^0 \gamma) - \Gamma(B^- \rightarrow \rho^- \gamma)}{c_\rho^2 \Gamma(\bar{B}^0 \rightarrow \rho^0 \gamma) + \Gamma(B^- \rightarrow \rho^- \gamma)}$$

- Current world average $A_I^{\text{exp}} = (30_{-13}^{+16}) \%$.


$$\begin{aligned} \mathcal{B}(B^+ \rightarrow \rho^+ \gamma) &= (12.85_{-1.92-1.13}^{+2.02+1.38}) \times 10^{-7}, \\ \mathcal{B}(B^0 \rightarrow \rho^0 \gamma) &= (7.45_{-1.27-0.79}^{+1.33+0.97}) \times 10^{-7}, \\ A_{\text{CP}}(B^+ \rightarrow \rho^+ \gamma) &= (-7.1_{-15.2-1.3}^{+15.3+1.4}) \%, \\ A_I(B \rightarrow \rho\gamma) &= (14.2_{-11.7-6.4-6.5}^{+11.0+6.6+6.0}) \%, \end{aligned}$$

- *Consistent with the SM prediction.*
- *Highest precision to date; supersede the previous measurements from Belle*



τ physics

B factory is also τ factory



@10.58 GeV:

$$\sigma(e^+e^- \rightarrow \tau^+\tau^-) = 0.92 \text{ nb}$$

$$\sigma(e^+e^- \rightarrow \Upsilon(4S)) = 1.11 \text{ nb}$$

- High luminosity.
- Well-defined initial state.
- High vertex resolution.
- Excellent calorimetry.
- Sophisticated particle ID.
- Ability to trigger low-multiplicity event


τ Lepton Flavor Universality Violation

preliminary


$$\left(\frac{g_\mu}{g_e}\right)_\tau = \sqrt{\frac{BF[\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau] f(m_e^2/m_\tau^2)}{BF[\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau] f(m_\mu^2/m_\tau^2)}}$$

In the SM: $\left(\frac{g_\mu}{g_e}\right)_\tau = 1$

Tests with 3x1 topology: same method as Babar



$$R_\mu = \frac{\mathcal{B}(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = 0.9675 \pm 0.0007 \pm 0.0036.$$





Consistent with previous measurements

Most precise to date

τ Lepton Flavor Violation

preliminary

- A search for the charged-lepton-flavour violating decay $\tau \rightarrow \mu\mu\mu$
- Provide indisputable evidence of physics beyond the SM.






- Novel inclusive tagging followed by a BDT-based selection.
 - 2.5 times higher efficiency than Belle and 37% higher efficiency than 1-prong tagging
- One event in signal region:
 - $B(\tau^- \rightarrow \mu^- \mu^+ \mu^-) = (3.1 + 8.7 \pm 0.1) \times 10^{-9}$
- $< 1.9(1.8) \times 10^{-8}$ for observed (expected) limit at 90% C.L.
- Less data, more restrictive than Belle

Search for $\Upsilon(10753) \rightarrow \omega\eta_b, \omega\chi_{b0}$

arXiv:2312.13043

- $\Upsilon(10753) \rightarrow \omega\eta_b(1S)$ is strongly enhanced in tetraquark model.
- Analogy to $Y(4230) \rightarrow \omega\chi_{c0}$ in searching for $\omega\chi_{b0}$



	$\eta_b(1S)\omega$	$\chi_{b0}(1P)\omega$
Born cross section (pb)	$0.5 \pm 1.1 \pm 0.5$	$2.8 \pm 3.1 \pm 2.1$
90% C.L. UL	< 2.4	< 9.1

UL of $\omega\eta_b(1S)$ is similar to $\pi\pi\Upsilon(nS)$. Strongly inconsistent with one of the tetraquark model.







UL with full reconstruction is 11.3 fb - similar sensitivities.

Measurement of $\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(nS)$, $n=1,2,3$

arXiv:2401.12021

- Full reconstruction of $\pi^+\pi^-\mu^+\mu^-$, clear signal of $\Upsilon(10753)$ as well as other $\Upsilon(2,3S)$ signals.
- Evident signal of $\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(1S)$ and $\pi^+\pi^-\Upsilon(2S)$; no evidence of $\pi^+\pi^-\Upsilon(3S)$.
- No $Z_b(10610/10650)$; $M(\pi^+\pi^-)$ could be parameterized like $\Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$

$p^*(\pi^+\pi^-\mu^+\mu^-) < 0.1 \text{ GeV}$



Fit with three coherent BW, convoluting a Gaussian modeling energy spread:

$$\sigma \propto \left| \sum_i^3 \frac{\sqrt{12\pi\Gamma_i\mathcal{B}_i}}{s - M_i + iM_i\Gamma_i} \cdot \sqrt{\frac{f(\sqrt{s})}{f(M_i)}} e^{i\phi_i} \right|^2 \otimes G(0, \delta E)$$

All parameters are free, except $\delta E = 0.0056$ GeV

Parameters of $\Upsilon(10753)$:


$$M = 10756.3 \pm 2.7_{(stat.)} \pm 0.6_{(syst.)} \text{ MeV}/c^2$$

$$\Gamma = 29.7 \pm 8.5_{(stat.)} \pm 1.1_{(syst.)} \text{ MeV}$$

Agree with previous Belle measurement.

Improve uncertainties ~2 times smaller

	resonance mass (MeV/ c^2)	width (MeV)
$\Upsilon(5S)$	10884.7 ± 1.2	38.7 ± 3.7
$\Upsilon(6S)$	10995.5 ± 4.2	34.6 ± 8.6



arXiv:2401.12021

Summary

- Belle II has collected 424/fb data before Long Shutdown 1.
- New data, new methods, new ideas lead to brand new results
 - First evidence of $B \rightarrow K\nu\nu$
 - First determination of ϕ_3 from Belle and Belle II
 - τ factory! Most precise τ properties measurement and search for NP
 - Properties study of $\Upsilon(10753)$, unique in Belle II
 - ...
- LS1 is finishing and new run has started.
 - More data, more new results

BACK UP

Reconstruction and background suppression

Charged particles:

- $p_t > 0.1 \text{ GeV}/c$
- close to collision point
- in central region

Neutral particles:

- $E > 100 \text{ MeV}$
- in central region

Signal candidate:

- **charged particle**
- Kaon identification
- Minimal q_{rec}^2

Event (pre-selection):

- $4 \leq N_{\text{track}} \leq 10$
- $E_{\text{total}} > 4 \text{ GeV}$
- $17^\circ < \vartheta_{\text{miss}} < 160^\circ$

BDT₁ (first filter):

- 12 event-shape based kinematic variables


BDT₂ (final selection):

- 35 input variables:
using signal, event, and their correlations

- Selection criteria for particles to ensure high and well-measured efficiency
- Signal candidate selected using mass of the neutrino pair q_{rec}^2 (computed as K^+ recoil)
- Three-step filter: basic event cuts, BDT-based filter (BDT₁) and final selection (BDT₂). BDT₂ improves performance in terms of $s/\sqrt{s+b}$ by almost factor 3


*Missing momentum is reconstructed using beam and all reconstructed particle 4-momenta

Examples of input variables for BDT₁ and BDT₂





- Example of input distributions at pre-selection level, 1% of data, with detector-level corrections applied but no physics modeling corrections
- Each variable is examined to have reasonable description by simulation and significant separation power

Signal efficiency validation



- Use cleanly reconstructed $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ decays with $\mu^+ \mu^-$ pair removed and K^+ kinematics adjusted to validate the **signal efficiency** in simulation. The ratio of data/simulation efficiency in the signal region is **1.00 ± 0.03**


Background from $B \rightarrow D(\rightarrow K^+ X) l \nu$ and $B \rightarrow K^+ D(\rightarrow K_L X)$



- Main backgrounds: semileptonic $B \rightarrow D(\rightarrow K^+ X) l \nu$ decays and prompt $B \rightarrow K^+ X$ production (>90%)
- Semileptonic decays suppressed by several MVA variables, checked at each selection step
- Prompt K^+ production studied using prompt π^+ from $B^+ \rightarrow \pi^+ X$ (and l^+ from $B^+ \rightarrow l^+ X$) decays
- Systematic uncertainties on decay branching fractions, enlarged for $D(\rightarrow K_L X)$ and $B \rightarrow D^{**} l \nu$

Background from $B^+ \rightarrow K^+ K^0 K^0$



Most signal-like backgrounds



← $B^+ \rightarrow K^+ K_S K_S$ decays




- Backgrounds from $B^+ \rightarrow K^+ nn$ and $B^+ \rightarrow K^+ K^0 K^0$ have branching fractions of few $\times 10^{-5}$, however K_L and neutrons can **escape** EM calorimeter
- $B^+ \rightarrow K^+ K^0 K^0$ modeled based on BaBar analysis ([arXiv:1201.5897](https://arxiv.org/abs/1201.5897))
- Dedicated checks of **K_L performance** in calorimeter using radiative φ production
- Dedicated checks using $B^+ \rightarrow K^+ K_c K_c$ and $B^0 \rightarrow K_c K^+ K^-$ control channels

High-precision vertexing and its importance for TDCPV





Improved vertex resolution
due to pixel detector
(despite lower boost)

γ measurement in $B^+ \rightarrow D(K_S^0 h^+ h^-) h^+$ with Belle and Belle II data




Determine bin-by-bin asymmetries
 $(N_-^{-i} - N_+^{+i}) / (N_-^{-i} + N_+^{+i})$
 in each Dalitz plot bin i



$$x_{\pm}^{\text{DK}} = r_B^{\text{DK}} \cos(\delta_B^{\text{DK}} \pm \phi_3)$$

$$y_{\pm}^{\text{DK}} = r_B^{\text{DK}} \sin(\delta_B^{\text{DK}} \pm \phi_3)$$



$$\phi_3 = (78.4 \pm 11.4 \pm 0.5 \pm 1.0)^\circ,$$

$$r_B^{\text{DK}} = 0.129 \pm 0.024 \pm 0.001 \pm 0.002,$$

$$\delta_B^{\text{DK}} = (124.8 \pm 12.9 \pm 0.5 \pm 1.7)^\circ,$$

$$r_B^{\text{D}\pi} = 0.017 \pm 0.006 \pm 0.001 \pm 0.001,$$

$$\delta_B^{\text{D}\pi} = (341.0 \pm 17.0 \pm 1.2 \pm 2.6)^\circ.$$

Measurement of $B^\pm \rightarrow D_{CP\pm} K^\pm$ with Belle and Belle II data

- Simultaneous fit to $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ with D decays to CP eigenstates

$$\mathcal{R}_{CP\pm} \equiv \frac{\mathcal{B}(B^- \rightarrow D_{CP\pm} K^-) + \mathcal{B}(B^+ \rightarrow D_{CP\pm} K^+)}{(\mathcal{B}(B^- \rightarrow D_{\text{flav}} K^-) + \mathcal{B}(B^+ \rightarrow \bar{D}_{\text{flav}} K^+))/2}$$





$$\mathcal{A}_{CP\pm} \equiv \frac{\mathcal{B}(B^- \rightarrow D_{CP\pm} K^-) - \mathcal{B}(B^+ \rightarrow D_{CP\pm} K^+)}{\mathcal{B}(B^- \rightarrow D_{CP\pm} K^-) + \mathcal{B}(B^+ \rightarrow D_{CP\pm} K^+)}$$

$$\mathcal{R}_{CP\pm} \approx \frac{R_{CP\pm}}{R_{\text{flav}}}$$

$$\mathcal{R}_{CP\pm} = 1 + r_B^2 \pm 2r_B \cos \delta_B \cos \phi_3$$

$$\mathcal{A}_{CP\pm} = \pm 2r_B \sin \delta_B \sin \phi_3 / \mathcal{R}_{CP\pm}$$

	68.3% CL	95.4% CL
ϕ_3 ($^\circ$)	[8.5, 16.5]	[5.0, 22.0]
	[84.5, 95.5]	[80.0, 100.0]
	[163.3, 171.5]	[157.5, 175.0]
r_B	[0.321, 0.465]	[0.241, 0.522]



R(X) Result

-The first results of $R(X) = \frac{\mathcal{B}(B \rightarrow X\tau\nu_\tau)}{\mathcal{B}(B \rightarrow X\ell\nu_\ell)}$ at B factory:



$$R(X_{\tau/e}) = 0.232 \pm 0.042, [0.020 \text{ (stat)}, 0.037 \text{ (syst)}]$$

$$R(X_{\tau/\mu}) = 0.222 \pm 0.057, [0.027 \text{ (stat)}, 0.050 \text{ (syst)}]$$

$$R(X_{\tau/\ell}) = 0.228 \pm 0.039, [0.016 \text{ (stat)}, 0.036 \text{ (syst)}]$$

Major systematics: MC statistics, PDF shape, BR of $B \rightarrow D^{**}\ell\nu$

-Consistent with SM prediction



$\sin 2\phi_1$ using $B^0 \rightarrow K_S^0 J/\psi$ preliminary

- Use $J/\psi \rightarrow e^+ e^-, \mu^+ \mu^-, K_S^0 \rightarrow \pi^+ \pi^-$

- Analysis method:

- Employ Graph Flavor tagger based on Dynamic Graph Convolution Neural Network (GFlaT)


- Determine signal yields and subtract background using sWeights from ΔE fit






- Fit Δt to extract S_{CP} and C_{CP} :

$$f_{CP}^{\text{true}} = \frac{1}{4\tau_B^0} e^{-|\Delta t|/\tau_B^0} (1 + q[S_{CP}\sin(\Delta m\Delta t) - C_{CP}\cos(\Delta m\Delta t)])$$

- SM expectation: $S_{CP} = \sin 2\phi_1$, and $C_{CP} = 0$

- Flavor resolution effect and resolution function taken from calibration with $B^0 \rightarrow D^{(*)+} \pi^-$





cross check:

SM expectation: $S_{cp} = C_{CP} = 0$

$$S_{cp} = 0.008 \pm 0.019$$

$$C_{cp} = -0.018 \pm 0.026$$

$$C_{CP} = -0.035 \pm 0.026 \text{ (stat)} \pm 0.012 \text{ (syst)}$$

$$S_{CP} = 0.724 \pm 0.035 \text{ (stat)} \pm 0.014 \text{ (syst)}$$

Previous stat. uncertainties:
 Belle II ICHEP22: $\sigma S_{CP} = 0.062$
 (improvement equivalent to 3.1X larger dataset)

Previous results (J/ψ KS only):
 Belle 2012: $\sigma S_{CP} = 0.029$
 BaBar 2009: $\sigma S_{CP} = 0.036$
 LHCb 2023: $\sigma S_{CP} = 0.015$



GFlaT established as standard tool for forthcoming TDCPV analyses

CP asymmetries in $B^0 \rightarrow K_S^0 \pi^0 \gamma$

- $b \rightarrow s\gamma$ proceeds via one-loop diagrams
 - Sensitive to BSM physics
- Mixing-induced time-dependent CP asymmetries (S) are expected to be small

- $S_{CP} = -0.035 \pm 0.017$ (arXiv:hep-ph/0406055)

preliminary



Most precise result up to date!

$$S(K^{*0}\gamma) = 0.00^{+0.27+0.03}_{-0.26-0.04},$$

$$C(K^{*0}\gamma) = 0.10 \pm 0.13 \pm 0.03,$$

$$S(K_S^0\pi^0\gamma) = 0.04^{+0.45}_{-0.44} \pm 0.10,$$

$$C(K_S^0\pi^0\gamma) = -0.06 \pm 0.25 \pm 0.07,$$

HFLAV:

$$K^{*0}\gamma: C_{CP} = -0.04 \pm 0.14 \quad S_{CP} = -0.16 \pm 0.22$$



$$K_S\pi^0\gamma: C_{CP} = -0.07 \pm 0.12 \quad S_{CP} = -0.15 \pm 0.20$$

22.08.2023 *The HFLAV $K_S\pi^0\gamma$ values include $K^{*0}\gamma$

CP asymmetries in $B^0 \rightarrow \eta' K_S^0$

preliminary

- Process $b \rightarrow sq\bar{q}$ via loop amplitude
- High transition rate relative to other gluonic penguins
- Additional source BSM could be involved
- Deviation from $\sin 2\phi_1$ would suggest BSM physics
- Cross checked with $B^+ \rightarrow \eta' K^+$, where no CP asymmetry is expected




$$C_{\eta' K_S^0} = -0.19 \pm 0.08 \pm 0.03,$$

$$S_{\eta' K_S^0} = 0.67 \pm 0.10 \pm 0.04,$$

$C = -0.05 \pm 0.04$ and $S = 0.63 \pm 0.06$ from HFLAV

Measurement of CPV in $B^0 \rightarrow \pi^0 K_S^0$

- Process $b \rightarrow s d \bar{d}$ via loop amplitude
- High transition rate relative to other gluonic penguins
- Additional source BSM could be involved
- Deviation from $\sin 2\phi_1$ would suggest BSM physics
- Cross checked with $B^+ \rightarrow \eta' K^+$, where no CP asymmetry is expected



Time dependent CPV using $B^0 \rightarrow K_S^0 K_S^0 K_S^0$

preliminary




- Current world average:


$$C = -0.15 \pm 0.12$$

$$S = -0.83 \pm 0.17$$

Use the likelihood from the fit,
resulting 2D confidence intervals

Need improvement on uncertainties.





PRD 108, 072012 (2023)

- Current world average:

$$C = 0.01 \pm 0.14$$



$$S = 0.74^{+0.11}_{-0.13}$$

whereas C is expected to be zero in SM, with $S = 0.02 \pm 0.01$

- Cross checked with $B^+ \rightarrow \phi K^+$, resulting $C = -0.12 \pm 0.10$ and $S = -0.09 \pm 0.12$

$$C = -0.31 \pm 0.20 \pm 0.05$$

$$S = 0.54 \pm 0.26^{+0.06}_{-0.08}$$





- *Compatible with previous determinations from Belle and BABAR.*
- *Similar uncertainty on C despite of smaller data sample.*
- *Improvement on the statistical uncertainty on S for the same number of signal events.*

First measurement of $R(X_{\tau/\ell})$

arXiv:2311.07248

- $R(X_{\tau/\ell}) = \frac{\mathcal{B}(B \rightarrow X\tau\nu_\tau)}{\mathcal{B}(B \rightarrow X\ell\nu_\ell)}$, measure inclusively. First measurement at B factories
- X reconstructed from remaining tracks and neutral clusters.
- Variables for yield extraction, 2D-fit to M_{miss}^2 and p_ℓ^B
- Results:
 - $R(X_{\tau/e}) = 0.232 \pm 0.020(\text{stat.}) \pm 0.037(\text{syst.})$
 - $R(X_{\tau/\mu}) = 0.222 \pm 0.027(\text{stat.}) \pm 0.050(\text{syst.})$
- Combining:
 - $R(X_{\tau/\mu}) = 0.228 \pm 0.016(\text{stat.}) \pm 0.036(\text{syst.})$



In agreement with SM prediction and $R(D^{(*)})$ measurements



Angular asymmetries using $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$

PRL 131, 181801 (2023)

- $B^0 \rightarrow D^{*-} \ell \nu$ is mediated in the SM via W -boson exchange.
- Characterized in terms of a recoil parameter and 3 helicity angles.
- Calculate the asymmetry: $\mathcal{A}_x(w) = \frac{N_x^+ w - N_x^-(w)}{N_x^+ w + N_x^-(w)}$
- Difference $\Delta A_x(w) = A_x^\mu(w) - A_x^e(w)$, is sensitive to LUV.
- Separate signal candidates into angular categories + and - based on the measured value of x .



First comprehensive tests of LU in the angular distributions of semileptonic B decays.

Agrees well with SM.


Direct CPV using $B \rightarrow K\pi$ and $B \rightarrow \pi\pi$

arXiv:2310.06381

- Charmless hadronic B meson decays feature non-negligible contributions from loop amplitudes.
- Sensitive to contributions from non-SM physics.

$$I_{K\pi} = A_{K^+\pi^-} + A_{K^0\pi^+} \frac{Br(K^0\pi^+)}{Br(K^+\pi^-)} \frac{\tau_{B^0}}{\tau_{B^+}} - 2A_{K^+\pi^0} \frac{Br(K^+\pi^0)}{Br(K^+\pi^-)} \frac{\tau_{B^0}}{\tau_{B^+}} - 2A_{K^0\pi^0} \frac{Br(K^0\pi^0)}{Br(K^+\pi^-)} \approx 0$$

- Belle II measures all modes in coherent way with unique access to $B^0 \rightarrow K_S^0\pi^0$





$I_{K\pi} = -0.03 \pm 0.13 \pm 0.05$ (world average: $I_{K\pi} = 0.13 \pm 0.11$)
 competitive with world average with 362 fb^{-1}

Observation of $Y(10753) \rightarrow \omega\chi_{bJ}$ in $e^+e^- \rightarrow \gamma\omega Y(1S)$

PRL 130, 091902 (2023)

Clear $\omega\chi_{bJ}$ signals at $\sqrt{s} = 10.745$ and 10.805 GeV



$\Gamma_{ee}\mathcal{B}_f$	Solution I	Solution II
$\Gamma_{ee}\mathcal{B}(Y(10753) \rightarrow \omega\chi_{b1})$	$(0.63 \pm 0.39 \pm 0.20)$ eV	$(2.01 \pm 0.38 \pm 0.76)$ eV
$\Gamma_{ee}\mathcal{B}(Y(10753) \rightarrow \omega\chi_{b2})$	$(0.53 \pm 0.46 \pm 0.15)$ eV	$(1.32 \pm 0.44 \pm 0.55)$ eV

- $\frac{\Gamma_{ee}\mathcal{B}(Y(10753) \rightarrow \omega\chi_{b1})}{\Gamma_{ee}\mathcal{B}(Y(10753) \rightarrow \omega\chi_{b2})} \sim 1.0$ agrees with the expectation for HQET^[3]
- $\frac{\Gamma_{ee}\mathcal{B}(\omega\chi_{b1/2})}{\Gamma_{ee}\mathcal{B}(\pi^+\pi^-Y(2S))}$ ^[2] ~ 1.5 for $Y(10753)$ and ~ 0.1 for $Y(10870)$


[1]PRL 113, 142001(2014); [2]. JHEP 10, 220(2019); [3]. arXiv:hep-ph/9908366;

○ Confirm the existence of $Y(10753)$.

○ Large difference of $\frac{\mathcal{B}(\omega\chi_{bJ})}{\mathcal{B}(\pi^+\pi^-Y(nS))}$ between $Y(10753)$ and $Y(10870)$.


$|V_{cb}|$ using $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$

PRD 108, 092013 (2023)







- $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$ is parameterized by the recoil parameter (w) and three decay angles ($\theta_l, \chi, \theta_\nu$)
- 2D-binned likelihood fit to $(\cos\theta_{BY}, \Delta M)$ for each bin of variables.

$$\cos\theta_{BY} = \frac{2E_B^{CM} E_Y^{CM} - m_B^2 - m_Y^2}{2|p_B^{CM}||p_Y^{CM}|}, \quad \Delta M = M(D^{*+}) - M(D)$$



- Include all measured w , θ_l , χ , θ_V to extract form factor & $|V_{cb}|$
- Fit differential shapes with form factor expansion based on Caprini-Lellouch-Neubert (CLN) [Nucl. Phys. B530, 153 (1998)] & Boyd-Grinstein-Lebed (BGL) parameterisations [Phys. Rev. D56, 6895 (1997)]



$$|V_{cb}| \eta_{\text{EW}} \mathcal{F}(1) = \frac{1}{\sqrt{m_B m_{D^*}}} \left(\frac{|\tilde{b}_0|}{P_f(0) \phi_f(0)} \right)$$

$$|V_{cb}|_{\text{BGL}} = (40.9 \pm 0.3 \pm 1.0 \pm 0.6) \times 10^{-3}$$

$$|V_{cb}|_{\text{CLN}} = (40.4 \pm 0.3 \pm 1.0 \pm 0.6) \times 10^{-3}$$

↑
Slow pion eff. plays
leading role in syst.

↑
Input from LQCD at
zero-recoil $\mathcal{F}(1)$

Measurement of τ mass

PRD 108, 032006 (2023)




Pseudomass method:

$$m_\tau^2 = (p_h + p_\nu)^2$$

$$= 2 E_h (E_\tau - E_h) + m_h^2 - 2 |\vec{p}_h| (E_\tau - E_h) \cos(\vec{p}_h, \vec{p}_\nu)$$

The direction of the neutrino is not known, since $\cos(\vec{p}_h, \vec{p}_\nu) \leq 1$

$$\text{Pseudomass: } M_{\min} = \sqrt{M_{3\pi}^2 + 2(\sqrt{s}/2 - E_{3\pi}^*)(E_{3\pi}^* - p_{3\pi}^*)} \leq m_\tau$$



$$m_\tau = 1777.09 \pm 0.08 \pm 0.11 \text{ MeV}/c^2$$

World's best!



Smaller data

BUT better statistical precision!

Measurement of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$

Preliminary

- Reconstruct B_{tag} with FEI
- Yield signals from simultaneous fit to M_{bc} (SR and SB)





Shape increase at $B\bar{B}^$ threshold. Suggestive of something?*

Measurement of $B \rightarrow K^* \gamma$

Preliminary

- CP and isospin asymmetries are theoretically clean to cancellation of form factor uncertainties.
- Sensitive to NP.
- Simultaneous fit to M_{bc} and ΔE for different modes



$$\mathcal{A}_{CP} = \frac{\Gamma(\bar{B} \rightarrow \bar{K}^* \gamma) - \Gamma(B \rightarrow K^* \gamma)}{\Gamma(\bar{B} \rightarrow \bar{K}^* \gamma) + \Gamma(B \rightarrow K^* \gamma)}$$

$$\Delta_{0+} = \frac{\Gamma(B^0 \rightarrow K^{*0} \gamma) - \Gamma(B^+ \rightarrow K^{*+} \gamma)}{\Gamma(B^0 \rightarrow K^{*0} \gamma) + \Gamma(B^+ \rightarrow K^{*+} \gamma)},$$

Results:

$$\mathcal{B}[B^0 \rightarrow K^{*0} \gamma] = (4.16 \pm 0.10 \pm 0.11) \times 10^{-5},$$

$$\mathcal{B}[B^+ \rightarrow K^{*+} \gamma] = (4.04 \pm 0.13 \pm 0.13) \times 10^{-5},$$

$$\mathcal{A}_{CP}[B^0 \rightarrow K^{*0} \gamma] = (-3.2 \pm 2.4 \pm 0.4)\%,$$

$$\mathcal{A}_{CP}[B^+ \rightarrow K^{*+} \gamma] = (-1.0 \pm 3.0 \pm 0.6)\%,$$

$$\Delta \mathcal{A}_{CP} = (2.2 \pm 3.8 \pm 0.7)\%, \text{ and}$$

$$\Delta_{0+} = (5.1 \pm 2.0 \pm 1.5)\%,$$