

清华大学  
Tsinghua University

中国物理学会高能物理分会  
HIGH ENERGY PHYSICS BRANCH OF CPS

14/8/2024-18/8/2024

# $\phi$ production in Au+Au collisions at $\sqrt{s_{NN}}=19.6$ , 14.6, and 7.7 GeV with the STAR experiment

Weiguang Yuan  
(for the STAR Collaboration)  
Tsinghua University

- Motivation
- Experimental analysis
- Results
  - ✓  $p_T$  spectra
  - ✓ Rapidity spectra
  - ✓ Nuclear modification factors
  - ✓  $\phi/K^-$ ,  $\Omega/\phi$  ratio
- Summary

Supported in part by the



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

# Motivation


## ➤ Beam Energy Scan (BES) program:

- Search for the critical point
- Search for the first-order phase transition
- Search for the threshold of QGP formation

## ➤ Energy dependency of QGP signature

- Strange baryon-to-meson ratio can be utilized to understand hadronization mechanism
- $R_{CP}$  may give insight into the parton energy loss


$$R_{CP} = \frac{[(dN/dp_T)/\langle N_{coll} \rangle]_{\text{central}}}{[(dN/dp_T)/\langle N_{coll} \rangle]_{\text{peripheral}}}$$



# Motivation: Why study $\phi$ ?


## ➤ Long lifetime and small reaction cross-section

- Lifetime:  $41 \text{ fm}/c \rightarrow$  the decay products are not disturbed by the late hadronic rescatterings
- Small cross-section  $\rightarrow \phi$  is more likely to remain unaffected by the later stage of hadronic interactions




## ➤ Enhanced yield in QGP

- Restored chiral symmetry in QGP  $\rightarrow$  the mass of  $s$  and  $\bar{s}$  is smaller and  $s\bar{s}$  pairs can be produced in large quantities by gluon fusion and light  $q\bar{q}$  pairs annihilation




# Motivation: Why $\Omega/\phi$ ratio?

STAR: Phys. Rev. C 93 (2016) 2, 021903




- $\Omega/\phi$  ratio for BES-I energies

- At  $\sqrt{s_{NN}} = 200$  GeV, the enhanced  $\Omega/\phi$  ratios from p+p collision to central Au+Au collision may indicate the existence of QGP
- For BES-I energy, the uncertainties are too large to draw a firm conclusion below 11.5 GeV.



- $\Omega/\phi$  ratio for 200 GeV energies


# Motivation: More precise measurement



## BES-II vs BES-I

$\sqrt{s_{NN}}$ (GeV)	Events BES-I ( $10^6$ )	Events BES-II ( $10^6$ )
7.7	3	<b>45</b>
<b>9.2</b>	-	<b>78</b>
11.5	7	<b>110</b>
14.5	20	<b>178</b>
<b>17.3</b>	-	<b>116</b>
19.6	15	<b>270</b>
27	30	<b>220</b>


- BES-II compared to BES-I:  $\sim 10$ - $18$  times larger statistics  
 → higher precision and wider  $\mu_B$  coverage



- Large and uniform acceptance
- Excellent particle identification


iTPC upgrade

- $|\eta|: 1.0 \rightarrow 1.5$
- Improved  $dE/dx$  resolution
- $p_T$  threshold:  $120\text{MeV} \rightarrow 60\text{MeV}$




# Particle identification and reconstruction

Au+Au  $\sqrt{s_{NN}} = 19.6$  GeV




➤ Particle identification with upgraded TPC and bTOF



$\phi \rightarrow K^+ + K^-$  (49.1%)

- Combinational background have been removed by Mix-Event Method.



# $p_T$ spectra of $\phi$ at $\sqrt{s_{NN}} = 19.6, 14.6$ and $7.7$ GeV



➤ For  $\phi$ : Levy function fit to extrapolate down to zero  $p_T$




# $p_T$ spectra of $\Omega$ at $\sqrt{s_{NN}} = 19.6, 14.6$ and $7.7$ GeV




➤ For  $\Omega$ : Boltzmann function fit to extrapolate down to zero  $p_T$

# Rapidity spectra of $\phi$



- Rapidity spectra of  $\phi$  are **Gaussian-like** distributions
- Rapidity distribution **become wider with increasing energy**



# Nuclear modification factor



$$R_{CP} = \frac{[(dN/dp_T)/\langle N_{coll} \rangle]_{\text{central}}}{[(dN/dp_T)/\langle N_{coll} \rangle]_{\text{peripheral}}}$$



- $R_{CP} < 1$  for higher  $p_T$  at  $\sqrt{s_{NN}} = 200$  GeV  $\rightarrow$  Partonic energy loss in the QGP medium
- $R_{CP} > 1$  for higher  $p_T$  at  $\sqrt{s_{NN}} = 19.6$  GeV and lower energies  $\rightarrow$  Cronin-type interactions, radial flow and/or coalescence hadronization
- $R_{CP}$  of  $\phi$  at  $\sqrt{s_{NN}} = 7.7$  GeV is significantly **different** from that at  $\sqrt{s_{NN}} = 14.6$  and 19.6 GeV

# Centrality dependence of $\phi$ yields (dN/dy)






- Fit function:  $(dN/dy)/(N_{part}/2) = k \times N_{part}^{\alpha-1}$
- $\alpha$  parameter for  $\phi$  is slightly larger than that for  $\Lambda$ , K and **less than UrQMD predictions**

# Centrality and Energy dependence of $\phi/K^-$ ratio



- The  $\phi/K^-$  ratio exhibits no clear dependency on centrality or energy across the range of  $\sqrt{s_{NN}} = 7.7$  to 19.6 GeV
- The  $\phi/K^-$  ratio **reaches the GCE limit** at  $\sqrt{s_{NN}} = 7.7, 14.6$  and 19.6 GeV

# $\Omega(sss)/\phi(s\bar{s})$ ratio



- Similar to the observation at  $\sqrt{s_{NN}} = 200$  GeV, the  $\Omega/\phi$  ratio increases from peripheral to central collisions at intermediated  $p_T$ , which is **compatible with the existence of QGP at  $\sqrt{s_{NN}} \geq 7.7$  GeV**

# Summary

## Summary:



- The  $p_T$ , centrality and rapidity dependences of  $\phi$  production at  $\sqrt{s_{NN}} = 7.7, 14.6$  and  $19.6$  GeV have been presented
- Hadronic transport model UrQMD cannot describe centrality dependence well from  $\sqrt{s_{NN}} = 7.7$  to  $19.6$  GeV
- For  $\phi/K^-$  ratio, both GCE and CE calculations **are consistent with** the data across the range of  $\sqrt{s_{NN}} = 7.7$  to  $19.6$  GeV
- The  $\phi R_{CP}$  at low energies shows the radial flow and quark coalescence effects
- The  $\Omega(sss)/\phi(s\bar{s})$  ratio **is compatible with the existence of QGP signals at  $\sqrt{s_{NN}} \geq 7.7$  GeV**

## Outlook:

- The measurements in other BES-II datasets at different energies will be conducted
- Other BES-II energies:  $\sqrt{s_{NN}} = 9.2, 11.5$  and  $17.3$  GeV

**Thanks!!**

# Back up: Rapidity spectra of $\phi$ in UrQMD at 19.6 GeV





## Back up:

## Coalescence model

➤ According to recombination model, **if exist QGP**, mesons and baryons can be **formed by combining quarks**.

• The yield distribution of a produced meson with momentum  $p$ :

$$p^0 \frac{dN_B}{dp} = \int \frac{dp_1}{p_1} \frac{dp_2}{p_2} \frac{dp_3}{p_3} F_{qq'q''}(p_1, p_2, p_3) R_B(p_1, p_2, p_3, p).$$

• The yield distribution of a produced baryon with momentum  $p$ :

$$p^0 \frac{dN_M}{dp} = \int \frac{dp_1}{p_1} \frac{dp_2}{p_2} F_{q\bar{q}'}(p_1, p_2) R_M(p_1, p_2, p),$$

$p_0$ 为E



$$F_{s\bar{s}} = \mathcal{T}_s \mathcal{T}_s + \mathcal{T}_s \mathcal{S}_s + \{\mathcal{S}_s \mathcal{S}_s\},$$

$$F_{sss} = \mathcal{T}_s \mathcal{T}_s \mathcal{T}_s + \mathcal{T}_s \mathcal{T}_s \mathcal{S}_s + \mathcal{T}_s \{\mathcal{S}_s \mathcal{S}_s\} + \{\mathcal{S}_s \mathcal{S}_s \mathcal{S}_s\}.$$



$$\frac{dN_\phi}{pdp} = \frac{g_\phi}{pp_0} F_{s\bar{s}}(p/2, p/2),$$

$$\frac{dN_\Omega}{pdp} = \frac{g_\Omega}{pp_0} F_{sss}(p/3, p/3, p/3),$$

The yield distribution of  $\Omega$  and  $\phi$ .

$$\mathcal{T}(p_1) = p_1 \frac{dN_q^{\text{th}}}{dp_1} = C p_1 \exp(-p_1/T),$$

$$\mathcal{S}(p_2) = \xi \sum_i \int_{k_0}^{\infty} dk k f_i(k) S_i(p_2/k).$$

$\mathcal{T}_s$  is the thermal parton distribution comes from QGP.

$\mathcal{S}_s$  is the shower parton distribution comes from hard scattering.

# Back up:

# Coalescence model

➤ Just consider the contribution of thermal partons:

$$\frac{dN_\phi}{pdp} = g_\phi C_s^2 \frac{p}{4p_0} e^{-p/T_s},$$



$$\frac{dN_\Omega}{pdp} = g_\Omega C_s^3 \frac{p^2}{27p_0} e^{-p/T_s},$$




$$R_{\Omega/\phi}^{\text{th}}(p) = \frac{4g_\Omega C_s}{27g_\phi} p,$$


Ratio of  $\Omega/\phi$  is proportional to  $p$ .

➤ The  $\Omega/\phi$  ratio distribution with  $p_T$



The model prediction result of  $\Omega/\phi$  ratio in QGP!





No obvious Centrality dependence  
for 7.7 GeV.