



Towards ultimate fragmentation functions at future lepton colliders

Bin Zhou

INPAC, Shanghai Jiao Tong University

Based on 2407.10059 with Jun Gao

第十四届全国粒子物理学术会议, August 15, 2024

Outline

① Introduction

② Pseudo–data generation

③ Results and discussions

④ Summary

Outline

① Introduction

② Pseudo–data generation

③ Results and discussions

④ Summary

Fragmentation functions

- Fragmentations of quarks and gluons into hadrons have been the central topic of QCD since only hadrons are observed experimentally
- In its simplest form, fragmentation functions (FFs) describe number density of the identified hadron wrt the fraction of momentum of the initial parton it carries, as measured in single inclusive hadron production, e.g., from single-inclusive annihilation (SIA), semi-inclusive DIS (SIDIS), pp collisions¹




Figure 1: Representation of the scattering amplitude for processes whose QCD description involves FFs: Single-inclusive hadron production in (a) e^+e^- annihilation, (b) deep-inelastic lepton-nucleon scattering, (c) proton-proton scattering.

- As the FFs are related to a nonperturbative aspect of QCD, they cannot be calculated with precision using theoretical methods and in general extracted from fits to a variety of experimental data.

¹A. Metz and A. Vossen, “Parton Fragmentation Functions”, *Prog. Part. Nucl. Phys.* **91**, 136–202 (2016).

Current situations

Many studies have already been conducted on a comprehensive fit involving various data samples to extract FFs.

- DSS, AKK, NNFF, MAPFF, JAM, and NPC23, which combine data sets from SIA, SIDIS, or pp collisions
 - HKNS, which only includes SIA data
- } NLO in QCD

Furthermore, determinations of FFs at next-to-next-to-leading order (NNLO) with SIA data only^{1 2 3 4} and at approximate NNLO with SIA and SIDIS data^{5 6} have also been made.

In the present study, we focus on the fit based solely on the data from the lepton collider.

- the separation between quark and antiquark FFs
- the gluon FF

¹V. Bertone et al. (NNPDF), “A determination of the fragmentation functions of pions, kaons, and protons with faithful uncertainties”, *Eur. Phys. J. C* **77**, 516 (2017).

²M. Soleymaninia et al., “First QCD analysis of charged hadron fragmentation functions and their uncertainties at next-to-next-to-leading order”, *Phys. Rev. D* **98**, 074002 (2018).

³M. Soleymaninia et al., “Impact of unidentified light charged hadron data on the determination of pion fragmentation functions”, *Phys. Rev. D* **99**, 034024 (2019).

⁴M. Soleymaninia et al., “Simultaneous extraction of fragmentation functions of light charged hadrons with mass corrections”, *Phys. Rev. D* **103**, 054045 (2021).

⁵I. Borsa et al., “Towards a Global QCD Analysis of Fragmentation Functions at Next-to-Next-to-Leading Order Accuracy”, *Phys. Rev. Lett.* **129**, 012002 (2022).

⁶R. Abdul Khalek et al. (MAP (Multi-dimensional Analyses of Partonic distributions)), “Pion and kaon fragmentation functions at next-to-next-to-leading order”, *Phys. Lett. B* **833**, 137456 (2022) ➔

Opportunities with future lepton colliders

The run of future lepton colliders, such as ILC¹, CEPC², and FCC–ee³, will collect high-quality and diverse data on hadron production, sufficiently for precision determination of fragmentation functions alone

Particle	E _{c.m.} (GeV)	Years	SR Power (MW)	Lumi. /IP (10 ³⁴ cm ⁻² s ⁻¹)	Integrated Lumi. /yr (ab ⁻¹ , 2 IPs)	Total Integrated L (ab ⁻¹ , 2 IPs)	Total no. of events
H*	240	10	50	8.3	2.2	21.6	4.3 × 10 ⁶
			30	5	1.3	13	2.6 × 10 ⁶
Z	91	2	50	192**	50	100	4.1 × 10 ¹²
			30	115**	30	60	2.5 × 10 ¹²
W	160	1	50	26.7	6.9	6.9	2.1 × 10 ⁸
			30	16	4.2	4.2	1.3 × 10 ⁸
t̄t	360	5	50	0.8	0.2	1.0	0.6 × 10 ⁶
			30	0.5	0.13	0.65	0.4 × 10 ⁶

CEPC operation plans

Working point	Luminosity (10 ³⁴ cm ⁻² s ⁻¹)	Tot. lum./year (ab ⁻¹)/year	Goal (ab ⁻¹)	Run time (years)
Z first two years	100	24	150	4
Z other years	200	48		
W	25	6	10	1–2
H	7.0	1.7	5	3
RF reconfiguration				1
t̄t 350 GeV (first year)	0.8	0.20	0.2	1
t̄t 365 GeV	1.4	0.34	1.5	4


FCC–ee operation plans

$\int \mathcal{L} dt$ [fb ⁻¹]			
\sqrt{s}	G20	H20	I20
250 GeV	500	2000	500
350 GeV	200	200	1700
500 GeV	5000	4000	4000
			Snow
			1150
			200
			1600


ILC operation plans

¹A. Aryshev et al. (ILC International Development Team), “The International Linear Collider: Report to Snowmass 2021”, (2022).²W. Abdallah et al. (CEPC Study Group), “CEPC Technical Design Report – Accelerator (v2)”, (2023).³A. Abada et al. (FCC), “FCC-ee: The Lepton Collider: Future Circular Collider Conceptual Design Report Volume 2”, Eur. Phys. J. ST **228**, 261–623 (2019).

Opportunities with future lepton colliders



FFs-sensitive processes



LO cross section for various light quarks and antiquarks production

- producing $q\bar{q}$ samples at various energies with kinematic cuts, important for light quark flavor and charge separation
- heavy-quark enriched samples and gluon samples from Higgs hadronic decays
- further quark flavor and charge separation from W -boson production with hadronic decays

Outline

① Introduction

② Pseudo–data generation

③ Results and discussions

④ Summary

Pseudo–data generation

If σ_i^{th} is the theoretical cross-section for bin i of a given process, then the central values of the pseudo–data σ_i^{exp} is constructed by means of

$$\sigma_i^{\text{exp}} = \sigma_i^{\text{th}} \times (1 + r_i \cdot \delta_{\text{tot}, i}^{\text{exp}} + s \cdot \delta_{\text{cor}}^{\text{exp}})$$

- σ_i^{th} : obtained using FMNLO¹ with the NPC23 set as input. Given that the present study is based on pseudo–data, we focus on the analysis carried out at NLO in QCD
- r_i, s : univariate Gaussian random numbers
- $\delta_{\text{tot}, i}^{\text{exp}}$: defined as $\left((\delta_{\text{stat}, i}^{\text{exp}})^2 + (\delta_{\text{sys}, i}^{\text{exp}})^2 \right)^{1/2}$, where statistical uncertainties are evaluated from the expected number of events per bin and systematic uncertainties are set to be those of the baseline measurement from SLD at Z–pole
- $\delta_{\text{cor}}^{\text{exp}}$: derived from the reference SLD measurement at Z–pole, 1%

¹C. Liu et al., “Automated calculation of jet fragmentation at NLO in QCD”, JHEP 09, 108 (2023).

Selection of data

- $N \times 2$
- $W^- W^{+,*}$ production
- CEPC: 1831 points in total

$e^+ e^-$ annihilation									
\sqrt{s} (GeV)	luminosity (ab^{-1})			final state	kinematic cuts	hadrons	N_{pt}		
	CEPC	FCC-ee	ILC						
91.2	60	150	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	132		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	65		
160	4.2	-	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	168		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	83		
161	-	10	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	168		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	83		
240	13	5	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	186		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	92		
250	-	-	2	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	186		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	92		
350	-	0.2	0.2	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	198		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	98		
360	0.65	-	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	198		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	98		
365	-	1.5	-	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	198		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	98		
500	-	-	4	$q\bar{q}$	$\cos(\theta) > 0$	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	198		
				$c\bar{c}/bb$	-	$\pi^\pm, K^\pm, p/\bar{p}$	98		
W boson decay channels									
\sqrt{s} (GeV)	# events (million)			final state	kinematic cuts	hadrons	N_{pt}		
	CEPC	FCC-ee	ILC						
80.419	116	68	62	$W^- W^{**} \rightarrow W^- q\bar{q}$	-	$\pi^+, \pi^-, K^+, K^-, p, \bar{p}$	120		
	58	34	31	$W^- W^{**} \rightarrow W^- c\bar{s}$					
Higgs boson decay channels									
\sqrt{s} (GeV)	# events (million)			final state	kinematic cuts	hadrons	N_{pt}		
	CEPC	FCC-ee	ILC						
125	0.23	0.09	0.07	gg	-	$\pi^\pm, K^\pm, p/\bar{p}$	77		
	0.08	0.03	0.02	$c\bar{c}$					
	1.53	0.59	0.47	bb					

A summary of the main features of the CEPC, FCC-ee, and ILC pseudo-data generated for the present study

Fit of FFs

Choosing a framework^{1,2} on global analysis of fragmentation functions to identified charged hadrons, including charged pion, kaon and proton, using pseudo–data from future lepton colliders

parametrization of FFs to charged pion/kaon/ proton at initial scale (Q=5 GeV):

$$z D_i^h(z, Q_0) = z^{\alpha_i^h} (1-z)^{\beta_i^h} \exp \left(\sum_{n=0}^m a_{i,n}^h (\sqrt{z})^n \right)$$

parton-to- π^+	favored	α	β	a_0	a_1	a_2	d.o.f.
u	Y						5
$d \simeq u$	Y	-	-	-	-		1
$\bar{u} = d$	N				x		4
$s = \bar{s} \simeq \bar{u}$	N	-			x		3
$c = \bar{c}$	N				x		4
$b = \bar{b}$	N				x		4
g	N		F				4

parton-to- K^+	favored	α	β	a_0	a_1	a_2	d.o.f.
u	Y					x	4
$\bar{s} \simeq u$	Y	-	-	-	x		1
$\bar{u} = d = \bar{d} = s$	N				x		4
$c = \bar{c}$	N				x		4
$b = \bar{b}$	N				x		4
g	N		F		x		3

parton-to- p	favored	α	β	a_0	a_1	a_2	d.o.f.
$u = 2d$	Y					x	4
$\bar{u} = d = s = \bar{s}$	N			x	x		3
$c = \bar{c}$	N				x		4
$b = \bar{b}$	N				x		4
g	N		F		x		3

- a joint determination of FFs to charged pion, kaon and proton at NLO in QCD (63 parameters) including estimation of uncertainties with Hessian sets
- apply a strong selection criteria on the kinematics of fragmentation processes to ensure validity of leading-twist factorization and perturbative calculations ($z > 0.01$ and $E_h > 4\text{GeV}$)
- including theory uncertainties (residual scale variations) into the covariance matrix

¹J. Gao et al., “Simultaneous Determination of Fragmentation Functions and Test on Momentum Sum Rule”, *Phys. Rev. Lett.* **132**, 261903 (2024).

²J. Gao et al., “Global analysis of fragmentation functions to charged hadrons with high-precision data from the LHC”, (2024).

Outline

① Introduction

② Pseudo–data generation


③ Results and discussions

④ Summary

The constraints to FFs from CEPC

We are only interested in the relative reduction of the FFs uncertainties once the data from future lepton colliders is included in the fit.


- the total χ^2 is 1659.2 for a total number of data points of 1831, resulting in $\chi^2/N_{\text{pt}} = 0.91$
- we observe a marked reduction in the FFs uncertainties almost in all cases
- a slight increase of uncertainties for gluon FF to π^+ in the small x region due to the insufficient number of data points available



A comparison of the baseline NPC23 FFs with those based on CEPC pseudo-data for π^+ , K^+ , and p ↗ ↘

The constraints to FFs from future lepton colliders

The pseudo-data from CEPC, FCC-ee and ILC have a comparable impact on the uncertainties associated with the FFs




A comparison of the relative errors among the FFs obtained from CEPC, FCC-ee, and ILC measurements, respectively



Alternative fit

We utilize the FFs from CEPC as a case study to investigate the impact of individual data sets on the extracted FFs

- the $W^- W^{+,*}$ production data set has strong constraints on the FFs of light quarks
- the measurements from the decay of the Higgs boson exert a profound influence on the FFs of gluon
- the data sets on quark pair production at lepton colliders show a larger impact compared to the $W^- W^{+,*}$ production data set for the FFs of light quarks.



The effect of the higher-order QCD corrections





- The inclusion of NNLO corrections results in a further reduction in scale uncertainties.
- Furthermore, the resulting predictions using the NNFF10_PIp_nnlo set are comparable to those obtained using the NPC23_PIp_nlo set ¹.

¹<http://fmnlo.sjtu.edu.cn/~fmnlo/>

The effect of the higher-order QCD corrections


Alternative fits were conducted without including the theoretical uncertainties to discuss the effects of the higher-order corrections.



- A stronger constraint on the FFs when higher-order corrections are included in the global QCD fit

The constraints from three-jet production

The experiment is also capable of measuring the relevant observable in three-jet production.



A comparison between the scale uncertainties and the Hessian uncertainties for the distribution of the hadron energy fraction in each jet of three-jet events

- The NNLO corrections will be needed in the future to effectively include the data into a global analysis of FFs

Outline

① Introduction

② Pseudo–data generation

③ Results and discussions

④ Summary

Summary

- In this work, we study the constraining power of future lepton colliders on FFs to light charged hadrons from quarks and gluon
- We find that either CEPC, FCC-ee or ILC can significantly reduce the uncertainties of FFs in a wide kinematic range, compared to the NPC23 set
- We also discuss the impact of higher-order QCD corrections, and the potential constraints from measurements of three-jet production.
- Currently, hadron production cross sections at NNLO in QCD are available for SIDIS, SIA, and decays of the Higgs boson to gluons, and have been implemented into the FMNLO program

Thank you for listening!