

## Probing Neutral Triple Gauge Couplings via $Z\gamma(\ell+\ell-\gamma)$ Production at $e+e-$ Colliders

Ref. <https://arxiv.org/abs/2404.15937>

(Accepted by Frontier of Physics)

Neutral triple gauge couplings (nTGCs) are absent in the Standard Model (SM) and at the dimension-6 level in the Standard Model Effective Field Theory (SMEFT), arising first from dimension-8 operators. As such, they provide a unique window for probing new physics beyond the SM. These dimension-8 operators can be mapped to nTGC form factors whose structure is consistent with the spontaneously-broken electroweak gauge symmetry of the SM. In this work, we study the probes of nTGCs in the reaction  $e+e-\rightarrow Z\gamma$  with  $Z\rightarrow\ell+\ell-(\ell=e,\mu)$  at an  $e+e-$  collider. We perform a detector-level simulation and analysis of this reaction at the Circular Electron Positron Collider (CEPC) with collision energy  $\sqrt{s} = 240$  GeV and an integrated luminosity of  $5\text{ ab}^{-1}$ . We present the sensitivity limits on probing the new physics scales of dimension-8 nTGC operators via measurements of the corresponding nTGC form factors.

**Primary author:** LIU, Danning (TDLI, SJTU)

**Co-authors:** ELLIS, John (King's College London); HE, Hong-Jian (Shanghai Jiao Tong University); XIAO, Rui-Qing (SJTU)

**Presenter:** LIU, Danning (TDLI, SJTU)

**Session Classification:** 分会场一

**Track Classification:** TeV 物理和超出标准模型新物理