

Recent results on charmed baryon at BESIII and LHCb

Xiao-Rui Lyu (吕晓睿)

xiaorui@ucas.ac.cn

University of Chinese Academy Sciences






Outline

- Introduction to the charmed baryons
- BESIII progress in studying the Λ_c^+
- Recent LHCb results on charmed baryons
- Ongoing progress and future plan at BESIII
- Summary

Quark model picture

a heavy quark (c) with an unexcited spin-zero diquark ($u-d$)

→ *diquark correlation is enhanced by weak Color Magnetic Interaction with a heavy quark.*



→ Charmed meson ($D^+[c\bar{d}]$)
 $m_d \ll m_c \rightarrow$ **quark + heavy quark**

→ Strange baryons ($\Lambda[uds]$)
 $m_u, m_d \approx m_s \rightarrow$ **(qqq)** uniform

→ Charmed baryon ($\Lambda_c[udc]$)
 $m_u, m_d \ll m_c \rightarrow$ **diquark + quark**
(qq) (Q)

In some sense, more reliable prediction of heavy-light quark transition without dealing with light degrees of freedom that have net spin or isospin.

Λ_c^+ may provide complementary powerful test on internal dynamics to D/Ds does



Experimental studies on Λ_c^+ until 2014

Before 2014, the c -ed baryons have been produced and studied at many experiments, notably fixed-target experiments (such as FOCUS and SELEX) and e^+e^- B-factories (ARGUS, CLEO, BABAR, and BELLE).

- ✓ Total branching fraction $\sim 60\%$
- ✓ Lots of unknown decay channels
- ✓ Quite large uncertainties ($>20\%$)
- ✓ Most BF's are measured relative to $\Lambda_c^+ \rightarrow pK^-\pi^+$


Large uncertainties in experiment
 → slow development in theory

Λ_c^+ data in PDG2015

Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/ c)
Hadronic modes with a p: $S = -1$ final states			
$p\bar{K}^0$	(3.21 ± 0.30) %		
$pK^-\pi^+$	(6.84 ^{+0.32} _{-0.40}) %		
$p\bar{K}^*(892)^0$	[q] (2.13 ± 0.30) %		
$\Delta(1232)^{++}K^-$	(1.18 ± 0.27) %		22.5%
$\Lambda(1520)\pi^+$	[q] (2.4 ± 0.6) %		25.0%
$pK^-\pi^+$ nonresonant	(3.8 ± 0.4) %		10.5%
$p\bar{K}^0\pi^0$	(4.5 ± 0.6) %		13.3%
$p\bar{K}^0\eta$	(1.7 ± 0.4) %		23.5%
$p\bar{K}^0\pi^+\pi^-$	(3.5 ± 0.4) %		11.4%
$pK^-\pi^+\pi^0$	(4.6 ± 0.8) %		13.0%
$pK^*(892)^-\pi^+$	[q] (1.5 ± 0.5) %		33.3%
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(5.0 ± 0.9) %		18.0%
$\Delta(1232)K^*(892)$	seen		
$pK^-\pi^+\pi^+\pi^-$	(1.5 ± 1.0) × 10 ⁻³		66.7%
$pK^-\pi^+\pi^0\pi^0$	(1.1 ± 0.5) %		45.4%
Hadronic modes with a p: $S = 0$ final states			
$p\pi^+\pi^-$	(4.7 ± 2.5) × 10 ⁻³		45.4%
$p f_0(980)$	[q] (3.8 ± 2.5) × 10 ⁻³		53.2%
$p\pi^+\pi^+\pi^-\pi^-$	(2.5 ± 1.6) × 10 ⁻³		64.0%
pK^+K^-	(1.1 ± 0.4) × 10 ⁻³		36.4%
$p\phi$	[q] (1.12 ± 0.23) × 10 ⁻³		
pK^+K^- non- ϕ	(4.8 ± 1.9) × 10 ⁻⁴		
Hadronic modes with a hyperon: $S = -1$ final states			
$\Lambda\pi^+$	(1.46 ± 0.13) %		8.9%
$\Lambda\pi^+\pi^0$	(5.0 ± 1.3) %		26.0%
$\Lambda\rho^+$	< 6 %	CL=95%	
$\Lambda\pi^+\pi^+\pi^-$	(3.59 ± 0.28) %		7.8%
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow$	(1.0 ± 0.5) %		20.0%
$\Lambda\pi^+$			
$\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow$	(7.5 ± 1.4) × 10 ⁻³		18.7%
$\Lambda\pi^-$			



Charmed baryon thresholds



BESIII energy upgrades:
 4.6 GeV (Phase I, 2014)
 → 4.95 GeV (Phase II, 2021)
 → 5.6 GeV (Phase III, planned in 2024)



Phase I

In spring of 2013, a proposal of Λ_c^+ data taking was discussed and later approved inside the collaboration.


BESIII Near threshold production at BESIII



In 2014, BESIII took (only!) 35 days to run at 4.6GeV and collected $\sim 500/\text{pb}$ data.

Energy(GeV)	lum.(1/pb)
4.575	47.67
4.580	8.54
4.590	8.16
4.600	566.93



Corresponds to 0.1M Λ_c pairs



Measurement using the threshold pair-productions via e^+e^- annihilations is unique: *the most simple and straightforward*

First time to systematically study charmed baryon at threshold!

The absolute BF can be obtained by the ratio of DT yields to ST yields



- High efficiency and clean background
- Absolute measurement with many systematics cancel out
- Missing-mass technique: K_L /neutron, neutrino, ...
- Good photon resolution: Σ , Ξ , π^0 , ...

$$B_i = \frac{N_{ij}^{DT}}{N_j^{ST}} \frac{\epsilon_j}{\epsilon_{ij}}$$

Published 17 papers
(7 PRLs)

- A series of **precise absolute BF measurements**: hadronic, semi-leptonic and inclusive decays
- **Observation** of decays into neutron $\Lambda_c^+ \rightarrow nK_s\pi^+, \Sigma^-\pi^+\pi^+\pi^0$
- **Observation** of Cabbibo-suppressed decay $\Lambda_c^+ \rightarrow p\pi^+\pi^-$
- **First evidence** of Cabbibo-suppressed decay $\Lambda_c^+ \rightarrow p\eta$
- **First measurements** of many decay asymmetries
- Determination of Λ_c^+ spin
- Threshold cross section and form factors of Λ_c^+ pairs

Very productive for the data set taken in 35 days!

Hadronic decay

2014 : 0.567 fb⁻¹ at 4.6 GeV

$\Lambda_c^+ \rightarrow pK^-\pi^+ + 11 \text{ CF modes}$	PRL 116, 052001 (2016)
$\Lambda_c^+ \rightarrow pK^+K^-, p\pi^+\pi^-$	PRL 117, 232002 (2016)
$\Lambda_c^+ \rightarrow nK_s\pi^+$	PRL 118, 12001 (2017)
$\Lambda_c^+ \rightarrow p\eta, p\pi^0$	PRD 95, 111102(R) (2017)
$\Lambda_c^+ \rightarrow \Sigma^-\pi^+\pi^+\pi^0$	PLB 772, 388 (2017)
$\Lambda_c^+ \rightarrow \Xi^{0(*)}K^+$	PLB 783, 200 (2018)
$\Lambda_c^+ \rightarrow \Lambda\eta\pi^+$	PRD 99, 032010 (2019)
$\Lambda_c^+ \rightarrow \Sigma^+\eta, \Sigma^+\eta'$	CPC 43, 083002 (2019)
$\Lambda_c^+ \rightarrow \text{BP decay asymmetries}$	PRD 100, 072004 (2019)
$\Lambda_c^+ \rightarrow pK_s\eta$	PLB 817, 136327 (2021)
Λ_c^+ spin determination	PRD 103, L091101(2021)

Semi-leptonic decay

$\Lambda_c^+ \rightarrow \Lambda e^+\nu_e$	PRL 115, 221805(2015)
$\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu$	PLB 767, 42 (2017)


Inclusive decay

$\Lambda_c^+ \rightarrow \Lambda X$	PRL121, 062003 (2018)
$\Lambda_c^+ \rightarrow e^+ X$	PRL 121 251801(2018)
$\Lambda_c^+ \rightarrow K_s^0 X$	EPJC 80, 935 (2020)

Production

$\Lambda_c^+\Lambda_c^-$ cross section	PRL 120,132001(2018)
----------------------------------------	----------------------

- Absolute BF of Λ_c^+ decays are still not well determined since its discovery 30 years ago. PDG2014: $\delta B/B \sim 26\%$; BELLE2014: $\delta B/B \sim 5.3\%$
- Double tag technique is applied to control systematics



PRL 116, 052001 (2016)

Mode	This work (%)	PDG (%)	BELLE β
ρK_S^0	$1.52 \pm 0.08 \pm 0.03$	1.15 ± 0.30	
$\rho K^- \pi^+$	$5.84 \pm 0.27 \pm 0.23$	5.0 ± 1.3	$6.84 \pm 0.24^{+0.21}_{-0.27}$
$\rho K_S^0 \pi^0$	$1.87 \pm 0.13 \pm 0.05$	1.65 ± 0.50	
$\rho K_S^0 \pi^+ \pi^-$	$1.53 \pm 0.11 \pm 0.09$	1.30 ± 0.35	
$\rho K^- \pi^+ \pi^0$	$4.53 \pm 0.23 \pm 0.30$	3.4 ± 1.0	
$\Lambda \pi^+$	$1.24 \pm 0.07 \pm 0.03$	1.07 ± 0.28	
$\Lambda \pi^+ \pi^0$	$7.01 \pm 0.37 \pm 0.19$	3.6 ± 1.3	
$\Lambda \pi^+ \pi^- \pi^+$	$3.81 \pm 0.24 \pm 0.18$	2.6 ± 0.7	
$\Sigma^0 \pi^+$	$1.27 \pm 0.08 \pm 0.03$	1.05 ± 0.28	
$\Sigma^+ \pi^0$	$1.18 \pm 0.10 \pm 0.03$	1.00 ± 0.34	
$\Sigma^+ \pi^+ \pi^-$	$4.25 \pm 0.24 \pm 0.20$	3.6 ± 1.0	
$\Sigma^+ \omega$	$1.56 \pm 0.20 \pm 0.07$	2.7 ± 1.0	

- a global least square fit to 12 hadronic modes [Chin. Phys. C37(2013)106201]

- ✓ First direct measurement on Λ_c BFs at threshold
- ✓ $B(pK^- \pi^+)$: BESIII precision comparable with Belle's
- ✓ Improved precisions of the other 11 modes significantly

So far, the mostly cited BESIII charm paper



BF for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$



- $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ is a $c \rightarrow s l^+ \nu_l$ dominated process.
- Urgently needed for LQCD calculations.
- No direct absolute measurement for $\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)$ available.

$$\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (2.1 \pm 0.6)\% \quad \text{PDG 2014}$$

11 hadronic single tag modes are used



$$\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (3.63 \pm 0.38 \pm 0.20)\%$$

- First absolute measurement of the semi-leptonic decay
- Statistics limited
- Best precision to date: twofold improvement



First Lattice calculation on charmed baryon SL decays

PRL **118**, 082001 (2017)

PHYSICAL REVIEW LETTERS

week ending
24 FEBRUARY 2017

$\Lambda_c \rightarrow \Lambda \ell^+ \nu_\ell$ Form Factors and Decay Rates from Lattice QCD with Physical Quark Masses

Stefan Meinel

Department of Physics, University of Arizona, Tucson, Arizona 85721, USA and RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973, USA

(Received 1 December 2016; published 21 February 2017)


Input the measured BFs from BESIII

Triggered by BESIII

$$B(\Lambda_c \rightarrow \Lambda \ell^+ \nu_\ell) = \begin{cases} 0.0363(38)(20), & \ell = e, \\ 0.0349(46)(27), & \ell = \mu. \end{cases}$$

The first LQCD calculations on BFs and form factors

$$B(\Lambda_c \rightarrow \Lambda \ell^+ \nu_\ell) = \begin{cases} 0.0380(19)_{\text{LQCD}}(11)_{\tau_{\Lambda_c}}, & \ell = e, \\ 0.0369(19)_{\text{LQCD}}(11)_{\tau_{\Lambda_c}}, & \ell = \mu, \end{cases}$$



✓ The first determination of $|V_{cs}|$ based on BFs of $\Lambda_c^+ \rightarrow \Lambda \ell^+ \nu_\ell$ measured by BESIII


$$|V_{cs}| = \begin{cases} 0.951(24)_{\text{LQCD}}(14)_{\tau_{\Lambda_c}}(56)_B, & \ell = e, \\ 0.947(24)_{\text{LQCD}}(14)_{\tau_{\Lambda_c}}(72)_B, & \ell = \mu, \\ 0.949(24)_{\text{LQCD}}(14)_{\tau_{\Lambda_c}}(49)_B, & \ell = e, \mu, \end{cases}$$

✓ More data on Λ_c^+ will be collected at BESIII



Λ_c^+ DECAY MODES

Mode	Fraction (Γ_i/Γ)	S _c Confi
Hadronic modes with a p: S = -1 final states		
Γ_1 $p\bar{K}^0$	(3.21 ± 0.30) %	
Γ_2 $pK^-\pi^+$	(6.84 ± 0.32, 0.40) %	
Γ_3 $p\bar{K}^*(892)^0$	[a] (2.13 ± 0.30) %	
Γ_4 $\Delta(1232)^{++}K^-$	(1.18 ± 0.27) %	
Γ_5 $\Lambda(1520)\pi^+$	[a] (2.4 ± 0.6) %	
Γ_6 $pK^-\pi^+$ nonresonant	(3.8 ± 0.4) %	
Γ_7 $p\bar{K}^0\pi^0$	(4.5 ± 0.6) %	
Γ_8 $p\bar{K}^0\eta$	(1.7 ± 0.4) %	
Γ_9 $p\bar{K}^0\pi^+\pi^-$	(3.5 ± 0.4) %	
Γ_{10} $pK^-\pi^+\pi^0$	(4.6 ± 0.8) %	
Γ_{11} $pK^*(892)^-\pi^+$	[a] (1.5 ± 0.5) %	
Γ_{12} $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(5.0 ± 0.9) %	
Γ_{13} $\Delta(1232)K^*(892)$	seen	
Γ_{14} $pK^-\pi^+\pi^+\pi^-$	(1.5 ± 1.0) × 10 ⁻³	
Γ_{15} $pK^-\pi^+\pi^0\pi^0$	(1.1 ± 0.5) %	
Γ_{16} $pK^-\pi^+3\pi^0$		
Hadronic modes with a p: S = 0 final states		
Γ_{17} $p\pi^+\pi^-$	(4.7 ± 2.5) × 10 ⁻³	
Γ_{18} $p f_0(980)$	[a] (3.8 ± 2.5) × 10 ⁻³	
Γ_{19} $p\pi^+\pi^+\pi^-\pi^-$	(2.5 ± 1.6) × 10 ⁻³	
Γ_{20} pK^+K^-	(1.1 ± 0.4) × 10 ⁻³	
Γ_{21} $p\phi$	[a] (1.12 ± 0.23) × 10 ⁻³	
Γ_{22} $pK^+K^-\text{non-}\phi$	(4.8 ± 1.9) × 10 ⁻⁴	
Hadronic modes with a hyperon: S = -1 final states		
Γ_{23} $\Lambda\pi^+$	(1.46 ± 0.13) %	
Γ_{24} $\Lambda\pi^+\pi^0$	(5.0 ± 1.3) %	
Γ_{25} $\Lambda\rho^+$	< 6 %	CL=95%
Γ_{26} $\Lambda\pi^+\pi^+\pi^-$	(3.59 ± 0.28) %	
Γ_{27} $\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow$	(1.0 ± 0.5) %	
Γ_{28} $\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow$	(7.5 ± 1.4) × 10 ⁻³	
Γ_{29} $\Lambda\pi^+\rho^0$	(1.4 ± 0.6) %	
Γ_{30} $\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$	(5 ± 4) × 10 ⁻³	
Γ_{31} $\Lambda\pi^+\pi^+\pi^-\text{nonresonant}$	< 1.1 %	CL=90%
Γ_{32} $\Lambda\pi^+\pi^+\pi^-\pi^0\text{total}$	(2.5 ± 0.9) %	
Γ_{33} $\Lambda\pi^+\eta$	[a] (2.4 ± 0.5) %	
Γ_{34} $\Sigma(1385)^+\eta$	[a] (1.16 ± 0.35) %	
Γ_{35} $\Lambda\pi^+\omega$	[a] (1.6 ± 0.6) %	
Γ_{36} $\Lambda\pi^+\pi^+\pi^-\pi^0, \text{no } \eta \text{ or } \omega$	< 9 × 10 ⁻³	CL=90%
Γ_{37} $\Lambda K^+\bar{K}^0$	(6.4 ± 1.3) × 10 ⁻³	S=1.6
Γ_{38} $\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$	(1.8 ± 0.6) × 10 ⁻³	
Γ_{39} $\Sigma^0\pi^+$	(1.43 ± 0.14) %	
Γ_{40} $\Sigma^+\pi^0$	(1.37 ± 0.30) %	
Γ_{41} $\Sigma^+\eta$	(7.5 ± 2.5) × 10 ⁻³	
Γ_{42} $\Sigma^+\pi^+\pi^-$	(4.9 ± 0.5) %	
Γ_{43} $\Sigma^+\rho^0$	< 1.8 %	CL=95%
Γ_{44} $\Sigma^-\pi^+\pi^+$	(2.3 ± 0.4) %	
Γ_{45} $\Sigma^0\pi^+\pi^0$	(2.5 ± 0.9) %	
Semileptonic modes		
Γ_{64} $\Lambda\ell^+\nu_\ell$	[b] (2.8 ± 0.4) %	
Γ_{65} $\Lambda e^+\nu_e$	(2.9 ± 0.5) %	
Γ_{66} $\Lambda\mu^+\nu_\mu$	(2.7 ± 0.6) %	



Hadronic modes with a p or n: S = -1 final states

Γ_1 pK_S^0	(1.59 ± 0.08) %	↓44%	S=1.1
Γ_2 $pK^-\pi^+$	(6.28 ± 0.32) %		S=1.4
Γ_3 $p\bar{K}^*(892)^0$	[a] (1.96 ± 0.27) %		
Γ_4 $\Delta(1232)^{++}K^-$	(1.08 ± 0.25) %		
Γ_5 $\Lambda(1520)\pi^+$	[a] (2.2 ± 0.5) %		
Γ_6 $pK^-\pi^+$ nonresonant	(3.5 ± 0.4) %		
Γ_7 $pK_S^0\pi^0$	(1.97 ± 0.13) %	↓50%	S=1.1
Γ_8 $nK_S^0\pi^+$	(1.82 ± 0.25) %	First	
Γ_9 $p\bar{K}^0\eta$	(1.6 ± 0.4) %		
Γ_{10} $pK_S^0\pi^+\pi^-$	(1.60 ± 0.12) %	↓28%	S=1.1
Γ_{11} $pK^-\pi^+\pi^0$	(4.46 ± 0.30) %	↓61%	S=1.5
Γ_{12} $pK^*(892)^-\pi^+$	[a] (1.4 ± 0.5) %		
Γ_{13} $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(4.6 ± 0.8) %		
Γ_{14} $\Delta(1232)K^*(892)$	seen		
Γ_{15} $pK^-2\pi^+\pi^-$	(1.4 ± 0.9) × 10 ⁻³		
Γ_{16} $pK^-\pi^+2\pi^0$	(1.0 ± 0.5) %		

Hadronic modes with a p: S = 0 final states

Γ_{17} $p\pi^0$	< 2.7	× 10 ⁻⁴	CL=90%
Γ_{18} $p\eta$	(1.24 ± 0.30) × 10 ⁻³		First
Γ_{19} $p\omega(782)^0$	(9 ± 4) × 10 ⁻⁴		
Γ_{20} $p\pi^+\pi^-$	(4.61 ± 0.28) × 10 ⁻³		First
Γ_{21} $p f_0(980)$	[a] (3.5 ± 2.3) × 10 ⁻³		
Γ_{22} $p2\pi^+2\pi^-$	(2.3 ± 1.4) × 10 ⁻³		
Γ_{23} pK^+K^-	(1.06 ± 0.06) × 10 ⁻³		
Γ_{24} $p\phi$	[a] (1.06 ± 0.14) × 10 ⁻³	↓36%	
Γ_{25} $pK^+K^-\text{non-}\phi$	(5.3 ± 1.2) × 10 ⁻⁴		
Γ_{26} $p\phi\pi^0$	(10 ± 4) × 10 ⁻⁵		
Γ_{27} $pK^+K^-\pi^0\text{nonresonant}$	< 6.3	× 10 ⁻⁵	CL=90%

Hadronic modes with a hyperon: S = -1 final states

Γ_{28} $\Lambda\pi^+$	(1.30 ± 0.07) %		S=1.1
Γ_{29} $\Lambda\pi^+\pi^0$	(7.1 ± 0.4) %	↓78%	S=1.1
Γ_{30} $\Lambda\rho^+$	< 6 %		CL=95%
Γ_{31} $\Lambda\pi^-2\pi^+$	(3.64 ± 0.29) %		S=1.4
Γ_{44} $\Sigma^0\pi^+$	(1.29 ± 0.07) %	↓45%	S=1.1
Γ_{45} $\Sigma^+\pi^0$	(1.25 ± 0.10) %	↓33%	
Γ_{46} $\Sigma^+\eta$	(4.4 ± 2.0) × 10 ⁻³		
Γ_{47} $\Sigma^+\eta'$	(1.5 ± 0.6) %		
Γ_{48} $\Sigma^+\pi^+\pi^-$	(4.50 ± 0.25) %	↓46%	S=1.3
Γ_{49} $\Sigma^+\rho^0$	< 1.7 %		CL=95%
Γ_{50} $\Sigma^-2\pi^+$	(1.87 ± 0.18) %		
Γ_{51} $\Sigma^0\pi^+\pi^0$	(3.5 ± 0.4) %		
Γ_{52} $\Sigma^+\pi^0\pi^0$	(1.55 ± 0.15) %		
Γ_{53} $\Sigma^0\pi^-2\pi^+$	(1.11 ± 0.30) %		

Semileptonic modes

Γ_{72} $\Lambda e^+\nu_e$	(3.6 ± 0.4) %	
Γ_{73} $\Lambda\mu^+\nu_\mu$	(3.5 ± 0.5) %	↓35%



Experimental precision reaches of the charmed hadrons

	Golden hadronic mode	$\delta B/B$	Golden SL mode	$\delta B/B$
D^0	$B(K\pi)=(3.88\pm 0.05)\%$	1.3%	$B(K\ell\nu)=(3.55\pm 0.05)\%$	1.4%
D^+	$B(K\pi\pi)=(9.13\pm 0.19)\%$	2.1%	$B(K^0\ell\nu)=(8.83\pm 0.22)\%$	2.5%
D_s	$B(KK\pi)=(5.39\pm 0.21)\%$	3.9%	$B(\phi\ell\nu)=(2.49\pm 0.14)\%$	5.6%
Λ_c	$B(pK\pi)=(5.0\pm 1.3)\%$ (PDG2014) $= (6.8\pm 0.36)\%$ (BELLE) $= (5.84\pm 0.35)\%$ (BESIII) $= (6.46\pm 0.24)\%$ (HFLAV)	26% 5.3% 6.0% 3.7%	$B(\Lambda\ell\nu)=(2.1\pm 0.6)\%$ (PDG2014) $= (3.63\pm 0.43)\%$ (BESIII) $= (3.18\pm 0.32)\%$ (HFLAV)	29% 12% 10%



- The precisions of Λ_c decay rates reaches to the level of charmed mesons!
- However, search for more unknown modes, especially **Cabbibo-suppressed** mode, are important



Phase II

After success of the first Λ_c^+ data set, an energy upgrade on the BEPCII, $4.6 \text{ GeV} \rightarrow 4.95 \text{ GeV}$, has been implemented to study Λ_c^+ with more statistics

BESIII Physics Reports



Int. J. Mod. Phys. A 24, S1-794 (2009)
[arXiv:0809.1869 [hep-ex]].

Chin. Phys. C 44, 040001 (2020)
doi:10.1088/1674-1137/44/4/040001
[arXiv:1912.05983 [hep-ex]].



Planned future data set

Table 7.1: List of data samples collected by BESIII/BEPCII up to 2019, and the proposed samples for the remainder of the physics program. The most right column shows the number of required data taking days in current (T_C) or upgraded (T_U) machine. The machine upgrades include top-up implementation and beam current increase.


Energy	Physics motivations	Current data	Expected final data	T_C / T_U
1.8 - 2.0 GeV	R values Nucleon cross-sections	N/A	0.1 fb ⁻¹ (fine scan)	60/50 days
2.0 - 3.1 GeV	R values Cross-sections	Fine scan (20 energy points)	Complete scan (additional points)	250/180 days
✓ J/ψ peak	Light hadron & Glueball J/ψ decays	3.2 fb ⁻¹ (10 billion)	3.2 fb ⁻¹ (10 billion)	N/A
✓ $\psi(3686)$ peak	Light hadron & Glueball Charmonium decays	0.67 fb ⁻¹ (0.45 billion)	4.5 fb ⁻¹ (3.0 billion)	150/90 days
$\psi(3770)$ peak	D^0/D^\pm decays	2.9 fb ⁻¹	20.0 fb ⁻¹	610/360 days
3.8 - 4.6 GeV	R values XYZ /Open charm	Fine scan (105 energy points)	No requirement	N/A
4.180 GeV	D_s decay XYZ /Open charm	3.2 fb ⁻¹	6 fb ⁻¹	140/50 days
4.0 - 4.6 GeV	XYZ /Open charm Higher charmonia cross-sections	16.0 fb ⁻¹ at different \sqrt{s}	30 fb ⁻¹ at different \sqrt{s}	770/310 days
4.6 - 4.9 GeV	Charmed baryon/ XYZ cross-sections	0.56 fb ⁻¹ at 4.6 GeV	15 fb ⁻¹ at different \sqrt{s}	1490/600 days
4.74 GeV	$\Sigma_c^+ \Lambda_c^-$ cross-section	N/A	1.0 fb ⁻¹	100/40 days
4.91 GeV	$\Sigma_c \Sigma_c$ cross-section	N/A	1.0 fb ⁻¹	120/50 days
4.95 GeV	Ξ_c decays	N/A	1.0 fb ⁻¹	130/50 days

to be complete
in 2022-24

18 fb⁻¹
 Λ_c^+ data

in 2020-2021, 5.8 fb⁻¹ is taken

[Chin. Phys. C 46, 113003 (2022)]



in total, 6.4 fb^{-1} data above Λ_c^+ threshold ($\sim 8x$ times more Λ_c^+ statistics)

- First measurement of absolute form factors of $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$
- Observation of second SL decay $\Lambda_c^+ \rightarrow p K^- e^+ \nu_e$
- Many observations and improved precisions of Cabbibo-Suppressed modes
- First partial wave analysis of Λ_c^+ decays
- More studies of neutron-involved decay modes
- Search for rare decay $\Lambda_c^+ \rightarrow \gamma \Sigma^+$



Semi-leptonic decay

- ✓ Form factors of $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$
- ✓ $\Lambda_c^+ \rightarrow p K^- e^+ \nu_e$
- ✓ LFU test of $\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$,
- ✓ Search for $\Lambda_c^+ \rightarrow \Lambda \pi^+ \pi^- e^+ \nu_e$ and $p K_S \pi^- e^+ \nu_e$

PRL129, 231803 (2022)
PRD106, 112010 (2022)
PRD108, L031105 (2023)
PLB843, 137993 (2023)

Neutron-involved decay

- ✓ $\Lambda_c^+ \rightarrow n \pi^+$
- ✓ $\Lambda_c^+ \rightarrow n \pi^+ \pi^0, n \pi^+ \pi^+ \pi^-, n K^- \pi^+ \pi^+$
- ✓ $\Lambda_c^+ \rightarrow n K_S K^+$
- ✓ $\Lambda_c^+ \rightarrow \Sigma^- K^+ \pi^+$
- ✓ $\Lambda_c^+ \rightarrow n K_S \pi^+ \pi^0$

PRL 128, 142001 (2022)
CPC 47, 023001 (2023) (Cover Story)
arXiv:2311.17131
arXiv:2309.05484
arXiv:2401.06813

Hadronic CS decays

- ✓ $\Lambda_c^+ \rightarrow p \pi^0, p \eta, p \eta', p \omega$
- ✓ $\Lambda_c^+ \rightarrow \Lambda K^+, \Lambda K^+ \pi^0$
- ✓ $\Lambda_c^+ \rightarrow \Sigma^+ K_S, \Sigma^0 K^+$
- ✓ $\Lambda_c^+ \rightarrow \Sigma^+ K^+ \pi^-$

PRD106, 072002 (2022); JHEP11, 137 (2023); arXiv:2311.06883

PRD106, L11101 (2022); PRD109, 032003 (2024)

PRD106, 052003 (2022)
JHEP09, 125 (2023);

Hadronic CF decays

- ✓ PWA of $\Lambda_c^+ \rightarrow \Lambda \pi^+ \pi^0$
- ✓ W-exchange-only process $\Xi^0 K^+$
- ✓ $\Lambda_c^+ \rightarrow \Xi^0 K^+ \pi^0$

JHEP 12, 033(2022)
PRL132, 031801 (2024)
PRD109, 052001 (2024)

Inclusive decay

- ✓ Improved BF of $\Lambda_c^+ \rightarrow e^+ X$
- ✓ First BF of $\bar{\Lambda}_c^- \rightarrow \bar{n} X$

PRD107, 052005 (2023)
PRD108, L031101 (2023)

Rare decay

- ✓ $\Lambda_c^+ \rightarrow \gamma \Sigma^+$

PRD107, 052002 (2023)

Production and excited Λ_c^+


- ✓ $\Lambda_c^+ \bar{\Lambda}_c^-$ lineshape and form factor
- ✓ $\Lambda_c (2595)^+$ and $\Lambda_c (2625)^+$ production and decay

PRL107, 052002 (2023)
arXiv:2312.08414; arXiv:2401.09225


25 papers are published

Determination of form factors of

$$\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e \quad \text{PRL129, 231803 (2022)}$$




$$B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (3.56 \pm 0.11 \pm 0.07)\%$$



First direct comparisons on the differential decay rates and form factors with LQCD calculations

Observation of $\Lambda_c^+ \rightarrow pK^- e^+ \nu$

PRD106, 112010 (2022)




$$B(\Lambda_c^+ \rightarrow pK^- e^+ \nu) = (0.88 \pm 0.17 \pm 0.07) \times 10^{-3}$$

$$B(\Lambda_c^+ \rightarrow \Lambda(1405)e^+ \nu) = (1.69 \pm 0.76 \pm 0.16) \times 10^{-3}$$














$$B(\Lambda_c^+ \rightarrow \Lambda(1520)e^+ \nu) = (0.99 \pm 0.51 \pm 0.10) \times 10^{-3}$$

- Second leptonic decay of Λ_c^+ is observed!
- Good channel to study Λ excited states, such as $\Lambda(1405)$ and $\Lambda(1520)$

PLB 843, 137993 (2023)






PRD 108, L031105 (2023)



First evidence of $\Lambda_c^+ \rightarrow p\pi^0$

Most precise measurement of $\Lambda_c^+ \rightarrow p\eta$


arXiv:2311.06883



		$B(\Lambda_c^+ \rightarrow p\eta)$	$B(\Lambda_c^+ \rightarrow p\omega)$
BESIII		$1.24 \pm 0.28 \pm 0.10$ [22]	—
LHCb		—	$0.94 \pm 0.32 \pm 0.22$ [23]
Belle		$1.42 \pm 0.05 \pm 0.11$ [24]	$0.827 \pm 0.075 \pm 0.075$ [25]
This paper		$1.57 \pm 0.11 \pm 0.04$	$1.11 \pm 0.20 \pm 0.07$
Current algebra	Uppal [13]	0.3	—
	Cheng [26]	1.28	—
SU(3) flavor symmetry	Sharma [14]	$0.2^a(1.7^b)$	—
	Geng [27]	$1.25^{+0.38}_{-0.36}$	—
	Geng [28]	1.30 ± 0.10	—
	Hsiao [29]	1.24 ± 0.21	—
	Geng [30]	—	0.63 ± 0.34
	Hsiao [31]	—	1.14 ± 0.54
	Zhong [32]	$1.36^a(1.27^b)$	—
Topological diagram method	Hsiao [33]	1.42 ± 0.23^c (1.47 ± 0.28^d)	—
Heavy quark effective theory	Singer [34]	—	0.36 ± 0.02


Observation of $\Lambda_c^+ \rightarrow n\pi^+$

PRL128, 142001 (2022)




Determination of the BF for $\Lambda_c^+ \rightarrow \Sigma^+ K_S^0$ and $\Sigma^0 K^+$

PRD106, 052003 (2022)




$\Lambda_c^+ \rightarrow p\eta'$

PRD106, 072002 (2022)




$\Lambda_c^+ \rightarrow \Lambda K^+$

PRD106, 111101 (2022)




Many CS modes are explored.


$\Lambda_c^+ \rightarrow n\pi^+\pi^0$



$\Lambda_c^+ \rightarrow n\pi^+\pi^-\pi^+$



$\Lambda_c^+ \rightarrow nK^-\pi^+\pi^+$




CPC47, 023001 (2023)

Decay asymmetry in W-exchange-only process $\Lambda_c^+ \rightarrow \Xi^0 K^+$

- Previous theoretical calculation on the BF lower than exp. measurement, which all predicted zero decay asymmetry
- BESIII confirmed the exp. result of BF in 2018 [PLB 783, 200 (2018)]
- In theory, BF is enhanced by enhancing the decay asymmetry close to 1

$$\alpha_{\Xi^0 K^+} = 2\text{Re}(s^* p) / (|s|^2 + |p|^2),$$

$$\delta_p - \delta_s = \arctan(\sqrt{1 - \alpha_{\Xi^0 K^+}^2} \sin \Delta_{\Xi^0 K^+} / \alpha_{\Xi^0 K^+})$$







Theory or experiment	$\mathcal{B}(\Lambda_c^+ \rightarrow \Xi^0 K^+) (\times 10^{-3})$	$\alpha_{\Xi^0 K^+}$	$ A (\times 10^{-2} G_F \text{ GeV}^2)$	$ B (\times 10^{-2} G_F \text{ GeV}^2)$	$\delta_p - \delta_s \text{ (rad)}$
Körner (1992), CCQM [7]	2.6	0
Xu (1992), Pole [8]	1.0	0	0	7.94	...
Żencaykowski (1994), Pole [9]	3.6	0
Ivanov (1998), CCQM [10]	3.1	0
Sharma (1999), CA [11]	1.3	0
Geng (2019), SU(3) [12]	5.7 ± 0.9	$0.94^{+0.06}_{-0.11}$	2.7 ± 0.6	16.1 ± 2.6	...
Zou (2020), CA [6]	7.1	0.90	4.48	12.10	...
Zhong (2022), SU(3) ^a [13]	$3.8^{+0.4}_{-0.5}$	$0.91^{+0.03}_{-0.04}$	3.2 ± 0.2	$8.7^{+0.6}_{-0.8}$...
Zhong (2022), SU(3) ^b [13]	$5.0^{+0.6}_{-0.9}$	0.99 ± 0.01	$3.3^{+0.5}_{-0.7}$	$12.3^{+1.2}_{-1.8}$...
BESIII (2018) [14]	$5.90 \pm 0.86 \pm 0.39$
PDG fit (2022) [2]	5.5 ± 0.7

process $\Lambda_c^+ \rightarrow \Xi^0 K^+$

three-level cascade decay $\Lambda_c^+ \rightarrow \Xi^0 K^+, \Xi^0 \rightarrow \Lambda \pi^0, \Lambda \rightarrow p \pi^-$

PRL132, 031801(2024)




- First determination of decay asymmetry $\alpha_{\Xi^0 K^+} = 0.01 \pm 0.16 \pm 0.03$, consistent with zero
- No theoretical model explains the current results
- First determination on phase difference $\delta_p - \delta_s$, with two solutions of $\pi/2$ and $-\pi/2$



Λ_c^+ decay asymmetries

Predictions and measurements	$\alpha_{\Lambda_c^+}^{pK_s^0}$	$\alpha_{\Lambda_c^+}^{\Lambda\pi^+}$	$\alpha_{\Lambda_c^+}^{\Sigma^0\pi^+}$	$\alpha_{\Lambda_c^+}^{\Sigma^+\pi^0}$	$\alpha_{\Lambda_c^+}^{\Xi^0K^+}$
CLEO(1990) [1]	-	$-1.0^{+0.4}_{-0.1}$	-	-	-
ARGUS(1992) [2]	-	-0.96 ± 0.42	-	-	-
Körner(1992), CCQM [3]	-0.10	-0.70	0.70	0.71	0
Xu(1992), Pole [4]	0.51	-0.67	0.92	0.92	0
Cheng, Tseng(1992), Pole [5]	-0.49	-0.96	0.83	0.83	-
Cheng, Tseng(1993), Pole [6]	-0.49	-0.95	0.78	0.78	-
Żencaykowski(1994), Pole [7]	-0.90	-0.86	-0.76	-0.76	0
Żencaykowski(1994), Pole [8]	-0.66	-0.99	0.39	0.39	0
CLEO(1995) [9]	-	$-0.94^{+0.21+0.12}_{-0.06-0.06}$	-	$-0.45 \pm 0.31 \pm 0.06$	-
Alakabha Datta(1995), CA [10]	-0.91	-0.94	-0.47	-0.47	-
Ivanov(1998), CCQM [11]	-0.97	-0.95	0.43	0.43	0
Sharma(1999), CA [12]	-0.99	-0.99	-0.31	-0.31	0
FOCUS(2006) [13]	-	$-0.78 \pm 0.16 \pm 0.19$	-	-	-
BESIII(2018) [14]	$0.18 \pm 0.43 \pm 0.14$	$-0.80 \pm 0.11 \pm 0.02$	$-0.73 \pm 0.17 \pm 0.07$	$-0.57 \pm 0.10 \pm 0.07$	-
Geng(2019), SU(3) [15]	$-0.89^{+0.26}_{-0.11}$	-0.87 ± 0.10	-0.35 ± 0.27	-0.35 ± 0.27	$0.94^{+0.06}_{-0.11}$
Zou(2020), CA [16]	-0.75	-0.93	-0.76	-0.76	0.90
BELLE(2022) [17, 18]	-	$-0.755 \pm 0.005 \pm 0.003$	$-0.463 \pm 0.016 \pm 0.008$	$-0.48 \pm 0.02 \pm 0.02$	-
Zhong(2022), SU(3) ^a [19]	-0.57 ± 0.21	-0.75 ± 0.01	-0.47 ± 0.03	-0.47 ± 0.03	$0.91^{+0.03}_{-0.04}$
Zhong(2022), SU(3) ^b [19]	-0.29 ± 0.24	-0.75 ± 0.01	-0.47 ± 0.03	-0.47 ± 0.03	0.99 ± 0.01
Liu(2023), Pole [20]	-0.81 ± 0.05	-0.75 ± 0.01	-0.47 ± 0.01	-0.45 ± 0.04	0.95 ± 0.02
Liu(2023), LP [20]	-0.68 ± 0.01	-0.75 ± 0.01	-0.47 ± 0.01	-0.45 ± 0.04	0.92
BESIII(2023) [21]	-	-	-	-	0.01 ± 0.16
Geng(2023), SU(3) [22]	-0.40 ± 0.49	-0.75 ± 0.01	-0.47 ± 0.02	-0.47 ± 0.02	-0.15 ± 0.14
Zhong(2024), TDA [23]	0.01 ± 0.24	-0.76 ± 0.01	-0.48 ± 0.02	-0.48 ± 0.02	-0.16 ± 0.13
Zhong(2024), IRA [23]	0.03 ± 0.24	-0.76 ± 0.01	-0.48 ± 0.02	-0.48 ± 0.02	-0.19 ± 0.12
PDG(for now) [24]	0.20 ± 0.50 (only BESIII)	-0.84 ± 0.09	-0.73 ± 0.18 (only BESIII)	-0.55 ± 0.11	-



- First Amplitude analysis of charmed baryon multi-hadronic decays
- Based on **TF-PWA** package: <https://gitlab.com/jiangyi15/tf-pwa>



	Theoretical calculation		This work	PDG
$10^2 \times \mathcal{B}(\Lambda_c^+ \rightarrow \Lambda \rho(770)^+)$	4.81 ± 0.58 [13]	4.0 [14, 15]	4.06 ± 0.52	< 6
$10^3 \times \mathcal{B}(\Lambda_c^+ \rightarrow \Sigma(1385)^+ \pi^0)$	2.8 ± 0.4 [16]	2.2 ± 0.4 [17]	5.86 ± 0.80	—
$10^3 \times \mathcal{B}(\Lambda_c^+ \rightarrow \Sigma(1385)^0 \pi^+)$	2.8 ± 0.4 [16]	2.2 ± 0.4 [17]	6.47 ± 0.96	—
$\alpha_{\Lambda \rho(770)^+}$	-0.27 ± 0.04 [13]	-0.32 [14, 15]	-0.763 ± 0.066	—
$\alpha_{\Sigma(1385)^+ \pi^0}$	$-0.91^{+0.45}_{-0.10}$ [17]		-0.917 ± 0.083	—
$\alpha_{\Sigma(1385)^0 \pi^+}$	$-0.91^{+0.45}_{-0.10}$ [17]		-0.79 ± 0.11	—

Many first measurements of intermediate states!

PRL131, 191901 (2023)







- Negate the $Y(4630)$ in decaying into $\Lambda_c^+\Lambda_c^-$ reported by BELLE
- Energy-dependence of $|G_E/G_M|$ reveals an oscillation feature, which may imply a non-trivial structure of the lightest charmed baryon.

Observation of $e^+e^- \rightarrow \Lambda_c^+\bar{\Lambda}_c(2595)^-$ and $\Lambda_c^+\bar{\Lambda}_c(2625)^-$



arXiv:2312.08414

Datasets of 208/pb at 4.92 GeV and 159/pb at 4.95 GeV



$$\frac{d\sigma}{d\cos\theta} \propto (1 + \cos^2\theta)(|G_E|^2 + 3|G_M|^2) + \frac{1}{\tau}|G_C|^2 \sin^2\theta$$




Phase III

To better accomplish the physics potentials in the white paper, an upgrade plan of BEPCII is being implemented

An upgrade of BEPCII (**BEPCII-U**) has been approved in July 2021 and planned to be completed by the end of 2024

- ✓ **Improve luminosity by 3 times higher than current BEPCII at 4.7 GeV**
- ✓ **Extend the maximum energy to 5.6 GeV**



Capable of finishing the proposed luminosity of Λ_c^+ data in shorter time



1490 → 600 days



Energy	Physics motivations	Current data	Expected final data	T_C / T_U
4.6 - 4.9 GeV	Charmed baryon/XYZ cross-sections	0.56 fb ⁻¹ at 4.6 GeV	15 fb ⁻¹ at different \sqrt{s}	1490/600 days
4.74 GeV	$\Sigma_c^+ \Lambda_c^-$ cross-section	N/A	1.0 fb ⁻¹	100/40 days
4.91 GeV	$\Sigma_c \Sigma_c$ cross-section	N/A	1.0 fb ⁻¹	120/50 days
4.95 GeV	Ξ_c decays	N/A	1.0 fb ⁻¹	130/50 days

Heavier charmed baryons

R



- Energy thresholds

- ✓ $e^+e^- \rightarrow \Lambda_c^+ \bar{\Sigma}_c^-$ 4.74 GeV
- ✓ $e^+e^- \rightarrow \Lambda_c^+ \bar{\Sigma}_c \pi$ 4.88 GeV
- ✓ $e^+e^- \rightarrow \Sigma_c \bar{\Sigma}_c$ 4.91 GeV
- ✓ $e^+e^- \rightarrow \Xi_c \bar{\Xi}_c$ 4.94 GeV
- ✓ $e^+e^- \rightarrow \Omega_c^0 \bar{\Omega}_c^0$ 5.40 GeV

- Cover all the **ground-state charmed baryons**: studies on their production & decays, CPV search, **to help developing more reliable QCD-derived models in charm sector**
- Studies on the production and decays of **excited charmed baryons**



Yet-to-be-Explored $\Xi_c^{+,0} / \Omega_c^0$ Decays

- We (will) have precise Λ_c^+ data after BESIII efforts
- However, $\Xi_c^{+,0} / \Omega_c^0$ has insufficient data
- A new territory for BESIII!



Mode	Fraction (Γ_i / Γ)
▼ Cabibbo-favored ($S = -2$) decays	
Γ_1 $p2 K_S^0$	$(2.5 \pm 1.3) \times 10^{-3}$
Γ_2 $\Lambda \bar{K}^0 \pi^+$	
Γ_3 $\Sigma(1385)^+ \bar{K}^0$	⁽¹⁾ $(2.9 \pm 2.0)\%$
Γ_4 $\Lambda K^- 2 \pi^+$	$(9 \pm 4) \times 10^{-3}$
Γ_5 $\Lambda \bar{K}^*(892)^0 \pi^+$	⁽¹⁾ $< 5 \times 10^{-3}$
Γ_6 $\Sigma(1385)^+ K^- \pi^+$	⁽¹⁾ $< 6 \times 10^{-3}$
Γ_7 $\Sigma^+ K^- \pi^+$	$(2.7 \pm 1.2)\%$
Γ_8 $\Sigma^+ \bar{K}^*(892)^0$	⁽¹⁾ $(2.3 \pm 1.1)\%$
Γ_9 $\Sigma^0 K^- 2 \pi^+$	$(8 \pm 5) \times 10^{-3}$
Γ_{10} $\Xi^0 \pi^+$	$(1.6 \pm 0.8)\%$
Γ_{11} $\Xi^- 2 \pi^+$	$(2.9 \pm 1.3)\%$
Γ_{12} $\Xi(1530)^0 \pi^+$	⁽¹⁾ $< 2.9 \times 10^{-3}$
Γ_{13} $\Xi(1620)^0 \pi^+$	seen
Γ_{14} $\Xi(1690)^0 \pi^+$	seen
Γ_{15} $\Xi^0 \pi^+ \pi^0$	$(6.7 \pm 3.5)\%$
Γ_{16} $\Xi^0 \pi^- 2 \pi^+$	$(5.0 \pm 2.6)\%$
Γ_{17} $\Xi^0 e^+ \nu_e$	$(7 \pm 4)\%$
Γ_{18} $\Omega^- K^+ \pi^+$	$(2.0 \pm 1.5) \times 10^{-3}$

Ξ_c^+ PDG2023

Mode	Fraction (Γ_i / Γ)
► Cabibbo-favored ($S = -3$) decays – relative to $\Omega^- \pi^+$	
Γ_6 $\Xi^0 \bar{K}^0$	1.64 ± 0.29
Γ_7 $\Xi^0 K^- \pi^+$	1.20 ± 0.18
Γ_8 $\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	0.68 ± 0.16
Γ_9 $\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow \Xi^0 K^-$	0.12 ± 0.05
Γ_{10} $\Xi^- \bar{K}^0 \pi^+$	2.12 ± 0.28
Γ_{11} $\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow \Xi^- \bar{K}^0$	0.12 ± 0.06
Γ_{12} $\Xi^- K^- 2 \pi^+$	0.63 ± 0.09
Γ_{13} $\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow \Xi^- \pi^+$	0.21 ± 0.06
Γ_{14} $\Xi^- \bar{K}^{*0} \pi^+$	0.34 ± 0.11
Γ_{15} $p K^- K^- \pi^+$	seen
Γ_{16} $\Sigma^+ K^- K^- \pi^+$	< 0.32
Γ_{17} $\Lambda \bar{K}^0 \bar{K}^0$	1.72 ± 0.35



Ω_c^0 PDG2023

LHCb



Potentials @LHCb

- Huge Λ_c production at LHCb: $\sim 100\mu\text{b}$
- **Prompt charm**: using exclusive reconstruction
- **Secondary charm** from b -hadron decays with inclusive b triggers
- $\Lambda_c^+ \rightarrow pK^-\pi^+$ CF yields: 0.8M in 0.65/fb ($\sim 20\%$ of Run I data)
- CS samples $O(10^5)$ in Run I:
BF measurement and CPV
- DCS $\Lambda_c^+ \rightarrow pK^+\pi^-$ can be measured with best precision
- Potential to set up the SL modes $pK^-\mu^+\nu$ and $p\pi^-\mu^+\nu$
 → size of this BF is critical to understand the internal dynamics of Λ_c
- Search for CS SL mode: $p\pi^-\mu^+\nu$
- amplitude analysis of $\Lambda_c \rightarrow \Lambda\mu^+\nu$, to extract form factors
 → input to theoretical calculation
- Rare decays: $p\mu^+\mu^-$, 3μ , $p\mu^+\mu^+$, ...




tag in secondary Λ_b/Σ_c decays

- singly charmed baryon
 - ✓ Ξ_c^+ mass [Phys. Rev. Lett. 113, 032001 (2014)]
 - ✓ BF for $\Lambda_c^+ \rightarrow pK^+\pi^-, pK^+K^-, p\pi^+\pi^-$ [JHEP 03, 043 (2018)]
 - ✓ CPV search in $\Lambda_c^+ \rightarrow pK^+K^-, p\pi^+\pi^-$ [JHEP 03, 182 (2018)]
 - ✓ Rare decay of $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ [Phys. Rev. D 97, 091101 (2018)]
 - ✓ Lifetime of Ω_c^0 [Phys. Rev. Lett. 121, 092003 (2018)]
 - ✓ Lifetimes of Λ_c^+, Ξ_c^+ and Ξ_c^0 [Phys. Rev. D 100, 032001 (2019)]
 - ✓ CPV search in $\Xi_c^+ \rightarrow pK^-\pi^+$ [Eur. Phys. J. C 80, 986 (2020)]
 - ✓ Suppressed decay $\Xi_c^0 \rightarrow \Lambda_c^+\pi^-$ [Phys. Rev. D 102, 071101 (2020)]
 - ✓ Lifetime of Ω_c^0 and Ξ_c^0 [Sci. Bull. 67, 479 (2022)]
 - ✓ $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude analysis [Phys. Rev. D 108, 012023 (2023)]
 - ✓ Λ_c^+ polarimetry [JHEP 07, 228 (2023)]
 - ✓ $\Omega_c^0 \rightarrow \Omega^-K^+, \Xi^-\pi^+$ and Ω_c^0 mass [Phys. Rev. Lett. 132.081802 (2024)]
- doubly charmed baryon: Ξ_{cc}^{++} discovery and other researches since 2017
- charmed baryon spectroscopy
 - ✓ Λ_c^{*++} in $\Xi_c^+K^-$ via $\Lambda_b^0 \rightarrow D^0p\pi^-$ [JHEP 05, 030 (2017)]
 - ✓ Ω_c^{*+0} in $\Xi_c^+K^-$ via prompt production [Phys. Rev. Lett. 118,182001 (2017)]
 - ✓ Ξ_c^{*+0} in $\Lambda_c^+K^-$ final states [Phys. Rev. Lett. 124, 222001 (2020)]
 - ✓ Ω_c^{*+0} in $\Xi_c^+K^-$ from $\Omega_b^- \rightarrow \Xi_c^+K^-\pi^-$ [Phys. Rev. D 104, L091102 (2021)]
 - ✓ Ω_c^{*+0} in $\Xi_c^+K^-$ via prompt production [Phys. Rev. Lett. 131, 131902 (2023)]
- charmed baryon production in pp and pPb collisions
 - Λ_c^+ in pp [NPB 871 (2013), JHEP06, 147 (2017)]
 - Λ_c^+ in pPb [JHEP 02, 102 (2019), JHEP06(2023)132]
 - Ξ_c^+ in pPb [Phys. Rev. C 109, 044901 (2024)]

Lifetimes of charmed baryons

<i>arXiv:2111.09566</i>	unit: fs	$\tau(\Xi_c^+)$	$\tau(\Lambda_c^+)$	$\tau(\Xi_c^0)$	$\tau(\Omega_c^0)$
PDG (2004-2018) [10]		442 ± 26	200 ± 6	112^{+13}_{-10}	69 ± 12
LHCb (2018) [12]					268 ± 26
LHCb (2019) [14]		457 ± 6	203.5 ± 2.2	154.5 ± 2.6	
PDG (2020) [11]		456 ± 5	202.4 ± 3.1	153 ± 6	268 ± 26






- Lifetime of Ω_c^0 [Phys. Rev. Lett. 121, 092003 (2018)]
- Lifetimes of Λ_c^+ , Ξ_c^+ and Ξ_c^0 [Phys. Rev. D 100, 032001 (2019)]



$$\tau(\Xi_c^+) > \tau(\Lambda_c^+) > \tau(\Xi_c^0) > \tau(\Omega_c^0)$$



$$\tau(\Xi_c^+) > \tau(\Omega_c^0) > \tau(\Lambda_c^+) > \tau(\Xi_c^0).$$





Lifetime of Ω_c^0 and Ξ_c^0 [Sci. Bull. 67, 479 (2022)]






[arXiv:2305.00665](https://arxiv.org/abs/2305.00665)

unit: fs	$\tau(\Xi_c^+)$	$\tau(\Lambda_c^+)$	$\tau(\Xi_c^0)$	$\tau(\Omega_c^0)$
PDG (2004-2018) [3]	442 ± 26	200 ± 6	112^{+13}_{-10}	69 ± 12
LHCb (2018) [4]				268 ± 26
LHCb (2019) [6]	457 ± 6	203.5 ± 2.2	154.5 ± 2.6	
PDG (2020) [8]	456 ± 5	202.4 ± 3.1	153 ± 6	268 ± 26
LHCb (2021) [7]			148.0 ± 3.2	276.5 ± 14.1
PDG (2022) [1]	453 ± 5	201.5 ± 2.7	151.9 ± 2.4	268 ± 26
Belle II (2022) [9, 10]		203.20 ± 1.18		243 ± 49
WA values (2023)	453 ± 5	202.9 ± 1.1	150.5 ± 1.9	273 ± 12





BF ratios	This work	CA model	LFQM	Naive estimation
$\frac{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- K^+)}{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)}$	0.0608 ± 0.0064	-	-	0.0467
$\frac{\mathcal{B}(\Omega_c^0 \rightarrow \Xi^- \pi^+)}{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)}$	0.1581 ± 0.0099	0.1038	0.0345	-





unit: MeV	$m_{\Lambda_c^+}$	$m_{\Xi_c^+}$	$m_{\Xi_c^0}$	$m_{\Omega_c^0}$
之前国际平均值	2286.46 ± 0.14	$2467.93^{+0.28}_{-0.40}$	2470.44 ± 0.28	2695.2 ± 1.7
LHCb 2014		$2467.97 \pm 0.14 \pm 0.17$		
LHCb 2023				$2695.28 \pm 0.07 \pm 0.40$
当前国际平均值	2286.46 ± 0.14	2467.95 ± 0.19	2470.44 ± 0.28	2695.28 ± 0.40


$\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude analysis

Phys. Rev. D 108, 012023 (2023)

Λ_c^+ signals are selected via Λ_b^0
 $\rightarrow \Lambda_c^+ \mu^- \nu$ from dataset taken in 2016,
 where only a subset of 0.4 M signals
 are employed



5-dim fit



Resonance	Fit fraction (%)	Model $\sqrt{3}S$	α
$\Lambda(1405)$	7.7	$K^*(892) \sqrt{3}S$	0.662
$\Lambda(1520)$	1.86	$\Lambda(1405)$	0.873
$\Lambda(1600)$	5.2	$\Lambda(1520)$	-0.58
$\Lambda(1670)$	1.18	$\Lambda(1600)$	-0.925
$\Lambda(1690)$	1.19	$\Lambda(1670)$	-0.20
$\Lambda(2000)$	9.58	$\Lambda(1690)$	-0.817
$\Delta(1232)^{++}$	28.60	$\Lambda(2000)^{++}$	-0.958
$\Delta(1600)^{++}$	4.5	$\Delta(1232)^{++}$	-0.57
$\Delta(1700)^{++}$	3.90	$\Delta(1600)^{++}$	-0.548
$K_0^*(700)$	3.02	$\Delta(1700)^{++}$	-0.50
$K^*(892)$	22.14	$K_0^*(700)$	-0.216
$K_0^*(1430)$	14.7	$K^*(892)$	-0.06
		$K_0^*(1430)$	-0.34

Λ_c^+ polarization and $\Lambda_c^+ \rightarrow pK^-\pi^+$ polarimetry

Phys. Rev. D 108, 012023 (2023)


JHEP 07, 228 (2023)

Component	Value (%)
$P_x (lab)$	$60.32 \pm 0.68 \pm 0.98 \pm 0.21$
$P_y (lab)$	$-0.41 \pm 0.61 \pm 0.16 \pm 0.07$
$P_z (lab)$	$-24.7 \pm 0.6 \pm 0.3 \pm 1.1$
$P_x (\tilde{B})$	$21.65 \pm 0.68 \pm 0.36 \pm 0.15$
$P_y (\tilde{B})$	$1.08 \pm 0.61 \pm 0.09 \pm 0.08$
$P_z (\tilde{B})$	$-66.5 \pm 0.6 \pm 1.1 \pm 0.1$

A large Λ_c^+ polarization is found in b semi-leptonic decays $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- \nu$

- The obtained representation can facilitate polarization measurements of the Λ_c^+ baryon and eases inclusion of the $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay mode in hadronic amplitude analyses.
- At BESIII, the transverse polarization of Λ_c^+ can be obtained via $\Lambda_c^+ \rightarrow pK^-\pi^+$ polarimetry


The amplitude model is used to produce the distribution of the kinematic-dependent polarimeter vector in the space of Mandelstam variables to express the polarized decay rate in a model-independent way.



Future opportunity at LHCb



- RUN1&2: 9 fb^{-1}
- RUN3&4: 50 fb^{-1}
- ➔ x10 more statistics



- Further improvement on mass and lifetime measurement
- SCS and DCS hadronic decays
 - e.g. $\Xi_c^0 \rightarrow pK^-$, $\Xi_c^+ \rightarrow pK_S$, $\Omega_c^0 \rightarrow \Lambda K_S$, pK^-
- Semi-leptonic decays via b-baryon four-body decays
 - e.g. $\Lambda_c^+ \rightarrow pK^- \mu^+ \nu$, $p\pi^- \mu^+ \nu$; $\Xi_c^0 \rightarrow \Xi^- \mu^+ \nu$; $\Xi_c^+ \rightarrow \Lambda \mu^+ \nu$; $\Omega_c^0 \rightarrow \Omega^- \mu^+ \nu$
- Decay asymmetries and CPV search via prompt production or b-baryon decays
 - e.g. $\Lambda_c^+ \rightarrow pK_S$, $\Lambda\pi^+$, ΛK^+ ; $\Xi_c^0 \rightarrow \Lambda K_S$, $\Xi^- \pi^+$, $\Xi^- K^+$; $\Omega_c^0 \rightarrow \Omega^- \pi^+$, $\Omega^- K^+$, $\Xi^- \pi^+$
- Amplitude analysis of multi-body hadronic decays



Summary




- ◆ In recent years, experimental activities on charmed baryons are reviving, esp. at BESIII, LHCb and Belle (II)
- ◆ **Threshold data at BESIII** opens a new door to direct measurements of the decays → comprehensive and systematic studies of charmed baryon decays
 - BESIII has published several world-leading results based on ~ 80 M Λ_c samples
 - More efforts on hadronic decays w/ $n/\Sigma/\Xi$ particles & semi-leptonic decays
 - Plan to take data up to 5.6 GeV to cover all the ground-state charmed baryons
- ◆ **LHCb** has largest charmed baryon yields → large potential of best precisions
 - Search for more semi-leptonic decays
 - Precise determination of decay rate and amplitude analysis
 - Decay asymmetry and CPV test
 - Rare decays: LFV, FCNC, ...
- ◆ BESIII and LHCb will be complementary in charmed baryon decays



Backup

Discovery of the charmed heavy baryon

- Not exclusively clear about the first observation
- A number of experiments which published evidence for the charmed baryons beginning in 1975
 - ✓ First hint of charmed baryon $\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+$ in BNL [PRL34, 1125 \(1975\)](#)
 - ✓ First evidence of Λ_c^+ at Fermi Lab [PRL37, 882 \(1976\)](#)
- The first well established state is the Λ_c^+ at MarkII [PRL44, 10 \(1980\)](#)





CERN COURIER

brightrecruits.com/job/8901/senior-research-positions-ihep-cas-china

Latest Issue | [Archive](#) | [Jobs](#) | [Links](#) | [Buyer's guide](#) | [White papers](#) | [Events](#) | [Contact us](#)

Search

REGISTER NOW

Register as a member of cerncourier.com and get full access to all features of the site. Registration is free.

LATEST CERN COURIER ARTICLES

- ▶ Sneeze dynamics
- ▶ The longest proof
- ▶ Electron-hole collider
- ▶ Imaging with muons
- ▶ Towards a nuclear clock

SHARE THIS

- E-mail to a friend
- StumbleUpon
- Twitter
- Facebook
- CiteUlike
- SHARE ...

CERN COURIER


Mar 18, 2016

BESIII makes first direct measurement of the Λ_c at threshold

The charmed baryon, Λ_c , was first observed at Fermilab in 1976. Now, 40 years later, the Beijing Spectrometer (BESIII) experiment at the Beijing Electron-Positron Collider II (BEPCII) has measured the

absolute branching fraction of $\Lambda_c^+ \rightarrow pK^+\pi^+$ at threshold for the first time.

Because the decays of the Λ_c^+ to hadrons proceed only through the weak interaction, their branching fractions are key probes for understanding weak interactions inside of a baryon. In particular, precise measurements of the decays of the Λ_c^+ will provide important information on the final-state strong interaction in the charm sector, thereby improving the understanding of quantum chromodynamics in the non-perturbative energy region. In addition, because most of the



Beam-constrained mass distribution

DIGITAL EDITION

CERN Courier is now available as a regular digital edition. Click [here](#) to read the digital edition.

KEY SUPPLIERS



[More companies](#) ▶


FEATURED COMPANIES



HFLAV Fit to world BF data

- A fitter to constrain the 12 hadronic BFs and 1 SL BF, based on all the existing experimental data
- Correlated systematics are fully taken into account

Mode	HFAG 2016 (%)	BESIII (%)	PDG 2014 (%)	BELLE (%)
pK_S^0	1.59 ± 0.07	$1.52 \pm 0.08 \pm 0.03$	1.15 ± 0.30	
$pK^- \pi^+$	6.46 ± 0.24	$5.84 \pm 0.27 \pm 0.23$	5.0 ± 1.3	$6.84 \pm 0.24^{+0.21}_{-0.27}$
$pK_S^0 \pi^0$	2.03 ± 0.12	$1.87 \pm 0.13 \pm 0.05$	1.65 ± 0.50	
$pK_S^0 \pi^+ \pi^-$	1.69 ± 0.11	$1.53 \pm 0.11 \pm 0.09$	1.30 ± 0.35	
$pK^- \pi^+ \pi^0$	5.05 ± 0.29	$4.53 \pm 0.23 \pm 0.30$	3.4 ± 1.0	
$\Lambda \pi^+$	1.28 ± 0.06	$1.24 \pm 0.07 \pm 0.03$	1.07 ± 0.28	
$\Lambda \pi^+ \pi^0$	7.09 ± 0.36	$7.01 \pm 0.37 \pm 0.19$	3.6 ± 1.3	
$\Lambda \pi^+ \pi^- \pi^+$	3.73 ± 0.21	$3.81 \pm 0.24 \pm 0.18$	2.6 ± 0.7	
$\Sigma^0 \pi^+$	1.31 ± 0.07	$1.27 \pm 0.08 \pm 0.03$	1.05 ± 0.28	
$\Sigma^+ \pi^0$	1.25 ± 0.09	$1.18 \pm 0.10 \pm 0.03$	1.00 ± 0.34	
$\Sigma^+ \pi^+ \pi^-$	4.64 ± 0.24	$4.25 \pm 0.24 \pm 0.20$	3.6 ± 1.0	
$\Sigma^+ \omega$	1.77 ± 0.21	$1.56 \pm 0.20 \pm 0.07$	2.7 ± 1.0	
$\Lambda e^+ \nu_e$	3.18 ± 0.32	$3.63 \pm 0.38 \pm 0.20$	2.1 ± 0.6	



The least overall $\chi^2/\text{ndf}=30.0/23=1.3$

Precise $B(pK^- \pi^+)$ is useful for constrain V_{ub} determined via baryonic mode

A theoretical Framework for Charmed Hadrons

- Topological diagrams + Symmetries + Experimental inputs \Rightarrow to understand the decaying dynamics, predicting double-charm baryon decays, CPV, etc.

(predictive power)

- Λ_c^+ branching fractions used for global analysis

$\Rightarrow \Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ and $\Xi_c^+ \pi^+$ are large enough for observation.



$$Br(\Lambda_c^+ \rightarrow p\phi)/|V_{us}|^2 = 2\% \quad \rightarrow \quad Br(\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \bar{K}^{*0}) = O(\%)$$

[PRL 117, 232002 (2016)]

$$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+ K^- \pi^+$$

Large enough for observation

Important inputs to the Ξ_{cc}^{++} observation at LHCb

Λ_c^+ data from BESIII \rightarrow Stronger predictive power



Contribution to PDG

PDG2014

PDG2019

$\Gamma(\rho\bar{K}^0\pi^0)/\Gamma(\rho K^-\pi^+)$					Γ_7/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.66 \pm 0.05 \pm 0.07$	774	ALAM	98	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\rho\bar{K}^0\eta)/\Gamma(\rho K^-\pi^+)$					Γ_8/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.25 \pm 0.04 \pm 0.04$	57	AMMAR	95	CLE2	$e^+e^- \approx \Upsilon(4S)$

Unseen decay modes of the η are included.

$\Gamma(\rho\bar{K}^0\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$					Γ_9/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.51 ± 0.06 OUR AVERAGE					
$0.52 \pm 0.04 \pm 0.05$	985	ALAM	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
$0.43 \pm 0.12 \pm 0.04$	83	AVERY	91	CLEO	e^+e^- 10.5 GeV
$0.98 \pm 0.36 \pm 0.08$	12	BARLAG	90D	NA32	π^- 230 GeV

$\Gamma(\rho K^-\pi^+\pi^0)/\Gamma(\rho K^-\pi^+)$					Γ_{10}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.67 \pm 0.04 \pm 0.11$	2606	ALAM	98	CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\rho K^*(892)^-\pi^+)/\Gamma(\rho\bar{K}^0\pi^+\pi^-)$					Γ_{11}/Γ_9
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.44 ± 0.14	17	ALEEV	94	BIS2	nN 20-70 GeV

Unseen decay modes of the $K^*(892)^-$ are included.

$\Gamma(\rho(K^-\pi^+)_{\text{nonresonant}}\pi^0)/\Gamma(\rho K^-\pi^+)$					Γ_{12}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.73 \pm 0.12 \pm 0.05$	67	BOZEK	93	NA32	π^- Cu 230 GeV

$\Gamma(\rho K_S^0\pi^0)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.96 ± 0.13 OUR FIT	Error includes scale factor of 1.1				
$1.87 \pm 0.13 \pm 0.05$	558	ABLIKIM	16	BES3	$e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV

$\Gamma(\rho K_S^0\pi^0)/\Gamma(\rho K^-\pi^+)$					Γ_7/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.314 ± 0.018 OUR FIT					
$0.33 \pm 0.03 \pm 0.04$	774	ALAM	98	CLE2	$e^+e^- \approx \Upsilon(4S)$

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

$\Gamma(nK_S^0\pi^+)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.82 \pm 0.23 \pm 0.11$	83	ABLIKIM	17H	BES3	e^+e^- at 4.6 GeV

$\Gamma(\rho\bar{K}^0\eta)/\Gamma(\rho K^-\pi^+)$					Γ_9/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.25 \pm 0.04 \pm 0.04$	57	AMMAR	95	CLE2	$e^+e^- \approx \Upsilon(4S)$

Unseen decay modes of the η are included.

$\Gamma(\rho K_S^0\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.59 ± 0.12 OUR FIT	Error includes scale factor of 1.2				
$1.53 \pm 0.11 \pm 0.09$	485	ABLIKIM	16	BES3	$e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV

$\Gamma(\rho K_S^0\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$					Γ_{10}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.255 ± 0.015 OUR FIT	Error includes scale factor of 1.1.				
0.257 ± 0.031 OUR AVERAGE					
$0.26 \pm 0.02 \pm 0.03$	985	ALAM	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
$0.22 \pm 0.06 \pm 0.02$	83	AVERY	91	CLEO	e^+e^- 10.5 GeV
$0.49 \pm 0.18 \pm 0.04$	12	BARLAG	90D	NA32	π^- 230 GeV

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.


$\Gamma(\rho K^-\pi^+\pi^0)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.42 ± 0.31 OUR FIT	Error includes scale factor of 1.5				
$4.53 \pm 0.23 \pm 0.30$	1849	ABLIKIM	16	BES3	$e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV

$\Gamma(\rho K^-\pi^+\pi^0)/\Gamma(\rho K^-\pi^+)$					Γ_{11}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

- ◆ BESIII的贡献使得多数分支比结果由相对测量改为绝对测量。
- ◆ BESIII对黄金道 $\Lambda_c^+ \rightarrow pK\pi$ 的测量 \Rightarrow “模型依赖”变为“模型无关”。
- ◆ BESIII贡献发现了更多以前没发现的衰变道(比如含中子末态的衰变)。

- Energy thresholds

- ✓ $e^+e^- \rightarrow \Lambda_c^+ \bar{\Sigma}_c^-$ 4.74 GeV
- ✓ $e^+e^- \rightarrow \Lambda_c^+ \bar{\Sigma}_c^- \pi$ 4.88 GeV
- ✓ $e^+e^- \rightarrow \Sigma_c \bar{\Sigma}_c$ 4.91 GeV
- ✓ $e^+e^- \rightarrow \Xi_c \bar{\Xi}_c$ 4.94 GeV
- ✓ $e^+e^- \rightarrow \Omega_c^0 \bar{\Omega}_c^0$ 5.40 GeV





Charmed Baryons

	Structure	J^P	Mass, MeV	Width, MeV	Decay
Λ_c^+	udc	$(1/2)^+$	2286.46 ± 0.14	(200 ± 6) fs	weak
Ξ_c^+	usc	$(1/2)^+$	$2467.8_{-0.6}^{+0.4}$	(442 ± 26) fs	weak
Ξ_c^0	dsc	$(1/2)^+$	$2470.88_{-0.8}^{+0.34}$	112_{-10}^{+13} fs	weak
Σ_c^{++}	uuc	$(1/2)^+$	2454.02 ± 0.18	2.23 ± 0.30	$\Lambda_c^+ \pi^+$
Σ_c^+	udc	$(1/2)^+$	2452.9 ± 0.4	< 4.6	$\Lambda_c^+ \pi^0$
Σ_c^0	ddc	$(1/2)^+$	2453.76 ± 0.18	2.2 ± 0.4	$\Lambda_c^+ \pi^-$
$\Xi_c'^+$	usc	$(1/2)^+$	2575.6 ± 3.1	—	$\Xi_c^+ \gamma$
$\Xi_c'^0$	dsc	$(1/2)^+$	2577.9 ± 2.9	—	$\Xi_c^0 \gamma$
Ω_c^0	ssc	$(1/2)^+$	2695.2 ± 1.7	(69 ± 12) fs	weak
Σ_c^{*++}	uuc	$(3/2)^+$	2518.4 ± 0.6	14.9 ± 1.9	$\Lambda_c^+ \pi^+$
Σ_c^{*+}	udc	$(3/2)^+$	2517.5 ± 2.3	< 17	$\Lambda_c^+ \pi^0$
Σ_c^{*0}	ddc	$(3/2)^+$	2518.0 ± 0.5	16.1 ± 2.1	$\Lambda_c^+ \pi^-$
Ξ_c^{*+}	usc	$(3/2)^+$	$2645.9_{-0.6}^{+0.5}$	< 3.1	$\Xi_c \pi$
Ξ_c^{*0}	dsc	$(3/2)^+$	2645.9 ± 0.5	< 5.5	$\Xi_c \pi$
Ω_c^{*0}	ssc	$(3/2)^+$	2765.9 ± 2.0	—	$\Omega_c^0 \gamma$

Production near threshold and tag technique

$$E_{\text{cms}}=4600\text{MeV}$$

- $E_{\text{cms}}-2M_{\Lambda_c}=26\text{MeV}$ only!
- $\Lambda_c^+\Lambda_c^-$ produced in pairs with no additional accompany hadrons.
 - $e^+e^-\rightarrow\gamma^*\rightarrow\Lambda_c^+\Lambda_c^-$
- Clean backgrounds and well constrained kinematics.

• Typically, two ways to study Λ_c^+ decays:

• **Single Tag (ST):** detect only one of the $\Lambda_c^+\Lambda_c^-$

=> **Relative higher backgrounds**

=> **Higher efficiencies**

=> **Full reconstruction**


• **Double Tag (DT):** detect both of $\Lambda_c^+\Lambda_c^-$

=> **Smaller backgrounds**

=> **Missing mass technique**

=> **Lower efficiencies**




=> **Systematic in tag side are mostly cancelled**



Angular dependence analysis of $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ near threshold



PRL 120, 132001 (2018)




$$|G_E/G_M|^2(1 - \beta^2) = (1 - \alpha_{\Lambda_c})/(1 + \alpha_{\Lambda_c}).$$

\sqrt{s} (MeV)	α_{Λ_c}	$ G_E/G_M $
4574.5	$-0.13 \pm 0.12 \pm 0.08$	$1.14 \pm 0.14 \pm 0.07$
4599.5	$-0.20 \pm 0.04 \pm 0.02$	$1.23 \pm 0.05 \pm 0.03$


- One of the most basic observables that intimately related to **the internal structure** of the nucleon.
- One of the most challenging questions in contemporary physics is why and how quarks are confined into hadrons.
- The electromagnetic form factors (EMFFs) have been a powerful tool in understanding the structure of nucleons.
- First measurements of the EMFFs of the Λ_c^+

Observation of $\Lambda_c^+ \rightarrow n K_S \pi^+$

- ✓ No measurements for Λ_c^+ decay into a neutron yet.
- ✓ To confer the missing neutron, we define the variable M_{miss}^2 , which considers the beam energy constrain to improve resolution



567/pb data @ 4.6 GeV





scatter plots $M_{\pi^+\pi^-}$ versus M_{miss}^2

BESIII Preliminary results: $\mathbf{B}[\Lambda_c^+ \rightarrow n K_S \pi^+] = (1.82 \pm 0.23 \pm 0.11)\%$

First observation of Λ_c^+ decays to final states involving the neutron

Competition from Belle & BelleII





- Belle tags $\sim 36\text{K } \Lambda_c^+$, while BESIII now tags $15\text{K } \Lambda_c^+$ ($567/\text{pb}@4.6\text{GeV}$)
- By middle of 2019, BELLEII will have $5/\text{ab}$ data, 5x of BELLE data;
→ 180K tagged Λ_c^+ ;
- We will have 150K tagged Λ_c^+ , however, BESIII is very clean
- Many precise measurements at BESIII will reach to the level of systematic dominated
→ BESIII has advantages on backgrounds and systematics



World campaign on the Λ_c^+

	BESIII	Belle(-II)	LHCb
Λ_c^+ total yields	***	*****	*****
S/B ratio	*****	**	**
Systematic error	*****	***	**
Systematic research	*****	***	*
Semi-leptonic mode	*****	***	*
n/K_L -involved mode	*****	**	☆
Photon final state	*****	****	☆
Absolute measurement	*****	***	☆

- The threshold data at BESIII have systematic advantage over Belle(-II) and LHCb in the Λ_c^+ studies.
- This proposal holds an optimal time window to maximize the visibility of BESIII physics.

Measurement of the Decays $\Lambda_c \rightarrow \Sigma \pi \pi$ at Belle


Important Input for b physics

- stringent Fragmentation Function of b/c quark to baryon
 - [Eur. Phys. J. C12, 225 (2000); Eur. Phys. J. C 16, 597 (2000); Phys. Rev. D 85, 032008 (2012), Phys. Rev.D 66, 091101 (2002).]
 - Fragmentation Function (FF) is an important probe in experiment to test and calibrate QCD theory.

PhysRevD.85.032008

TABLE IV. Systematic uncertainties on the absolute scale of $f_{\Lambda_b}/(f_u + f_d)$.

Source	Error (%)
Bin-dependent errors	2.2
$\mathcal{B}(\Lambda_b^0 \rightarrow D^0 p X \mu^- \bar{\nu})$	2.0
Monte Carlo modelling	1.0
Backgrounds	3.0
Tracking efficiency	2.0
Γ_{sl}	2.0
Lifetime ratio	2.6
PID efficiency	2.5
Subtotal	6.3
$\mathcal{B}(\Lambda_c^+ \rightarrow p K^- \pi^+)$	26.0
Total	26.8





- Now $\mathcal{B}(pK^-\pi^+)$ are still dominated (6%)
- 20x data=> small than 2%

Contributions to Ξ_{cc}^{++} observation



- LHCb observed Ξ_{cc}^{++} from $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ and $\Xi_c^+ \pi^+$ decays
- Credits from theorists
 - $\tau(\Xi_{cc}^{++}) \approx 3 \tau(\Xi_{cc}^+)$ (Chang, Li, Wang, Karliner, et al.)
 - “Discovery channels of $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ and $\Xi_c^+ \pi^+$ was predicted benefited from BESIII Λ_c^+ measurements ” (Fu-Sheng Yu, et al, '17)



First Measurements of absolute BFs for Ξ_c



$$\begin{aligned}\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+) &= (1.80 \pm 0.50 \pm 0.14)\%, \\ \mathcal{B}(\Xi_c^0 \rightarrow \Lambda K^- \pi^+) &= (1.17 \pm 0.37 \pm 0.09)\%, \\ \mathcal{B}(\Xi_c^0 \rightarrow p K^- K^- \pi^+) &= (0.58 \pm 0.23 \pm 0.05)\%, \\ \mathcal{B}(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) &= (2.86 \pm 1.21 \pm 0.38)\%, \\ \mathcal{B}(\Xi_c^+ \rightarrow p K^- \pi^+) &= (0.45 \pm 0.21 \pm 0.07)\%.\end{aligned}$$


- Large errors
- Belle II will improve these to $\sim 10\%$
- BESIII has potential to improve these to be $< 5\%$

Λ_c^+ hadronic decay

- Study of SCS channels: $\Lambda_c^+ \rightarrow \Lambda K^+, p\pi^0, p\eta, n\pi^+, \Sigma^0 K^+, \Sigma^+ K^0$ etc.

From Prof. Hai-Yang Cheng's report.

Singly Cabibbo-suppressed modes: $\Lambda_c^+ \rightarrow p\pi^0, p\eta$




$$\pi^0 = (d\bar{d} - u\bar{u})/\sqrt{2}, \quad \eta = (d\bar{d} + u\bar{u} - s\bar{s})/\sqrt{3} \quad \text{for } \eta - \eta' \text{ mixing angle} = 19.5^\circ$$

$$A(\Lambda_c^+ \rightarrow p\pi^0) = (C_1 + C_2 + E_1 - E_2 - E_3)/\sqrt{2}$$

$$A(\Lambda_c^+ \rightarrow p\eta) = (2C_1 + C_2 + E_1 + E_2 + E_3)/\sqrt{3}$$

It is most likely that

$$\Gamma(\Lambda_c^+ \rightarrow p\eta) \gg \Gamma(\Lambda_c^+ \rightarrow p\pi^0)$$



$$\text{BESIII: } B[\Lambda_c^+ \rightarrow p\pi^0] < 2.7 \times 10^{-4}$$



$$M(\Lambda_c^+ \rightarrow n\pi^+) = \sqrt{2} M(\Lambda_c^+ \rightarrow p\pi^0),$$

- More precise comparison of the two BFs are desired to explore the interference of different non-factorizable diagrams and BESIII result support the theoretic prediction. It is predicted that $\text{Br}[\Lambda_c^+ \rightarrow n\pi^+] \sim 3.5 \times \text{Br}[\Lambda_c^+ \rightarrow p\pi^0]$ [Hai-Yang Cheng, arXiv: 1801.08625]



$\Lambda_c^+ \rightarrow \Lambda l^+ \nu_l$ decays

- Theoretical calculations on the BF ranges from 1.4% to
- BESIII performed the first absolute BF measurements.
- The BFs provide complementary information on determining



$$B[\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e] = (3.63 \pm 0.38 \pm 0.20)\%$$

$$B[\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu] = (3.49 \pm 0.46 \pm 0.26)\%$$

$$\Gamma[\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu] / \Gamma[\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e] = 0.96 \pm 0.16 \pm 0.04$$

Provides important input for calibrating the LQCD calculations.

Λ_c^+ data in PDG2015



Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)			
Hadronic modes with a p: $S = -1$ final states						
$p\bar{K}^0$	(3.21 ± 0.30) %					
$pK^-\pi^+$	(6.84 ^{+0.32} _{-0.40}) %					
$p\bar{K}^*(892)^0$	[q] (2.13 ± 0.30) %					
$\Delta(1232)^{++}K^-$	(1.18 ± 0.27) %			22.9%		
$\Lambda(1520)\pi^+$	[q] (2.4 ± 0.6) %			25.0%		
$pK^-\pi^+$ nonresonant	(3.8 ± 0.4) %			10.5%		
$p\bar{K}^0\pi^0$	(4.5 ± 0.6) %			13.3%		
$p\bar{K}^0\eta$	(1.7 ± 0.4) %			23.5%		
$p\bar{K}^0\pi^+\pi^-$	(3.5 ± 0.4) %			11.4%		
$pK^-\pi^+\pi^0$	(4.6 ± 0.8) %			13.0%		
$pK^*(892)^-\pi^+$	[q] (1.5 ± 0.5) %			33.3%		
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(5.0 ± 0.9) %			18.0%		
$\Delta(1232)K^*(892)$	seen					
$pK^-\pi^+\pi^+\pi^-$	(1.5 ± 1.0) × 10 ⁻³			66.7%		
$pK^-\pi^+\pi^0\pi^0$	(1.1 ± 0.5) %			45.4%		
Hadronic modes with a p: $S = 0$ final states						
$p\pi^+\pi^-$	(4.7 ± 2.5) × 10 ⁻³			45.4%		
$p f_0(980)$	[q] (3.8 ± 2.5) × 10 ⁻³			53.2%		
$p\pi^+\pi^+\pi^-\pi^-$	(2.5 ± 1.6) × 10 ⁻³			64.0%		
pK^+K^-	(1.1 ± 0.4) × 10 ⁻³			36.4%		
$p\phi$	[q] (1.12 ± 0.23) × 10 ⁻³					
$pK^+K^- \text{ non-}\phi$	(4.8 ± 1.9) × 10 ⁻⁴					
Hadronic modes with a hyperon: $S = -1$ final states						
$\Lambda\pi^+$	(1.46 ± 0.13) %			8.9%		
$\Lambda\pi^+\pi^0$	(5.0 ± 1.3) %			26.0%		
$\Lambda\rho^+$	< 6 %	CL=95%				
$\Lambda\pi^+\pi^+\pi^-$	(3.59 ± 0.28) %			7.8%		
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow$	(1.0 ± 0.5) %			20.0%		
$\Lambda\pi^+$						
$\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow$	(7.5 ± 1.4) × 10 ⁻³			18.7%		
$\Lambda\pi^-$						
Hadronic modes with a hyperon: $S = 0$ final states						
ΛK^+	(6.9 ± 1.4) × 10 ⁻⁴			20.3%		
$\Lambda K^+\pi^+\pi^-$	< 6 × 10 ⁻⁴	CL=90%				
$\Sigma^0 K^+$	(5.7 ± 1.0) × 10 ⁻⁴			17.5%		
$\Sigma^0 K^+\pi^+\pi^-$	< 2.9 × 10 ⁻⁴	CL=90%				
$\Sigma^+ K^+\pi^-$	(2.3 ± 0.7) × 10 ⁻³			30.4%		
$\Sigma^+ K^*(892)^0$	[q] (3.8 ± 1.2) × 10 ⁻³			31.6%		
$\Sigma^- K^+\pi^+$	< 1.3 × 10 ⁻³	CL=90%				
Doubly Cabibbo-suppressed modes						
$pK^+\pi^-$	< 3.1 × 10 ⁻⁴	CL=90%				
Semileptonic modes						
$\Lambda\ell^+\nu_\ell$	[r] (2.8 ± 0.4) %			17.2%		
$\Lambda e^+\nu_e$	(2.9 ± 0.5) %					
$\Lambda\mu^+\nu_\mu$	(2.7 ± 0.6) %			22.2%		





HTTP://PDG.LBL.GOV Page 32 Created: 10/6/2015 12

- ✓ Total branching fraction ~60%
- ✓ Lots of unknown decay channels
- ✓ Quite large uncertainties (>20%)

Λ_c decay asymmetries

4(6)-fold angular analysis of the cascade decays of $\Lambda_c \rightarrow pK_S, \Lambda\pi^+, \Sigma^+\pi^0$ and $\Sigma^0\pi^+$

PRD 100, 072004 (2019)



$$\sin \Delta_0 = -0.28 \pm 0.13 \pm 0.03$$

$\Lambda_c^+ \rightarrow$		pK_S^0	$\Lambda\pi^+$	$\Sigma^+\pi^0$	$\Sigma^0\pi^+$
α_{BP}^+	Predicted	-1.0 [16], 0.51 [11]	-0.70 [16], -0.67 [11]	0.71 [16], 0.92 [11]	0.70 [16], 0.92 [11]
		-0.49 [10], -0.90 [10]	-0.95 [10], -0.99 [10]	0.79 [10], -0.49 [10]	0.78 [10], -0.49 [10]
		-0.49 [17], -0.97 [18]	-0.96 [17], -0.95 [18]	0.83 [17], 0.43 [18]	0.83 [17], 0.43 [18]
		-0.66 [19], -0.90 [30]	-0.99 [19], -0.86 [30]	0.39 [19], -0.76 [30]	0.39 [19], -0.76 [30]
		-0.99 [20], -0.91 [31]	-0.99 [20], -0.94 [31]	-0.31 [20], -0.47 [31]	-0.31 [20], -0.47 [31]
PDG [2]		-0.91 ± 0.15	-0.45 ± 0.32		
This work		0.18 ± 0.43 ± 0.14	-0.80 ± 0.11 ± 0.02	-0.57 ± 0.10 ± 0.07	-0.73 ± 0.17 ± 0.07

- Best precisions on the hadronic weak decay asymmetries
- The transverse polarization is firstly studied and found to be non-zero with 2.1σ