Contribution ID: 35

## The SM expected branching ratio for $h\to\gamma\gamma$ and an excess for $h\to Z\gamma$

The recent measurements of  $h \to Z\gamma$  from ATLAS and CMS show an excess of the signal strength  $\mu_Z = (\sigma \cdot calB)_{\rm obs}/(\sigma \cdot calB)_{\rm SM} = 2.2 \pm 0.7$ , normalized as 1 in the standard model<sup>-</sup>(SM). If confirmed, it would be a signal of new physics (NP) beyond the SM. We study NP explanation for this excess. In general, for a given model, it also affects the process  $h \to \gamma\gamma$ . Since the measured branching ratio for this process agrees well with the SM prediction, the model is severely constrained. We find that a minimally fermion singlets and doublet extended NP model can explain simultaneously the current data for  $h \to Z\gamma$  and  $h \to \gamma\gamma$ . There are two solutions. Although both solutions enhance the amplitude of  $h \to Z\gamma$  to the observed one, in one of the solution although cannot be ruled out simply using branching ratio measurements alone. However, we find another solution that naturally enhances  $h \to Z\gamma$  to the measured value, but keeps the amplitude of  $h \to \gamma\gamma$  close to its SM prediction. We also comment on the phenomenology associated with these new fermions.

**Primary authors:** HE, Xiao-Gang (Shanghai Jiao Tong University); HUANG, Zhonglv (TDLI); LI, Ming-Wei (Shanghai Jiao Tong University); LIU, Chia-Wei (Tsung-Dao Lee Institute,Shanghai Jiao Tong University)

Presenter: LIU, Chia-Wei (Tsung-Dao Lee Institute, Shanghai Jiao Tong University)