



中国科学院大学

University of Chinese Academy of Sciences



CP violation at LHCb

Peilian Li (李佩莲)

on behalf of the LHCb collaboration

University of Chinese Academy of Sciences

第十届中国LHC物理会议 (CLHCP2024)

2024.11.15, Qingdao


Outline

- Introduction
- Recent results for
 - CKM angle γ and $\beta_{(s)}$
 - Direct CPV in B meson
 - CPV in Baryons
 - CPV in Charm (back up)
- Summary



*Disclaimer: many new and interesting results, only a few selected
More details in parallel sessions and [public page](#)*

Matter and anti-matter asymmetry



[PLB667 \(2008\) 1](#)

- Large matter-antimatter imbalance in the Universe




$$(n_B - n_{\bar{B}})/n_\gamma \sim 6 \times 10^{-10}$$

- CP violation in SM $\sim 10^{-17}$, not enough!
- Extra sources of CP violation necessitated



CKM matrix

$$V_{CKM} = \begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| e^{-i\beta} & -|V_{ts}| e^{i\beta_s} & |V_{tb}| \end{pmatrix} + \mathcal{O}(\lambda^5) \sim$$



- Key test of the SM: Verify unitarity of CKM matrix
- Only source of CP violation in SM



$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$$

Indirect search for New Physics

- Sensitivity to New Physics scale much **higher** than direct search: $1 \sim 10^4$ GeV

$$\mathcal{A}(\psi_i \rightarrow \psi_j + X) = \mathcal{A}_0 \left(\frac{c_{\text{SM}}}{v^2} + \frac{c_{\text{NP}}}{\Lambda^2} \right)$$




NP scale: Λ



- **Statistics or precision essential** for flavour physics
 - NP scale, i.e. Dim = 6, proportional to $\sqrt[4]{\text{statistics}}$
- Would tell not only if there is NP but also which flavour it couples to

Types of CP Violation




- CP-violating nature of weak interaction has multiple manifestations
- Requires **two interfering amplitudes** with different strong and weak phases




credit: Wenbin Qian

Types of CP Violation

- CP-violating nature of weak interaction has multiple manifestations
- Requires **two interfering amplitudes** with different strong and weak phases



credit: Wenbin Qian




We continue to discover new types of CP violation

[PRL124 \(2020\) 031801](#)
[PRD101 \(2020\) 012006](#)


CKM angle γ

- Relative weak phase γ in interference between $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$ transition
- Measured with tree-level decays, theoretically **clean observable** ($\delta\gamma/\gamma \sim 10^{-7}$)



CKM angle γ


- Relative weak phase γ in interference between $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$ transition
- Measured with tree-level decays, theoretically **clean observable** ($\delta\gamma/\gamma \sim 10^{-7}$)



CKM angle γ

- Relative weak phase γ in interference between $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$ transition
- Measured with tree-level decays, theoretically **clean observable** ($\delta\gamma/\gamma \sim 10^{-7}$)


$$\gamma = \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right)$$



Latest γ combination

LHCb-CONF-2024-004

- **19** LHCb B decay measurements + **11** D decay measurements + 27 external inputs
- **29** physics parameters of interest + additional nuisance parameters



$$\gamma = (64.6 \pm 2.8)^\circ$$


- Surpass LHCb design: 4°
- Consistent with SM predictions $(65.5^{+0.09}_{-2.65})^\circ$

γ measurement in $B^+ \rightarrow DK^{*+} (\rightarrow K_S^0 \pi^+)$

LHCb-PAPER-2024-023
arXiv: 2410.21115




- CP eigenstates $D^0 \rightarrow hh, \pi\pi\pi\pi$
- Suppressed decays $D^0 \rightarrow K\pi, K\pi\pi\pi$
- Self-conjugated multi-body decays $D^0 \rightarrow K_S^0 hh$
- Inputs for $D^0 \rightarrow K_S^0 hh, \pi\pi\pi\pi$ from BESIII

[PRD106.092004, PRD82(2010)112006, PRD102(2020)]



$$\gamma = (63 \pm 13)^\circ$$



First observation of $B^+ \rightarrow [D^0]_{\pi K, \pi K \pi \pi} K^{*+}$




γ measurement in $B_s^0 \rightarrow D_s^\mp K^\pm$

LHCb-PAPER-2024-020
in preparation

- Time-dependent CP violation measurement
- Interference between mixing and decay \rightarrow relative phase difference of $\gamma - 2\beta_s$



$\oplus\oplus B_s^0 \rightarrow D_s^- K^+$ $\oplus\oplus \bar{B}_s^0 \rightarrow D_s^- K^+$ $\oplus\oplus$ Untagged $D_s^- K^+$
 $\oplus\oplus B_s^0 \rightarrow D_s^+ K^-$ $\oplus\oplus \bar{B}_s^0 \rightarrow D_s^+ K^-$ $\oplus\oplus$ Untagged $D_s^+ K^-$




$$\gamma = (74 \pm 11)^\circ$$

CKM angles $\beta_{(s)}$

- Tree dominated $b \rightarrow c\bar{c}s$ transition offers access to the CKM angle $\beta_{(s)}$
- CP violation in the interference between mixing and decays

$$\beta_{(s)} = \arg\left(-\frac{V_{cq}V_{cb}^*}{V_{tq}V_{tb}^*}\right)$$

$$A_{CP}(t) = \frac{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) - \Gamma_{B_{(s)}^0 \rightarrow f}(t)}{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) + \Gamma_{B_{(s)}^0 \rightarrow f}(t)} \propto -D_{\text{tag}} \cdot D_t \cdot \eta_f \cdot \sin 2\beta_{(s)} \cdot \sin(\Delta m_{(s)}t)$$




CKM angles $\beta_{(s)}$

- Tree dominated $b \rightarrow c\bar{c}s$ transition offers access to the CKM angle $\beta_{(s)}$
- CP violation in the interference between mixing and decays

$$\beta_{(s)} = \arg\left(-\frac{V_{cq}V_{cb}^*}{V_{tq}V_{tb}^*}\right)$$



$$A_{CP}(t) = \frac{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) - \Gamma_{B_{(s)}^0 \rightarrow f}(t)}{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) + \Gamma_{B_{(s)}^0 \rightarrow f}(t)} \propto -D_{\text{tag}} \cdot D_t \cdot \eta_f \cdot \sin 2\beta_{(s)} \cdot \sin(\Delta m_{(s)}t)$$



World best measurements!

$$S_{\psi K_S^0}^{\text{Run 2}} = 0.716 \pm 0.013 \pm 0.008$$

$$\phi_s^{c\bar{c}s} = -0.031 \pm 0.018 \text{ rad}$$



$$\begin{aligned} \phi_s^{\text{CKMFitter}} &\approx -2\beta_s \\ &= (-0.0368_{-0.0009}^{+0.0006}) \text{ rad} \end{aligned}$$


Direct CPV in $B^+ \rightarrow J/\psi\pi^+$

New

LHCb-PAPER-2024-31
in preparation

- ⦿ O(1%) direct CP violation expected in $B^+ \rightarrow J/\psi\pi^+$ [PRD 49 (1994) 5904, PRD 52 (1995) 242]
- ⦿ Important control channel to understand penguin effects in $\sin 2\beta$ measurement in $B^0 \rightarrow J/\psi K^0$

[PRD 79 (2009) 014030, JHEP 03 (2015) 145]




Direct CPV in $B^+ \rightarrow J/\psi\pi^+$

New





LHCb-PAPER-2024-31
in preparation

- O(1%) direct CP violation expected in $B^+ \rightarrow J/\psi\pi^+$ [PRD 49 (1994) 5904, PRD 52 (1995) 242]
- Important control channel to understand penguin effects in $\sin 2\beta$ measurement in $B^0 \rightarrow J/\psi K^0$

[PRD 79 (2009) 014030, JHEP 03 (2015) 145]



$$\mathcal{R}_{\pi/K} \equiv \frac{B(B^+ \rightarrow J/\psi\pi^+)}{B(B^+ \rightarrow J/\psi K^+)} = \frac{N_\pi}{N_K} \times \frac{\epsilon_K}{\epsilon_\pi}$$




Direct CPV in $B^+ \rightarrow J/\psi\pi^+$

New

LHCb-PAPER-2024-31
in preparation


First evidence for direct CP violation in beauty decays to charmonium final states (3.2σ)



- Taking previous LHCb result of $A^{CP}(B^+ \rightarrow J/\psi K^+)$ to extract: [Phys. Rev. D 95, 052005 (2017)]

$$A^{CP}(B^+ \rightarrow J/\psi\pi^+) = (1.51 \pm 0.50 \pm 0.11) \times 10^{-2}$$

$$A(B^+ \rightarrow J/\psi\pi^+) = -\lambda\mathcal{A}(1 + ae^{i\theta}e^{i\gamma}) \quad A(B^+ \rightarrow J/\psi K^+) = (1 - \lambda^2/2)\mathcal{A}'(1 + \epsilon a'e^{i\theta'}e^{i\gamma})$$






- SU(3) flavour symmetry: $a = a', \theta = \theta'$
- Constraints on the relative size (a) and strong phase difference (θ) between penguin and tree contributions

See more details in Manshu Li's talk (Thursday afternoon)



CP violation in baryonic decays?

- Baryons crucial for asymmetries in Universe, **no CP violation in baryons observed yet**
- CPV: *b* baryons $\mathcal{O}(1 - 10\%)$, *c* baryons $\mathcal{O}(0.1\%)$, hyperon $\mathcal{O}(0.001 - 0.01\%)$




- **Puzzling situation:** similar Λ_b^0 production as B^+ , huge significance of CPV in B^+ , **none in Λ_b^0 ?**

Charmless three-body b decays



- Complex CP violation pattern in multi-body B decays, as large as 80%
- Interesting to search for CP violation in $\Lambda_b^0 \rightarrow \Lambda h^+ h'^-$

PRL124 (2020) 031801
 PRD101 (2020) 012006



First observation of $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda\pi^+\pi^-(K^-)$

New

CERN Seminar by W. Qian

LHCb-PAPER-2024-043
in preparation

- Systematic study of $\Lambda_b^0/\Xi_b^0 \rightarrow \Lambda h^+ h'^-$ with control mode to reduce systematic uncertainty

$$\frac{\mathcal{B}(\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-)} = \frac{N_{\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-}}{N_{\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-}} \times \frac{\epsilon_{\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-}}{\epsilon_{\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-}} \times \frac{f_{\Lambda_b^0}}{f_{\Lambda_b^0(\Xi_b^0)}}$$

First observation of $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda\pi^+\pi^-(K^-)$

New

- Systematic study of $\Lambda_b^0/\Xi_b^0 \rightarrow \Lambda h^+ h'^-$ with control mode to reduce systematic uncertainty



$$\frac{\mathcal{B}(\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-)} = \frac{N_{\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-}}{N_{\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-}} \times \frac{\epsilon_{\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda \pi^+) \pi^-}}{\epsilon_{\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda h^+ h'^-}} \times \frac{f_{\Lambda_b^0}}{f_{\Lambda_b^0(\Xi_b^0)}}$$

First observations of $\Xi_b^0 \rightarrow \Lambda K^- \pi^+$ & $\Lambda_b^0 \rightarrow \Lambda \pi^+ \pi^-$ and evidence for $\Xi_b^0 \rightarrow \Lambda \pi^+ \pi^- (4\sigma)$

$$\mathcal{B} = (10.4 \pm 1.4 \pm 1.2 \pm 3.5) \times 10^{-6}$$


$$\mathcal{B} = (5.3 \pm 0.4 \pm 0.5 \pm 0.5) \times 10^{-6}$$

$$\mathcal{B} = (11.0 \pm 2.6 \pm 1.4 \pm 3.8) \times 10^{-6}$$




Evidence of CP violation in baryonic decays

New



$$\mathcal{A}^{CP}(\Lambda_b^0/\Xi_b^0 \rightarrow f) \equiv \frac{\Gamma(\Lambda_b^0/\Xi_b^0 \rightarrow f) - \Gamma(\bar{\Lambda}_b^0/\bar{\Xi}_b^0 \rightarrow \bar{f})}{\Gamma(\Lambda_b^0/\Xi_b^0 \rightarrow f) + \Gamma(\bar{\Lambda}_b^0/\bar{\Xi}_b^0 \rightarrow \bar{f})}$$


Preliminary



Evidence of CP violation in baryonic decays


New

LHCb-PAPER-2024-043
in preparation




$$\mathcal{A}^{CP}(\Lambda_b^0/\Xi_b^0 \rightarrow f) \equiv \frac{\Gamma(\Lambda_b^0/\Xi_b^0 \rightarrow f) - \Gamma(\bar{\Lambda}_b^0/\bar{\Xi}_b^0 \rightarrow \bar{f})}{\Gamma(\Lambda_b^0/\Xi_b^0 \rightarrow f) + \Gamma(\bar{\Lambda}_b^0/\bar{\Xi}_b^0 \rightarrow \bar{f})}$$

Preliminary



$\Lambda_b^0 \rightarrow \Lambda K^+ K^-$




$\bar{\Lambda}_b^0 \rightarrow \bar{\Lambda} K^+ K^-$

$$\Delta A_{CP} = 0.083 \pm 0.023 \pm 0.016$$

First evidence for CP violation, 3.1σ

$\Lambda_b^0 \rightarrow \Lambda K^+ K^-$ yields




Preliminary $m^2(\Lambda K^+)$ [GeV^2/c^4]

Evidence of CP violation in baryonic decays

CERN Seminar by W. Qian

New



LHCb-PAPER-2024-043
in preparation



[PRD107 \(2023\) 053009](#)



Evidence of CP violation in baryonic decays

New



Evidence of CP violation in baryonic decays

New



region
 $\Delta A_{CP}(\Lambda\phi) = 0.150 \pm 0.055 \pm 0.021$
Consistent with 0 within 2.5σ
Predicted CPV (resonant), $\sim 1.5\%$

Preliminary $m^2(\Lambda K^+)$ [GeV²/c⁴]

PRD107 (2023) 053009

See more details in Chenxu Yu's talk (Friday afternoon)

CP violation in $\Lambda_b^0 \rightarrow ph$



New LHCb-PAPER-2024-048
in preparation

- Sizeable CP violation found in $B_{(s)}^0 \rightarrow h^+h'^-$
- Golden channel in baryon decays, with predictions of CPV ranges from few percent to 30%

- No clear asymmetry observed directly

$$A_{CP}^{pK^-} = (-1.14 \pm 0.67 \pm 0.36) \%$$


$$A_{CP}^{p\pi^-} = (-0.20 \pm 0.83 \pm 0.37) \%$$



CP violation in $\Lambda_b^0 \rightarrow ph$

New LHCb-PAPER-2024-048
in preparation

- Sizeable CP violation found in $B_{(s)}^0 \rightarrow h^+h'^-$
- Golden channel in baryon decays, with predictions of CPV ranges from few percent to 30%



- No clear asymmetry observed directly

$$A_{CP}^{pK^-} = (-1.14 \pm 0.67 \pm 0.36) \%$$

$$A_{CP}^{p\pi^-} = (-0.20 \pm 0.83 \pm 0.37) \%$$

Why CP violation so small?

$$A = a_1 e^{i(\delta_1 + \phi_1)} + a_2 e^{i(\delta_2 + \phi_2)}$$


$$\bar{A} = a_1 e^{i(\delta_1 - \phi_1)} + a_2 e^{i(\delta_2 - \phi_2)}$$

$$A_{CP} = \frac{|A|^2 - |\bar{A}|^2}{|A|^2 + |\bar{A}|^2} \propto \sin(\delta_1 - \delta_2) \sin(\phi_1 - \phi_2)$$

$$\delta_1 - \delta_2 \sim 0?$$

- Small strong phase difference?
- Cancellation between S, P waves? [arXiv:2409.02821](https://arxiv.org/abs/2409.02821)

See more details in Xinchun Dai's talk (Thursday afternoon)



What's more?

- Decay parameter, first proposed by Lee and Yang (1959) to study hyperon decays

The diagram illustrates the decay of a hyperon (Λ) into a nucleon (p) and a pion (π^-). A coordinate system (x', y', z') is centered on the nucleon. The pion momentum vector P_π is shown at an angle θ relative to the z' -axis. The nucleon momentum vector P_p is shown at an angle γ relative to the z' -axis. The pion polarization vector is $-\alpha P_\pi \cos\theta$. The decay parameter P_p is defined as:

$$P_p = \frac{(\alpha + P_\Lambda \cos\theta)z' + \beta P_\Lambda x' + \gamma P_\Lambda y'}{1 + \alpha P_\Lambda \cos\theta}$$

Below the diagram are two black and white portraits: Chen Ning Yang on the left and Tsung-Dao Lee on the right.

$$\alpha_{\mp} = \pm \frac{2\Re(S^*P)}{|S|^2 + |P|^2} = \pm \frac{2|S||P| \cos(\delta \pm \phi)}{|S|^2 + |P|^2}$$

$$\beta_{\mp} = \pm \frac{2\Im(S^*P)}{|S|^2 + |P|^2} = \pm \frac{2|S||P| \sin(\delta \pm \phi)}{|S|^2 + |P|^2}$$

$$\gamma = \frac{|S|^2 - |P|^2}{|S|^2 + |P|^2} \quad \alpha^2 + \beta^2 + \gamma^2 = 1$$

- New CP observables

* δ, ϕ : strong and weak phase difference between S and P waves


$$A_\alpha = \frac{\alpha_+ + \alpha_-}{\alpha_+ - \alpha_-} = -\tan(\delta) \tan(\phi) \quad R_{\beta_1} = \frac{\beta_+ + \beta_-}{\alpha_+ - \alpha_-} = \tan(\phi) \quad R_{\beta_2} = \frac{\beta_+ - \beta_-}{\alpha_+ - \alpha_-} = -\tan(\delta)$$

Decay parameters of $\Lambda_b^0 \rightarrow \Lambda_c^+ h^-$




$$\Lambda_b^0 \rightarrow \Lambda_c^+ (\rightarrow \Lambda h^+) h^-, \Lambda \rightarrow p \pi^-$$

$$\alpha_c, \beta_c, \gamma_c: \Lambda_c^+ \rightarrow \Lambda h^+$$

$$\frac{d\Phi}{d\Omega} \propto (1 + \alpha_b \alpha_c \cos \theta_1 + \alpha_c \alpha_s \cos \theta_2 + \alpha_b \alpha_s \cos \theta_1 \cos \theta_2 - \alpha_b \gamma_c \alpha_s \sin \theta_1 \sin \theta_2 \cos \phi_2 + \alpha_b \beta_c \alpha_s \sin \theta_1 \sin \theta_2 \sin \phi_2)$$




- First determination of α in $\Lambda_b^0 \rightarrow \Lambda_c^+ h^-$ decays $\mathcal{O}(0.9\%)$
- Most precise determinations $\alpha, (\beta, \gamma)$
- Confirmation of $\alpha(\Lambda \rightarrow p \pi^-)$ from BESIII
- Pave the way for other decay parameter measurements




See more details in Yuhao Wang's talk (Friday afternoon)


Summary

- ✓ LHCb dominates the world average of many measurements in CKM and CPV
- ✓ Various precise measurement of CP violation in beauty and charm decays using LHCb Run 1+2 data
- ✓ First evidence of CP violation in $B^+ \rightarrow J/\psi\pi^+$ and $\Lambda_b^0 \rightarrow \Lambda K^+ K^-$
- ✓ Run 3 provides great opportunities for further test of the SM and search for new physics



Run 3 has collected more data than Run 1+2, a lot more new results to come!







Thank you

Back up slides


LHCb detector

General purpose detector specialised in beauty and charm hadrons



$$2 < \eta < 5$$




LHCb performance:
[JINST 14 \(2019\) P04013](https://arxiv.org/abs/1904.01153)



LHCb detector


General purpose detector specialised in beauty and charm hadrons

- Daughters of b & c hadron decays: $p_T \sim \mathcal{O}(1 \text{ GeV}/c)$, flight distance $L \sim 1\text{mm}$



$$2 < \eta < 5$$

LHCb performance:
JINST 14 (2019) P04013



LHCb detector


General purpose detector specialised in beauty and charm hadrons

- Daughters of b & c hadron decays: $p_T \sim \mathcal{O}(1 \text{ GeV}/c)$, flight distance $L \sim 1\text{mm}$


Precise vertex measurements,
 $\sigma(IP_x) \sim 35 \mu\text{m}$

Excellent decay time resolution $\sim 43 \text{ fs}$ for B_s^0

proton beam




Excellent momentum resolution $\sim 0.5\%$



$2 < \eta < 5$

Excellent particle identification
 $\epsilon(K) \approx 95\%$
 misID $p(\pi \rightarrow K) \approx 5\%$
 $\epsilon(\mu) \approx 97\%$




LHCb performance:
[JINST 14 \(2019\) P04013](https://arxiv.org/abs/1904.00001)

Control of penguin contribution


- $\sigma(\phi_s) \sim 0.016$ comparable with the theoretical estimation of $\Delta\phi_s^{penguin} \sim 1^\circ \approx 0.017$, better control of penguin effect necessary
- Combined analysis of penguin contributions in ϕ_s and ϕ_d ($\sin 2\beta$), using SU(3) flavour symmetry

$$\phi_d = \sin(2\beta^{tree}) + \Delta\phi_d^{penguin} + \phi_d^{NP}$$

$$\phi_s = \phi_s^{tree} + \Delta\phi_s^{penguin} + \phi_s^{NP}$$




J.Phys.G 48 (2021) 6, 065002



Method to measure ΔA^{CP}

➤ CP asymmetries



➤ CP asymmetry difference

$$\Delta A^{CP} \equiv A^{CP}(B^{\mp} \rightarrow J/\psi\pi^{\mp}) - A^{CP}(B^{\mp} \rightarrow J/\psi K^{\mp})$$

$$= \Delta a^{\text{raw}} - \cancel{\Delta a^{\text{prod}}} - \Delta a^{\text{det}} - \Delta a^{\text{PID}}$$

credit: Manshu Li

γ measurements

B decay	D decay	Ref.	Dataset	Status since Ref. [14]
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^\pm h'^\mp$	[35]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+ h^- \pi^+ \pi^-$	[19]	Run 1&2	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K^\pm \pi^\mp \pi^+ \pi^-$	[36]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^\pm h'^\mp \pi^0$	[37]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 h^+ h^-$	[38]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 K^\pm \pi^\mp$	[39]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow D^* h^\pm$	$D \rightarrow h^\pm h'^\mp$ (PR)	[35]	Run 1&2	<i>As before</i>
$B^\pm \rightarrow D^* h^\pm$	$D \rightarrow K_S^0 h^+ h^-$ (PR)	[20]	Run 1&2	New
$B^\pm \rightarrow D^* h^\pm$	$D \rightarrow K_S^0 h^+ h^-$ (FR)	[21]	Run 1&2	New
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^\pm h'^\mp$	[22] [†]	Run 1&2	Updated
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^\pm \pi^\mp \pi^+ \pi^-$	[22] [†]	Run 1&2	Updated
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow K_S^0 h^+ h^-$	[22] [†]	Run 1&2	New
$B^\pm \rightarrow Dh^\pm \pi^+ \pi^-$	$D \rightarrow h^\pm h'^\mp$	[40]	Run 1	<i>As before</i>
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^\pm h'^\mp$	[23]	Run 1&2	Updated
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^\pm \pi^\mp \pi^+ \pi^-$	[23]	Run 1&2	Updated
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_S^0 h^+ h^-$	[24]	Run 1&2	Updated
$B^0 \rightarrow D^\mp \pi^\pm$	$D^+ \rightarrow K^- \pi^+ \pi^+$	[41]	Run 1	<i>As before</i>
$B_s^0 \rightarrow D_s^\mp K^\pm$	$D_s^+ \rightarrow h^+ h^- \pi^+$	[25, 42] [†]	Run 1&2	Updated
$B_s^0 \rightarrow D_s^\mp K^\pm \pi^+ \pi^-$	$D_s^+ \rightarrow h^+ h^- \pi^+$	[43]	Run 1&2	<i>As before</i>


D decay	Observable(s)	Ref.	Dataset	Status since Ref. [14]
$D^0 \rightarrow h^+ h^-$	ΔA_{CP}	[44–46]	Run 1&2	<i>As before</i>
$D^0 \rightarrow K^+ K^-$	$A_{CP}(K^+ K^-)$	[46–48]	Run 2	<i>As before</i>
$D^0 \rightarrow h^+ h^-$	$y_{CP} - y_{CP}^{K^- \pi^+}$	[49, 50]	Run 1&2	<i>As before</i>
$D^0 \rightarrow h^+ h^-$	ΔY	[51–54]	Run 1&2	<i>As before</i>
$D^0 \rightarrow K^+ \pi^-$ (double tag)	$R^\pm, (x'^\pm)^2, y'^\pm$	[55]	Run 1	<i>As before</i>
$D^0 \rightarrow K^+ \pi^-$ (single tag)	$R_{K\pi}, A_{K\pi}, c_{K\pi}^{(i)}, \Delta c_{K\pi}^{(i)}$	[27, 56]	Run 1&2	Updated
$D^0 \rightarrow K^\pm \pi^\mp \pi^+ \pi^-$	$(x^2 + y^2)/4$	[57]	Run 1	<i>As before</i>
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	x, y	[58]	Run 1	<i>As before</i>
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[59]	Run 1	<i>As before</i>
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[60, 61]	Run 2	<i>As before</i>
$D^0 \rightarrow \pi^+ \pi^- \pi^0$	ΔY^{eff}	[26]	Run 2	New

Time-dependent CPV in $B^0 \rightarrow D^+D^-$

New

arXiv: 2409.03009

$$\frac{d\Gamma(t, d)}{dt} \propto e^{-t/\tau_{B^0}} (1 + d C_{D^+D^-} \cos \Delta m_d t - d S_{D^+D^-} \sin \Delta m_d t)$$



- CP asymmetry observed in $B^0 \rightarrow D^+D^-$ for the first time with a significance exceeding 6σ

$$S_{D^+D^-} = -0.552 \pm 0.100 \text{ (stat)} \pm 0.010 \text{ (syst)}$$




$$C_{D^+D^-} = 0.128 \pm 0.103 \text{ (stat)} \pm 0.010 \text{ (syst)}$$

[PRL117 (2016) 261801, BaBar PRD79 (2009) 032002] [Belle PRD85 (2012) 091106]

- LHCb combination:



$$S_{D^+D^-} = -0.549 \pm 0.085 \text{ (stat)} \pm 0.015 \text{ (syst)}$$

$$C_{D^+D^-} = 0.162 \pm 0.088 \text{ (stat)} \pm 0.009 \text{ (syst)}$$





CP violation in charm sector

- GIM mechanism very effective for charm decays, SM loops highly suppressed
- Tiny weak phases in first two generations of CKM matrix
- Oscillation and CPV ($\leq 10^{-3}$)
- Long distance contribution comparable/larger than short distance



CP violation in charm sector

- GIM mechanism very effective for charm decays, SM loops highly suppressed
- Tiny weak phases in first two generations of CKM matrix
- Oscillation and CPV ($\leq 10^{-3}$)
- Long distance contribution comparable/larger than short distance



Breakthroughs by LHCb thanks to huge statistics:

First observation of CPV in $D^0 \rightarrow h^+ h^-$ decays

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-15.4 \pm 2.9) \times 10^{-4} \quad [\text{PRL}(2019)211803]$$

Evidence of CPV in $D^0 \rightarrow \pi^+ \pi^-$ decay

$$A_{CP}(\pi^+ \pi^-) = (23.2 \pm 6.1) \times 10^{-4} \quad (3.8\sigma) \quad [\text{PRL}(2023)211803]$$


Direct CP violation in $D^+ \rightarrow K^+ K^- \pi^+$

New

[arXiv:2409.01414](https://arxiv.org/abs/2409.01414)

- Search for localised CP violation in the phase space of $D^+ \rightarrow K^+ K^- \pi^+$ (S) decay
- Control channel $D_s^+ \rightarrow K^+ K^- \pi^+$ (C) to subtract nuisance asymmetries

$$\Delta A_{CP}^i = A_{\text{raw}}^{i,S} - A_{\text{raw}}^{i,C} - \Delta A_{\text{raw}}^{\text{global}}$$



Direct CP violation in $D^+ \rightarrow K^+ K^- \pi^+$

New

arXiv:2409.01414



- Search for localised CP violation in the phase space of $D^+ \rightarrow K^+ K^- \pi^+$ (S) decay
- Control channel $D_s^+ \rightarrow K^+ K^- \pi^+$ (C) to subtract nuisance asymmetries

$$\Delta A_{CP}^i = A_{\text{raw}}^{i,S} - A_{\text{raw}}^{i,C} - \Delta A_{\text{raw}}^{\text{global}}$$

$$A_{CP|S}^{\phi\pi^+} = (0.95 \pm 0.43 \pm 0.26) \times 10^{-3}$$

$$A_{CP|S}^{\bar{K}^{*0}K^+} = (-0.26 \pm 0.56 \pm 0.18) \times 10^{-3}$$

p-values (2.3-14.1%) compatible with no CPV



Direct CP violation in $D^+ \rightarrow K^+ K^- \pi^+$

New


[arXiv:2409.01414](https://arxiv.org/abs/2409.01414)

- Search for localised CP violation in the phase space of $D^+ \rightarrow K^+ K^- \pi^+$ (S) decay
- Control channel $D_s^+ \rightarrow K^+ K^- \pi^+$ (C) to subtract nuisance asymmetries


$$\Delta A_{CP}^i = A_{\text{raw}}^{i,S} - A_{\text{raw}}^{i,C} - \Delta A_{\text{raw}}^{\text{global}}$$

- Test-statistic to extract a p-value for the hypothesis of no localised CP violation

$$\chi^2(\mathcal{S}_{\Delta CP}) = \sum_i^{N_{\text{bins}}} (\mathcal{S}_{\Delta CP}^i)^2, \quad \mathcal{S}_{\Delta CP}^i = \frac{\Delta A_{CP}^i}{\sigma_{\Delta A_{CP}^i}}$$



$D^+ \rightarrow K^+ K^- \pi^+$ (S)



$D_s^+ \rightarrow K^+ K^- \pi^+$ (C)

Direct CP violation in $D^+ \rightarrow K^+ K^- \pi^+$

New

arXiv:2409.01414


- Search for localised CP violation in the phase space of $D^+ \rightarrow K^+ K^- \pi^+$ (S) decay
- Control channel $D_s^+ \rightarrow K^+ K^- \pi^+$ (C) to subtract nuisance asymmetries

$$\Delta A_{CP}^i = A_{\text{raw}}^{i,S} - A_{\text{raw}}^{i,C} - \Delta A_{\text{raw}}^{\text{global}}$$


- p-values (2.3-14.1%) compatible with absence of localised CP violation in Dalitz plot

- Test-statistic to extract a p-value for the hypothesis of no localised CP violation


$$\chi^2(\mathcal{S}_{\Delta CP}) = \sum_i^{N_{\text{bins}}} (\mathcal{S}_{\Delta CP}^i)^2, \quad \mathcal{S}_{\Delta CP}^i = \frac{\Delta A_{CP}^i}{\sigma_{\Delta A_{CP}^i}}$$



$D^+ \rightarrow K^+ K^- \pi^+$ (S)





$D_s^+ \rightarrow K^+ K^- \pi^+$ (C)



Time-dependent CP violation in $D^0 \rightarrow \pi^+ \pi^- \pi^0$

Phys. Rev. Lett. 133 (2024) 101803


- First measurement of time-dependent CP violation in SCS mode



$$A_{CP}(f_{CP}, t) \equiv \frac{\Gamma_{D^0 \rightarrow f_{CP}}(t) - \Gamma_{\bar{D}^0 \rightarrow f_{CP}}(t)}{\Gamma_{D^0 \rightarrow f_{CP}}(t) + \Gamma_{\bar{D}^0 \rightarrow f_{CP}}(t)}$$

$$\approx a_{f_{CP}}^{\text{dir}} + \Delta Y_{f_{CP}} \frac{t}{\tau_{D^0}}$$

$$A_{\text{meas}}(\langle t/\tau_{D^0} \rangle_i) \equiv \frac{N_{D^0}^i - N_{\bar{D}^0}^i}{N_{D^0}^i + N_{\bar{D}^0}^i}$$



$$\Delta Y_{f_{CP}} \approx \frac{\eta_{f_{CP}}}{2} \left[\left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) x \sin \phi - \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) y \cos \phi \right]$$

- No evidence for time-dependent CP violation, constant with world average

$$\Delta Y \equiv \eta_{CP} \Delta Y_{f_{CP}} = (-1.3 \pm 6.3 \pm 2.4) \times 10^{-4}$$

Time-dependent CP violation in $D^0 \rightarrow K\pi$

LHCb-PAPER-2024-008

- Interference between mixing and decay for favoured RS and suppressed WS decays



$$R_{K\pi}^+ = \frac{\Gamma(D^0(t) \rightarrow K^+\pi^-)}{\Gamma(\bar{D}^0 \rightarrow K^-\pi^+)}; \quad R_{K\pi}^- = \frac{\Gamma(\bar{D}^0(t) \rightarrow K^-\pi^+)}{\Gamma(D^0 \rightarrow K^+\pi^-)};$$

DCS over CF amplitude

$$R_{K\pi}^\pm(t) \approx \boxed{R_{K\pi}} (1 \pm A_{K\pi}) + R_{K\pi} (1 \pm A_{K\pi}) (c_{K\pi} \pm \Delta c_{K\pi}) \left(\frac{t}{\tau_{D^0}}\right) + (c'_{K\pi} \pm \Delta c'_{K\pi}) \left(\frac{t}{\tau_{D^0}}\right)^2$$




CPV observables: $A_{K\pi}$ (in decays), $\Delta c_{K\pi}$ (in interference), $\Delta c'_{K\pi}$ (in mixing).

Mixing observables: $c_{K\pi}$, $c'_{K\pi}$

Time-dependent CP violation in $D^0 \rightarrow K\pi$

arXiv:2407.18001

- Measured with yields: RS ~ 400 M, WS ~ 1.6 M



$R_{K\pi}$	$(343.1 \pm 2.0) \times 10^{-5}$	
$c_{K\pi}$	$(51.4 \pm 3.5) \times 10^{-4}$	Mixing parameter Evidence of non 0
$c'_{K\pi}$	$(13.1 \pm 3.7) \times 10^{-6}$	
$A_{K\pi}$	$(-7.1 \pm 6.0) \times 10^{-3}$	} No CPV
$\Delta c_{K\pi}$	$(3.0 \pm 3.6) \times 10^{-4}$	
$\Delta c'_{K\pi}$	$(-1.9 \pm 3.8) \times 10^{-6}$	

$$c_{K\pi} \approx y_{12} \cos \phi_f^\Gamma \cos \Delta_f + x_{12} \cos \phi_f^M \sin \Delta_f$$