



# Higgs mass and width at CMS

**C.Zhang (IHEP)**





**Institute of High Energy Physics**  
Chinese Academy of Sciences



14Nov2024, Qingdao



# Introduction to $m_H$

- $m_H$  is the only one free parameter in the SM Higgs sector and determines all other properties of Higgs boson ( couplings, BR... )
- SM self-consistency
- $m_H$  and  $m_t$  determine EW vacuum



# Introduction to $H \rightarrow ZZ, H \rightarrow \gamma\gamma$


- The Higgs boson discovery has marked the LHC Run1
- LHC Run2 and Run3 are the eras of precision measurements of the Higgs boson
- $H \rightarrow ZZ, H \rightarrow \gamma\gamma$ , two golden channels serve Higgs measurements with clear signals





# $m_H$ measurement in $H \rightarrow \gamma\gamma$

- Previous analysis with 2016 Run2 dataset ( $35.9\text{fb}^{-1}, 13\text{TeV}$ )
- Expected event yield  $\sim 1900$  events with  $S/B \sim 1:10$  at peak range
- Larger event yield and worse reconstruction precision (w.r.t  $H \rightarrow ZZ$ )
  - Final precision is driven by systematic uncertainties
  - Put a lot of effort on calibrations for detector effects, energy leak, material effect, non-linearity responses and etc.



# $m_H$ measurement in $H \rightarrow \gamma\gamma$

- Event categorisation

- Di-photon MVA discriminant  $D_{\gamma\gamma}$   
(higher score for better mass resolution and higher photon purity)

- VBF MVA discriminant  $D_{VBF}$  (trained vs  $gg \rightarrow H$  and bkg.)


- Enhanced VBF MVA discriminant

$$D_{eVBF}$$

- Systematic uncertainties

- $Z \rightarrow ee$  for energy calibration


- $e/\gamma$  extrapolation



**Table 1**

The observed impact of the different uncertainties on the measurement of  $m_H$ .

Source	Contribution (GeV)
Electron energy scale and resolution corrections	0.10
Residual $p_T$ dependence of the photon energy scale	0.11
Modelling of the material budget	0.03
Nonuniformity of the light collection	0.11
Total systematic uncertainty	0.18
Statistical uncertainty	0.18
Total uncertainty	0.26







# $m_H$ measurement in $H \rightarrow \gamma\gamma$

- Modelling

- Multiple gaussian for sig. pdf, data sideband for bkg.
- Maximum likelihood,  $N \times PDF(m_{\gamma\gamma} | m_H) + BKG$
- Simultaneous fit on 7 categories




Final  $m_H$  in  $H \rightarrow \gamma\gamma$  combined with 2016  $H \rightarrow ZZ$  and CMS Run1 results

Only 2016  $H \rightarrow \gamma\gamma$   
 $m_H = 125.78 \pm 0.26(0.18) \text{ GeV}$

# $m_H$ measurement in $H \rightarrow ZZ$

- Newest result with full Run2 dataset (  $138\text{fb}^{-1}$ ,  $13\text{TeV}$  )
- Expected event yield  $\sim 600$  events with  $S/B \sim 1:1$
- Lower statistics with better momentum measurement ( w.r.t  $H \rightarrow \gamma\gamma$  )
  - Final  $m_H$  precision is determined by statistical uncertainty
  - More effort on reducing statistical part, detector resolution optimisation, sig-bkg separation, etc.






# $m_H$ measurement in $H \rightarrow ZZ$

- Observables

- Four-lepton invariant mass ( $m_{4\ell}$ )
- ME-based discriminant ( $D$ )
- Event by event mass error (ebe)

- Improvements on momentum resolution

- $Z_1$  mass constraint
  - A kinematic fit using the intermediate on-shell Z line-shape to calibrate leading lepton pair momentum
- Beam-spot constraint
  - Muon tracks are constrained to beam-spot by KM algorithm, inject more information provided by BS position to track reconstruction
  - Improve Higgs mass resolution by 5~8% in  $4\mu$  final state, smaller impacts for  $2e2\mu$  and  $2\mu2e$





# $m_H$ measurement in $H \rightarrow ZZ$

- Systematic uncertainties



Difference(%)	4 $\mu$	4e	2e2 $\mu$	2 $\mu$ 2e
Muon momentum scale	0.03%	-	0.03%	0.03%
Electron energy scale	-	0.15%	0.15%	0.15%
Muon momentum resolution	3%	-	3%	3%
Electron energy resolution	-	10%	10%	10%

- Modelling

- 9 categories based on per-event mass error
- $N_{ebe} \times PDF(m_{4\ell} | m_H) \times PDF(D | m_{4\ell}) + BKG$
- Combination with CMS Run1  $H \rightarrow ZZ$

$$m_H = 125.08 \pm 0.12 (\pm 0.10) \text{ GeV}$$


Expected: 0.12 GeV



# $\Gamma_H$ measurement in $H \rightarrow ZZ$

- On-shell

- The same procedure as  $m_H$  measurement, treat  $\Gamma_H$  as POI and signal strength and  $m_H$  as free parameters
- Bounded parameter, Feldman-Cousins algorithm implemented




- Off-shell

- Aim to measure width from the ratio of on-shell and off-shell yields

$$\frac{\sigma_{gg \rightarrow H \rightarrow ZZ^*}^{on-shell}}{\sigma_{gg \rightarrow H^* \rightarrow ZZ}^{off-shell}} \sim \frac{g_{ggH}^2 g_{HZZ}^2}{m_H \Gamma_H}$$

$$\sim \frac{g_{ggH}^2 g_{HZZ}^2}{(2m_Z)^2}$$



# Summary

- Higgs mass measurement has been introduced from aspects of achievable precision, methods, advantages/drawbacks of the two golden channels
  - Statistical uncertainty dominates  $H \rightarrow ZZ$ . How to improve mass resolution and efficiency
  - Systematic uncertainty dominates  $H \rightarrow \gamma\gamma$ . How to calibrate ECAL energy reconstruction
- Cross reference from ATLAS

LHC full Run2	ATLAS	CMS
$H \rightarrow ZZ$	$124.99 \pm 0.19 \text{ GeV}$	$125.04 \pm 0.12 \text{ GeV}$
$H \rightarrow \gamma\gamma$	$125.17 \pm 0.14 \text{ GeV}$	-

- $\Gamma_H$  from  $H \rightarrow ZZ$ 
  - Off-shell:  $\Gamma_H < 2.9^{+2.3}_{-1.7} \text{ MeV}$
  - On-shell:  $\Gamma_H < 0.05 \text{ GeV}$

