

# Radiation Semiconductor Detector


史欣



2024年7月23日

RASER Workshop 2024

# Building Blocks of Elementary Particles



# Standard Model of FUNDAMENTAL PARTICLES AND INTERACTIONS

The Standard Model summarizes the current knowledge in Particle Physics. It is the quantum theory that includes the theory of strong interactions (quantum chromodynamics or QCD) and the unified theory of weak and electromagnetic interactions (electroweak). Gravity is included on this chart because it is one of the fundamental interactions even though not part of the "Standard Model."

## FERMIONS

**matter constituents**  
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_e$ electron neutrino	$<1 \times 10^{-8}$	0	<b>u</b> up	0.003	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.006	-1/3
$\nu_\mu$ muon neutrino	$<0.0002$	0	<b>c</b> charm	1.3	2/3
<b><math>\mu</math></b> muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_\tau$ tau neutrino	$<0.02$	0	<b>t</b> top	175	2/3
<b><math>\tau</math></b> tau	1.7771	-1	<b>b</b> bottom	4.3	-1/3

**Spin** is the intrinsic angular momentum of particles. Spin is given in units of  $\hbar$ , which is the quantum unit of angular momentum, where  $\hbar = h/2\pi = 6.58 \times 10^{-25}$  GeV s =  $1.05 \times 10^{-34}$  J s.

**Electric charges** are given in units of the proton's charge. In SI units the electric charge of the proton is  $1.60 \times 10^{-19}$  coulombs.

The **energy** unit of particle physics is the electronvolt (eV), the energy gained by one electron in crossing a potential difference of one volt. **Masses** are given in GeV/c<sup>2</sup> (remember  $E = mc^2$ ), where  $1 \text{ GeV} = 10^9 \text{ eV} = 1.60 \times 10^{-10}$  joule. The mass of the proton is  $0.938 \text{ GeV}/c^2 = 1.67 \times 10^{-27}$  kg.

## BOSONS

**force carriers**  
spin = 0, 1, 2, ...

Unified Electroweak spin = 1			Strong (color) spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge	Name	Mass GeV/c <sup>2</sup>	Electric charge
$\gamma$ photon	0	0	<b>g</b> gluon	0	0
<b>W<sup>-</sup></b>	80.4	-1			
<b>W<sup>+</sup></b>	80.4	+1			
<b>Z<sup>0</sup></b>	91.187	0			

### Color Charge


Each quark carries one of three types of "strong charge," also called "color charge." These charges have nothing to do with the colors of visible light. There are eight possible types of color charge for gluons. Just as electrically-charged particles interact by exchanging photons, in strong interactions color-charged particles interact by exchanging gluons. Leptons, photons, and **W** and **Z** bosons have no strong interactions and hence no color charge.

### Quarks Confined in Mesons and Baryons

One cannot isolate quarks and gluons; they are confined in color-neutral particles called **hadrons**. This confinement (binding) results from multiple exchanges of gluons among the color-charged constituents. As color-charged particles (quarks and gluons) move apart, the energy in the color-force field between them increases. This energy eventually is converted into additional quark-antiquark pairs (see figure below). The quarks and antiquarks then combine into hadrons; these are the particles seen to emerge. Two types of hadrons have been observed in nature: **mesons**  $q\bar{q}$  and **baryons**  $qqq$ .

### Residual Strong Interaction

The strong binding of color-neutral protons and neutrons to form nuclei is due to residual strong interactions between their color-charged constituents. It is similar to the residual electrical interaction that binds electrically neutral atoms to form molecules. It can also be viewed as the exchange of mesons between the hadrons.



If the protons and neutrons in this picture were 10 cm across, then the quarks and electrons would be less than 0.1 mm in size and the entire atom would be about 10 km across.

## PROPERTIES OF THE INTERACTIONS

Baryons $qqq$ and Antibaryons $\bar{q}\bar{q}\bar{q}$					
Baryons are fermionic hadrons. There are about 120 types of baryons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c <sup>2</sup>	Spin
<b>p</b>	proton	<b>uud</b>	1	0.938	1/2
$\bar{p}$	anti-proton	$\bar{u}\bar{u}\bar{d}$	-1	0.938	1/2
<b>n</b>	neutron	<b>udd</b>	0	0.940	1/2
$\Lambda$	lambda	<b>uds</b>	0	1.116	1/2
$\Omega^-$	omega	<b>sss</b>	-1	1.672	3/2

### Matter and Antimatter




For every particle type there is a corresponding antiparticle type, denoted by a bar over the particle symbol (unless + or - charge is shown). Particle and antiparticle have identical mass and spin but opposite charges. Some electrically neutral bosons (e.g.,  $Z^0$ ,  $\gamma$ , and  $\eta_c = c\bar{c}$ , but not  $K^0 = d\bar{s}$ ) are their own antiparticles.

### Figures

These diagrams are an artist's conception of physical processes. They are not exact and have no meaningful scale. Green shaded areas represent the cloud of gluons or the gluon field, and red lines the quark paths.

Property	Gravitational		Weak (Electroweak)		Electromagnetic		Strong	
	Mass - Energy	Flavor	Electric Charge	Color Charge	See Residual Strong Interaction Note	Fundamental	Residual	
Acts on:	All	Quarks, Leptons	Electrically charged	Quarks, Gluons				
Particles experiencing:	All	Quarks, Leptons	Electrically charged	Quarks, Gluons				
Particles mediating:	Graviton (not yet observed)	<b>W<sup>+</sup> W<sup>-</sup> Z<sup>0</sup></b>	$\gamma$	Gluons				
Strength relative to electromag for two u quarks at:	$10^{-41}$ $10^{-41}$ $10^{-36}$	0.8 $10^{-4}$ $10^{-7}$	1 1 1	25 60 Not applicable to hadrons			Not applicable to quarks 20	

Mesons $q\bar{q}$					
Mesons are bosonic hadrons. There are about 140 types of mesons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c <sup>2</sup>	Spin
$\pi^+$	pion	<b>u<math>\bar{d}</math></b>	+1	0.140	0
<b>K<sup>-</sup></b>	kaon	<b>s<math>\bar{u}</math></b>	-1	0.494	0
$\rho^+$	rho	<b>u<math>\bar{d}</math></b>	+1	0.770	1
<b>B<sup>0</sup></b>	B-zero	<b>d<math>\bar{b}</math></b>	0	5.279	0
$\eta_c$	eta-c	<b>c<math>\bar{c}</math></b>	0	2.980	0



### The Particle Adventure

Visit the award-winning web feature *The Particle Adventure* at <http://ParticleAdventure.org>

This chart has been made possible by the generous support of:


U.S. Department of Energy  
U.S. National Science Foundation  
Lawrence Berkeley National Laboratory  
Stanford Linear Accelerator Center  
American Physical Society, Division of Particles and Fields  
**BURLE INDUSTRIES, INC.**

©2000 Contemporary Physics Education Project. CPEP is a non-profit organization of teachers, physicists, and educators. Send mail to: CPEP, MS 50-308, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720. For information on charts, text materials, hands-on classroom activities, and workshops, see:

<http://CPEPweb.org>

# Particle accelerator

- Device produces a beam of fast-moving, electrically charged atomic or subatomic particles





© 2012 Encyclopædia Britannica, Inc.

# Particle detector


Year	Name	Detector principle	Discovery	Nobel Prize
1896	H. Becquerel	photographic plate	radioactivity	1903
1908	H. Geiger	gas amplification		
1911	E. Rutherford	scintillation screen	atomic nucleus	
1912	C.T.R. Wilson	cloud chamber	many new particles	1927
1912	V. Hess	electrometer	cosmic rays	1936
1924	W. Bothe	coincidence method		1954
1933	P. Blackett	triggered cloud chamber	$e^+e^-$ pairs	1948
1934	P.A. Cherenkov	Cherenkov radiation	$\nu$ oscillation	1958
1947	C.F. Powell	photoemulsion	pion	1950
1953	D.A. Glaser	bubble chamber	$\Omega^-$ , neutral currents	1960
1968	G. Charpak	multiwire prop. chamber		1992
1980		Si microstrip detector	$B\bar{B}$ oscillation	

# Historical Particle Detectors


Bubble chamber



Cloud chamber




# Large particle detector apparatus



# Particle detection by semiconductor detectors


- charged particle loses energy by ionisation or a photon absorbed in semiconductor part of released energy used to generate electron-hole pairs
- charge carrier pairs separated in electric field applied to the semiconductor and drift inside the bulk toward the electrodes
- electronic signal determined in size and shape
  - generated of e/h pairs
  - velocity of their drift movement
  - electrode geometry
- drift velocity depends on carrier mobility  $\mu$  and the magnitude of the electric field






# Single-sided silicon detectors

- single-sided pro
  - simpler and cost effective
  - good for large area detectors



(a) pn area diode.




(d) Strip detector.


Kolanoski, Wermes 2015

# Two-dimensional position information


- 2D position with crossed strip layers
  - cons: material thickness doubles
  - increased multiple scattering and secondary interactions
    - double-sided strip
- small stereo angle allows several modules aligned as a 'ladder'



# Ambiguities in crossed strip layers – ‘ghosts’





(a) Orthogonal strips.




(b) Small angle crossing.


# Double-sided microstrip detectors



# Silicon strip detectors as vertex detectors




(a) Fixed-target experiment (HERA-B).




(b) Collider experiment (H1).

# Hybrid pixel detector




(a) Hybrid pixel cell.



(b) Pixel matrix.

# Monolithic CMOS pixel detectors

- Monolithic active pixel sensors (MAPS)



- deposited charge is very small ( $\sim 1500e$ ), slow readout

# Acknowledgement

- Particle Detectors, 1st Edition, Edited by H. Kolanoski and N. Wermes