

Discovery of a Glueball-like Particle $X(2370)$ from J Decays

Shan JIN

**for the BESIII Collaboration
Nanjing University**


50 Years Discovery of the J Particle
Beijing , October 20, 2024

New forms of hadrons


- Hadrons consist of 2 or 3 quarks :

Naive Quark Model:

Meson ($q \bar{q}$)



Baryon ($q q q$)



- **New forms of hadrons:**


- Multi-quark states : Number of quarks ≥ 4
- Hybrids : $q\bar{q}g$, $qqqg$...
- Glueballs : gg , ggg ...

Lots of candidates, but new forms of hadrons have not been established yet!

Glueballs


- Unique particles formed by gauge bosons due to non-Abel self-interactions
 - Unique kind of matter made of pure force (usually matter formed by fermions)
 - Glueballs to QCD is just as important as Higgs boson to EW.
- Direct test of QCD and SM
- Lattice QCD predictions
 - 0^{++} ground state: $1.3 \sim 2 \text{ GeV}/c^2$
 - 2^{++} ground state: $2.2 \sim 2.8 \text{ GeV}/c^2$
 - 0^{-+} ground state: $2.3 \sim 2.8 \text{ GeV}/c^2$

arxiv:2305.04869



J particle radiative decays

- Gluon rich environment



- Isospin filter
 - final states dominated by $I=0$ processes
- Spin-parity filter
 - C parity must be +, so $J^{PC} = 0^{-+}, 0^{++}, 1^{++}, 2^{++}, 2^{-+} \dots$



→ Clean environment

very different from proton-antiproton collision processes

→ Ideal place to search for glueballs

Glueball productions in J radiative decays

- Rich production in J radiative decays — glueball production rate in J radiative decays could be higher than normal hadrons.



→ J particle radiative decays are believed to be an ideal/golden place to search for glueballs.

Glueball decays

- via gluons
 - flavor symmetric decays
- No rigorous predictions on decay patterns and their branching ratios.
- The glueball decays should have similar decay patterns to the charmonium families since they all decay via gluons.
 - e.g. the 0^{-+} glueball should have similar decays to η_c
 - One of the largest decay modes of η_c is $\eta'\pi^+\pi^-$, so $J/\psi \rightarrow \gamma\eta'\pi^+\pi^-$ could be a good place to search for the 0^{-+} glueball.

Golden decay modes for 0^{-+} Glueball searches

- Typically, PPP (3 pseudoscalar mesons, such as $\pi\pi\eta$, $\pi\pi\eta'$, $KK\pi$) modes are believed as golden decay modes in 0^{-+} glueball searches.
 - S wave decays for 0^{-+} mesons, no suppression factor, major decay modes
 - PPP modes are either forbidden or strongly suppressed in 0^{++} , 2^{++} mesons decays — spin-parity filter
- PP (2 pseudoscalar mesons) modes are mostly forbidden for 0^{-+} mesons
- VV modes (2 vector mesons, such as $\omega\omega$, $\phi\phi$, $\rho\rho$, K^*K^*)
 - P wave decays for 0^{-+} mesons — suppressed decays, especially near mass threshold
 - All J^{PC} mesons allowed, not a spin-parity filter
- Baryon modes
 - All J^{PC} mesons allowed, not a spin-parity filter
- Multi-pion modes
 - All J^{PC} mesons allowed, not a spin-parity filter
 - 0^{-+} mesons decay mainly via 2 body sequential decays, i.e., mainly via $f_2(1270)$, $a_1(1260)$ pair intermediate states — 0^{-+} glueball mass may not be high enough, i.e., PS is not allowed.

Historical Glueball Candidates

- 0^{++} scalar glueball candidates
 - $f_0(1710)$
 - Discovered by MarkII in 1980's as $\theta_2(1720)$ at beginning, only based on a fit to the angular distribution. Lots of studies at MarkII, DM2, BESII.
 - J^{PC} was firstly changed to 0^{++} at BESII based on a full PWA in $J/\psi \rightarrow \gamma KK$
 - With PWA, it was carefully studied at BESIII in $J/\psi \rightarrow \gamma KK, \gamma\pi\pi, \gamma\eta\eta, \gamma\eta\eta'$.
 - **The high production rate of $J/\psi \rightarrow \gamma f_0(1710)$ and the suppression of $f_0(1710) \rightarrow \gamma\eta\eta'$ measured from BESIII strongly support its interpretation as a scalar glueball or it has large glueball content if it is a mixture of glueball and normal meson.**
 - **Difficulty:** With phase space subtracted, from BESII and BESIII measurements, we obtained $\Gamma(f_0(1710) \rightarrow$

- $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma\pi\pi) = (4.01 \pm 1.0) \times 10^{-4}$
BESII: PLB 642 441 (2006)
- $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma K_s K_s) = (2.00_{-0.02}^{+0.03+0.31}) \times 10^{-4}$
BESIII: PRD 98 072003 (2018)
- $B(f_0(1500) \rightarrow \eta\eta')/B(f_0(1500) \rightarrow \pi\pi) = (1.66_{-0.40}^{+0.42}) \times 10^{-1}$
- $B(f_0(1710) \rightarrow \eta\eta')/B(f_0(1710) \rightarrow \pi\pi) < (2.9_{-0.9}^{+1.1}) \times 10^{-3}$
BESIII: PRD 106 072012(2022)

Historical Glueball Candidates

- 0^{++} scalar glueball candidates
 - $f_0(1500)$
 - Discovered by Crystal Barrel in 1990's as a unique 0^{++} candidate since $f_0(1710)$ was f_2 at that time.
 - Difficulty: compared with $f_0(1710)$, much lower production rate than $f_0(1710)$ disfavors its interpretation as a scalar glueball.
 - Mixing between $f_0(1500)/f_0(1710)$, or even with $f_0(1790)$?
 - Difficulty: dynamic mixing mechanism needs to be understood from the first principle of QCD (not just phenomenological understanding).

BESIII:

$$B(J/\psi \rightarrow \gamma f_0(1500)) \sim 0.29 \times 10^{-3}$$

$$B(J/\psi \rightarrow \gamma f_0(1710)) \sim 2.2 \times 10^{-3}$$

Historical Glueball Candidates


• 2^{++} Tensor Glueball Candidates

• $\xi(2230)$

- First observed by MarkIII is $J/\psi \rightarrow \gamma KK$ in 1980's, then by BES I in 1990's in $J/\psi \rightarrow \gamma KK, \gamma\pi\pi, \gamma p\bar{p}$ with very narrow mass peak.
- It was a tensor glueball candidate due to good flavor symmetric decay property.
- **Difficulty: it was not confirmed by BESII, nor BESIII.**

• $f_2(2340)$

- Many wide f_2 mesons in the mass region of 2.3 GeV from the LQCD predictions
- **Difficulty: no clear mass peak of these f_2 mesons can be directly observed in J/ψ radiative decays due to large overlaps among various wide resonances. They can only be seen as PWA components.**
- $f_2(2340)$: its large production rate in J/ψ radiative decays favors its interpretation as a tensor glueball candidate.
- **More PWA studies are needed to check the consistency among various decays modes. However, due to large overlaps again, no independent mass and width scan can be performed in PWA, i.e., the masses and widths of these f_2 mesons have to be fixed to previous measurements.**



Resonance	M (MeV/c ²)	Γ (MeV/c ²)	B.F. ($\times 10^{-4}$)	Sig.
$\eta(2225)$	2216^{+4+21}_{-5-11}	185^{+12+43}_{-14-17}	$(2.40 \pm 0.10^{+2.47}_{-0.18})$	28σ
$\eta(2100)$	2050^{+30+75}_{-24-26}	$250^{+36+181}_{-30-164}$	$(3.30 \pm 0.09^{+0.18}_{-3.04})$	22σ
$X(2500)$	$2470^{+15+101}_{-19-23}$	230^{+64+56}_{-35-33}	$(0.17 \pm 0.02^{+0.02}_{-0.08})$	8.8σ
$f_0(2100)$	2101	224	$(0.43 \pm 0.04^{+0.24}_{-0.03})$	24σ
$f_2(2010)$	2011	202	$(0.35 \pm 0.05^{+0.28}_{-0.15})$	9.5σ
$f_2(2300)$	2297	149	$(0.44 \pm 0.07^{+0.09}_{-0.13})$	6.4σ
$f_2(2340)$	2339	319	$(1.91 \pm 0.14^{+0.72}_{-0.73})$	11σ
0^{-+} PHSP			$(2.74 \pm 0.15^{+0.16}_{-1.48})$	6.8σ

Historical Glueball Candidates

- 0^{-+} Pseudoscalar Glueball Candidate
 - $\eta(1405)$ first discovered by MarkII in 1980's, named as $\iota(1440)$ with complicated structures. Lots of studies at MarkII, MarkIII, DM2 and BES.
 - Believed as the first glueball candidate due to its large production rate in J/ψ radiative decays and lack of reliable LQCD predictions in 1980's
 - No longer to be believed as 0^{-+} glueball candidate due to its large different mass from LQCD prediction.


BESIII Data samples

Totally about 50fb^{-1} integrated luminosity

Data sets collected so far include

- ◆ 10×10^9 J/ψ events
- ◆ 2.7×10^9 $\psi(2S)$ events
- ◆ 20fb^{-1} $\psi(3770)$
- ◆ Scan data between 2.0 and 3.08 GeV, and above 3.74 GeV
- ◆ Large datasets for XYZ studies:

Scan with $>500\text{pb}^{-1}$ per energy point space 10–20 MeV apart



World largest J/ψ data sample : ~10 billion

Key scientific question to be answered at BESIII: Glueballs exist or not?

- Searching for glueballs has been the top physics goal of BEPC and BEPCII for about 40 years!
- With 10 billion J/ψ data, we should be able to answer this key question — Either we find them, or exclude them.
 - [BESIII Physics Yellow Book](#)

→ We started to find the answer...





X(2370)

- Discovered by BESIII in $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$ in 2011.

Resonance	M (MeV/c ²)	Γ (MeV/c ²)	Stat.Sig.
X(1835)	$1836.5 \pm 3.0^{+5.6}_{-2.1}$	$190.1 \pm 9.0^{+38}_{-36}$	$>20\sigma$
X(2120)	$2122.4 \pm 6.7^{+4.7}_{-2.7}$	$83 \pm 16^{+31}_{-11}$	7.2σ
X(2370)	$2376.3 \pm 8.7^{+3.2}_{-4.3}$	$83 \pm 17^{+44}_{-6}$	6.4σ

- Confirmed by BESIII in $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$, $\gamma K K \eta'$ (new mode)

PRL106, 072002 (2011)




PRL 117, 042002 (2016)

X(2370) — good candidate of 0^{-+} glueball when it was first observed at beginning

- Its mass is consistent with LQCD prediction on the 0^{-+} glueball.
- Observed in the best place to search for the 0^{-+} glueball:
 - In J/ψ radiative decays
 - In golden decays modes of 0^{-+} glueball searches: first in $\pi\pi\eta'$ & later in $KK\eta'$
- Flavor symmetric decay — observed in both $\pi^+\pi^-\eta'$ and $KK\eta'$ decay modes.

→ Determination of its spin-parity is crucial!

Spin-parity determination of $X(2370)$ in $J/\psi \rightarrow \gamma K_s K_s \eta'$



- Almost background free channel
- 10 billion J/ψ data
- Very good BESIII detector performance

$J^{PC} = 0^{++}$ with significance $>9.8\sigma$

$M = 2395 \pm 11^{+26}_{-94}$ MeV

$\Gamma = 188^{+18}_{-17} {}^{+124}_{-33}$ MeV

$B(J/\psi \rightarrow \gamma X(2370)) B(X(2370) \rightarrow f_0(980) \eta') B(f_0(980) \rightarrow K_s^0 K_s^0)$
 $= (1.31 \pm 0.22 {}^{+2.85}_{-0.84}) \times 10^{-5}$

Compared with LQCD prediction on lightest 0^{-+} Glueball

$X(2370)$ measurements: [PRL 132 \(2024\) 181901](#)

$J^{PC} = 0^{-+}$ with significance $>9.8\sigma$

$$M = 2395 \pm 11^{+26}_{-94} \text{ MeV}$$

$$\Gamma = 188^{+18}_{-17}{}^{+124}_{-33} \text{ MeV}$$

$$\begin{aligned} & \mathbf{B}(J/\psi \rightarrow \gamma X(2370)) \mathbf{B}(X(2370) \rightarrow f_0(980) \eta') \mathbf{B}(f_0(980) \rightarrow K_s^0 K_s^0) \\ & = (1.31 \pm 0.22^{+2.85}_{-0.84}) \times 10^{-5} \end{aligned}$$

LQCD prediction on lightest pseudoscalar glueball:

$J^{PC} = 0^{-+}$ [PRD 100 \(2019\) 054511](#)


$$M = 2395 \pm 14 \text{ MeV}$$

$$\mathbf{B}(J/\psi \rightarrow \gamma G_{0^{-+}}) = (2.31 \pm 0.80) \times 10^{-4}$$


- ◇ The measurements are in a agreement with the predictions on **lightest pseudoscalar glueball**
- ◆ The spin-parity of the $X(2370)$ is determined to be 0^{-+} for the first time
- ◆ Mass is in a good agreement with LQCD predictions
- ◆ The estimation on $\mathbf{B}(J/\psi \rightarrow \gamma X(2370))$ and prediction on $\mathbf{B}(J/\psi \rightarrow \gamma G_{0^{-+}})$ are consistent within errors (assuming $\sim 5\%$ decay rate, $\mathbf{B}(J/\psi \rightarrow \gamma X(2370)) = (10.7^{+22.8}_{-7}) \times 10^{-4}$)

New (preliminary) results on X(2370) at ICHEP2024

Observation of new decay mode: $X(2370) \rightarrow K_S^0 K_S^0 \pi^0$




- Almost background free channel
- 1D mass spectrum fit
- Statistical significance : $\gg 5\sigma$
- Mass and width



$$M_{X(2370)} = 2321 \pm 4(\text{stat}) \pm 65(\text{syst.}) \text{ MeV}$$

$$\Gamma_{X(2370)} = 182 \pm 16(\text{stat}) \pm 59(\text{syst.}) \text{ MeV}$$


Observation of new decay mode: $X(2370) \rightarrow \pi^0\pi^0\eta$




- Almost background free channel
- 1D mass spectrum fit
- Statistical significance : $\gg 5\sigma$
- Mass and width

$$M_{X(2370)} = 2370 \pm 2(stat) \pm 52(syst.) \text{ MeV}$$


$$\Gamma_{X(2370)} = 134 \pm 8(stat) \pm 30(syst.) \text{ MeV}$$



Observation of new decay mode: $X(2370) \rightarrow a_0(980)\pi^0$



- Clear $a_0(980)$ signal in $M_{\pi\eta}$ spectrum
- 1D mass spectrum fit
- Statistical significance : $\gg 5\sigma$
- Mass and width



$$M_{X(2370)} = 2352 \pm 3(\text{stat}) \pm 74(\text{syst.}) \text{ MeV}$$

$$\Gamma_{X(2370)} = 134 \pm 4(\text{stat}) \pm 62(\text{syst.}) \text{ MeV}$$


$a_0(980)$ signal region
 $|m_{\pi^0\eta} - 0.98| < 0.05 \text{ GeV}$

$$J/\psi \rightarrow \gamma K_S K_S \eta$$

In the 2D mass plot of M_{KK} vs. $M_{KK\eta}$ in the BESIII paper on the spin-parity determination of the X(1835), qualitatively, we can clearly observe:

- In the upper M_{KK} mass band of 1.5-1.7 GeV range, clear signals of both X(2370) and η_c .
- In the lower M_{KK} mass band of $f_0(980)$, no X(2370), nor η_c .

PRL 115, 091803 (2015)



5 major η_c decay modes (from PDG)
 — 5 “Golden” modes in 0^{-+} glueball traditional searches

Decays involving hadronic resonances

Γ_1	$\eta'(958) \pi \pi$	(1.87 ± 0.26) %
Γ_2	$\eta'(958) K \bar{K}$	(1.61 ± 0.25) %

Decays into stable hadrons

Γ_{34}	$K \bar{K} \pi$	(7.0 ± 0.4) %
Γ_{35}	$K \bar{K} \eta$	(1.32 ± 0.15) %
Γ_{36}	$\eta \pi^+ \pi^-$	(1.7 ± 0.5) %

X(2370) is observed in all 5 modes

→ High similarities between X(2370) and η_c

Discussion on $X(2370)$ properties

- With personal understanding

Discussion on $X(2370)$ properties — Decays

- Normal $q\bar{q}$ mesons, hybrids and multiquark states can hardly explain all 5 decay modes ($\eta'\pi\pi$, $\eta'KK$, $\eta\pi\pi$, ηKK , $KK\pi$) with different quark flavor combinations.
 - e.g., Strong mixing between $u\bar{u}+d\bar{d}$ and $s\bar{s}$ would be needed for a normal $q\bar{q}$ meson to have all these 5 decay modes, however, LQCD calculation (PRD 83 (2011) 11502) shows that the mixing between $u\bar{u}+d\bar{d}$ and $s\bar{s}$ components should be very small for a 0^- $q\bar{q}$ meson around 2GeV.
- The high similarities between $X(2370)$ and η_c decay modes strongly suggest it decays via gluons.
- Narrow decay partial widths → next page

Narrow decay partial widths of X(2370)

- For 5 golden PPP decay modes: similar number of events under the X(2370) peak — No dominant decay modes, similar to η_c !
- Naïve estimation on the BR of each mode ~ 5 -10%, i.e., partial width of each decay mode is ~ 10 MeV!
- This would be very hard to be explained if there were quark content (qqbar, qqg, or multiquark) in X(2370) for OZI allowed decays:
 - Typical OZI allowed decay partial width ~ 100 MeV (see all PDG mesons)
 - OZI allowed decays usually have dominant decay modes
- X(2370) decay should be OZI suppressed decays as η_c , i.e., via gluons!

Discussion on X(2370) properties — Production

- Richly produced in J radiative decays — just as glueball expectation
 - In the above 2.3 GeV mass region as LQCD 0-+ glueball prediction, X(2370) is the **unique** 0-+ particle produced in these “5 golden modes” and in J/ψ radiative decays.
 - No other qualified candidates for 0-+ glueball since their production rates should be very low in J radiative decays if they have not shown up in 10 billion such a huge J data sample.
- The unique qualified 0-+ glueball candidate in 10 billion J data make us face a situation: Either X(2370) is the lightest 0-+ glueball, or LQCD may meet a big challenge in the glueball predictions.
- similar to the situation before the Higgs boson was discovered.

Conclusions

- BESIII discovers a glueball like particle X(2370) from J decays. The mass, spin-parity quantum numbers, production and decay properties of X(2370) are consistent with the features of the lightest pseudoscalar glueball.
- The J particle and its family are extremely important in the discovery of glueballs, both in the glueball production and in understanding the glueball decays.

Conclusions

- BESIII discovers a glueball like particle $X(2370)$ from J decays. The mass, spin-parity quantum numbers, production and decay properties of $X(2370)$ are consistent with the features of the lightest pseudoscalar glueball.
- The J particle and its family are extremely important in the discovery of glueballs, both in the glueball production and in understanding the glueball decays.


Thanks!

backup





Observation of $X(1835)$ in $J/\psi \rightarrow \gamma K_S K_S \eta$

- Use 1.3×10^9 J/ψ events collected by BESIII in 2009 and 2012
- Clear structure on mass spectrum of $K_S K_S \eta$ around $1.85 \text{ GeV}/c^2$
- Strongly correlated to $f_0(980)$
- PWA for $M(K_S K_S) < 1.1 \text{ GeV}/c^2$

PRL 115, 091803 (2015)



Confirmation of the X(2370) in $J/\psi \rightarrow \gamma K K \eta'$



EPJC (2020) 80:746

◆ Combination with 1.31×10^9 J/ψ events

- $J/\psi \rightarrow \gamma K^+ K^- \eta'$ and $J/\psi \rightarrow \gamma K_s K_{s\eta}'$
- $\eta' \rightarrow \gamma \pi \pi$ and $\eta' \rightarrow \pi \pi \eta$


◆ Confirmation of the X(2370) with 8.3σ

- $M = 2341.6 \pm 6.5(\text{stat.}) \pm 5.7(\text{syst.}) \text{ MeV}$
- $\Gamma = 117 \pm 10(\text{stat.}) \pm 8(\text{syst.}) \text{ MeV}$
- $\text{Br}(J/\psi \rightarrow \gamma X(2370) \rightarrow \gamma K^+ K^- \eta') = (1.79 \pm 0.23 \pm 0.65) \times 10^{-5}$
- $\text{Br}(J/\psi \rightarrow \gamma X(2370) \rightarrow \gamma K_s K_{s\eta}') = (1.18 \pm 0.32 \pm 0.39) \times 10^{-5}$


Observation: X(2370) new decay mode of $K K \eta'$

$\pi^0\pi^0\eta$ invariant mass spectrum in $a_0(980)$ signal region


- After requiring $|m_{\pi^0\pi^0} - 1.5| > 0.15$ GeV to veto $f_0(1500)$ signal, there is the clear $X(2370)$ signal in the $a_0(980)$ signal region, but not in the $a_0(980)$ sideband region.



$a_0(980)$ signal region
 $|m_{\pi^0\eta} - 0.98| < 0.05$ GeV



$a_0(980)$ sideband region
 $0.2 < |m_{\pi^0\eta} - 0.98| < 0.25$ GeV




$M(\pi^0\pi^0\eta) > 2$ GeV




$$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$$

- $\pi^+ \pi^- \eta'$ is the first mode to discover the X(2370)
- In the 2D plot of $M_{\pi\pi}$ vs. $M_{\pi\pi\eta'}$, in the $f_0(980)$ mass band, clear signals of both X(2370) and η_c
- Additional important information:
 - $X(2370) \rightarrow \pi\pi\eta'$ almost 100% via $f_0(980)\eta'$
 - $\eta_c \rightarrow \pi\pi\eta'$ dominantly via $X(2000)\eta'$
 - NO phase space for $X(2370) \rightarrow X(2000)\eta'$









$$J/\psi \rightarrow \gamma K_s K_s \eta'$$

- In the $f_0(980)$ mass band of $K_s K_s$ mass, clear signals of both $X(2370)$ and η_c
- Additional important information:
 - $X(2370) \rightarrow K K \eta'$ almost 100% via $f_0(980) \eta'$
 - $\eta_c \rightarrow K K \eta'$ dominantly via $X(1800) \eta'$
 - NO phase space for $X(2370) \rightarrow X(1800) \eta'$



Observation of new decay modes of $X(2370)$



- $X(2370) \rightarrow K_S^0 K_S^0 \pi^0, \eta \pi^0 \pi^0, a_0^0(980) \pi^0$ firstly observed, all accompanied with η_c

* $\eta(2320) \rightarrow \eta \eta \eta, \eta \pi \pi$ [PL B496 145(2000)] could be the current $X(2370)$ at BESIII 36