

# Spin Correlations

Shu-Yi Wei 魏树一 (Shandong University)

shuyi@sdu.edu.cn

*H.C. Zhang, S.Y. Wei; PLB 839, 137821 (2023)*

*X.W. Li, Z.X. Chen, S. Cao, S.Y. Wei, PRD 109, 014035 (2024)*

*Z.X. Chen, H. Dong, S.Y. Wei, PRD 110, 056040 (2024)*

*L. Yang, Y.K. Song, S.Y. Wei, PRD 111, 054035 (2025)*

*F. Huang, T. Liu, Y.K. Song, S.Y. Wei, PLB 862, 139346 (2025)*



# Contents

- ☑ Introduction
- ☑ Spin correlation of back-to-back dihadron
- ☑ Helicity correlation of neighboring dihadron
- ☑ Summary

## QCD factorization

partonic interaction, perturbative

Cross Section = short distance  $\otimes$  long distance

non-perturbative, universal

**Fragmentation Functions:** quarks  $(k, \lambda_q, S_{T,q}) \rightarrow$  hadrons  $(p = zk, \lambda_h, S_{T,h})$

$$\mathcal{D}_q^h(z; \lambda_q, \lambda_h; S_{T,q}, S_{T,h}) = D_{1,q}^h(z) + \lambda_q \lambda_h G_{1L,q}^h(z) + S_{T,q} \cdot S_{T,h} H_{1T}(z)$$

$$\mathcal{D}_g^h(z; \lambda_g, \lambda_h; S_{T,h}) = D_{1,g}^h(z) + \lambda_g \lambda_h G_{1L,g}^h(z)$$

QCD dominant process: Parity Symmetry.

## QCD factorization

		Baryons		
		Unpolarized	L	T
Quarks	Unpolarized	$D_1$		
	L		$G_{1L}$	
	T			$H_{1T}$ quark only


☑  $G_{1L}$ , aka, the longitudinal spin transfer

Number density of longitudinally polarized hadrons produced from longitudinally polarized quarks.

polarized beams  
or  
weak interaction



## Single Inclusive $\Lambda$ Production in $e^+e^-$ Annihilation Experiment



Final state quarks gain polarization through weak interaction

spin transfer

$$\mathcal{P}_L^\Lambda = \lambda_q \frac{G_{1Lq}^\Lambda}{D_{1q}^\Lambda}$$

quark polarization

$$\frac{d\sigma}{dPS} = \sigma_0 \left[ D_{1q}^\Lambda(z) + \lambda_q \lambda_\Lambda G_{1Lq}^\Lambda(z) \right]$$


Belle  
Energy

LEP  
Energy

# Introduction

## Single Inclusive $\Lambda$ Production in $e^+e^-$ Annihilation Experiment


Final state quarks gain polarization through weak interaction



spin transfer

$$\mathcal{P}_L^\Lambda = \lambda_q \frac{G_{1Lq}^\Lambda}{D_{1q}^\Lambda}$$


quark polarization



**DSV: Phys.Rev.D 57 (1998) 5811**

**See also: Chen, Yang, Liang, Zhou, PRD 95, 034009 (2017)**

## Single Inclusive $\Lambda$ Production in $e^+e^-$ Annihilation Experiment



Final state quarks gain polarization through weak interaction

spin transfer

$$\mathcal{P}_L^\Lambda = \lambda_q \frac{G_{1Lq}^\Lambda}{D_{1q}^\Lambda}$$


quark polarization

$$\frac{d\sigma}{dPS} = \sigma_0 \left[ D_{1q}^\Lambda(z) + \lambda_q \lambda_\Lambda G_{1Lq}^\Lambda(z) \right]$$

Belle  
Energy

LEP  
Energy

## $\Lambda\bar{\Lambda}$ -pair Production in $e^+e^-$ Annihilation Experiment



Belle  
Energy

$$\frac{d\sigma}{dPS} = \sigma_0 \left[ D_{1q}^{\Lambda}(z_1) D_{1\bar{q}}^{\bar{\Lambda}}(z_2) - \lambda_{\Lambda} \lambda_{\bar{\Lambda}} G_{1Lq}^{\Lambda}(z_1) G_{1L\bar{q}}^{\bar{\Lambda}}(z_2) \right]$$

### ☑ Helicity Conservation

$q$  and  $\bar{q}$  are on the same fermion line.  
They must have opposite helicities.

### ☑ Helicity Correlation

A novel probe to the spin-dependent fragmentation functions

### ☑ Entangled states

$$\frac{1}{\sqrt{2}} |A^{\uparrow} B^{\downarrow} \pm A^{\downarrow} B^{\uparrow}\rangle$$

$$\frac{1}{\sqrt{2}} |A^{\uparrow} B^{\uparrow} \pm A^{\downarrow} B^{\downarrow}\rangle$$


*H.C. Zhang, SYW; PLB 839 (2023) 137821*  
see also Nucl. Phys. B 445 (1995) 380.



# Helicity correlation of B2B dihadron

## Helicity Correlation of $\Lambda\bar{\Lambda}$ -pair

$$C_{LL} = \frac{\text{same signs} - \text{opposite signs}}{\text{total cross section}} = - \frac{\sum_q \sigma_0 G_{1Lq}^\Lambda(z_1) G_{1L\bar{q}}^{\bar{\Lambda}}(z_2)}{\sum_q \sigma_0 D_{1q}^\Lambda(z_1) D_{1\bar{q}}^{\bar{\Lambda}}(z_2)} \propto \langle \cos \theta_1^* \cos \theta_2^* \rangle$$




☑ The helicity correlation at the Belle energy has a similar magnitude with that at the LEP energy.

☑ It is now possible to extract the longitudinal spin transfer at Belle experiment.

*H.C. Zhang, SYW; PLB 839 (2023) 137821*

# Helicity correlation of B2B dihadron

Applying to the unpolarized pp collisions




$$a + b \rightarrow c(\lambda_c) + d(\lambda_d)$$

☒ Are  $\lambda_c$  and  $\lambda_d$  correlated?

Yes!

“s-channel diagrams”: just like  $e^+e^-$  annihilation, maximum correlation



$$g + g \rightarrow q + \bar{q}$$

$$q_i + \bar{q}_i \rightarrow q_j + \bar{q}_j$$

$$q + \bar{q} \rightarrow g + g$$



negative correlation

*H.C. Zhang, SYW; PLB 839 (2023) 137821*



# Helicity correlation of B2B dihadron

“t-channel diagrams”: prefer same-sign correlation




$$\begin{aligned}
 &q + g \rightarrow q + g \\
 &q_i + q_j \rightarrow q_i + q_j \\
 &g + g \rightarrow g + g \\
 &q_i + q_i \rightarrow q_i + q_i \\
 &q_i + \bar{q}_i \rightarrow q_i + \bar{q}_i
 \end{aligned}$$

To summarize




- ☑ “s-channel”:  $\sigma_{+-} = \sigma_{-+} > \sigma_{++} = \sigma_{--} = 0$
- ☑ “t-channel”:  $\sigma_{++} = \sigma_{--} > \sigma_{+-} = \sigma_{-+} > 0$
- ☑ Probe polarized FF in unpolarized pp collisions
- ☑ Explore the circularly polarized gluon FF

*H.C. Zhang, SYW; PLB 839 (2023) 137821*



# Helicity correlation of B2B dihadron

## Polarization Correlation in unpolarized pp collisions




- ✓ Smaller, but none-zero
- ✓ Distinguish different scenarios
- ✓ Avoid contamination of polarized PDF
- ✓ Probe gluon spin transfer

*H.C. Zhang, SYW; PLB 839 (2023) 137821*



## Keywords of Jet Quenching




# Helicity correlation of B2B dihadron

## Helicity Correlation in central and peripheral AA collisions

A toy model:  $\left. \frac{d\sigma}{dPS} \right|_{AA} = \text{Energy Loss} \otimes \left. \frac{d\sigma}{dPS} \right|_{pp}$


*X. Li, Z.X. Chen, S. Cao, S.Y. Wei;*  
Phys.Rev.D 109, 014035 (2024)



Clear Enhancement in  
central AA collisions

☑ Much larger luminosity


☑ Jet Quenching + Polarization





# Helicity correlation of B2B dihadron

## Helicity Correlation in ultra-peripheral AA collisions




$\gamma + \gamma$

$\gamma + \mathbb{P}$


✓ Much larger luminosity

✓ Pomeron + Polarization




X. Li, Z.X. Chen, S. Cao, S.Y. Wei;  
Phys.Rev.D 109, 014035 (2024)

## Helicity Correlation at future EIC



Transverse  
photon




Z.X. Chen, H. Dong, S.Y. Wei;  
 PRD 110, 056040 (2024)



Longitudinal  
photon






## Helicity Correlation at future EIC



Z.X. Chen, H. Dong, S.Y. Wei;  
PRD 110, 056040 (2024)

## Helicity Correlation at future EIC



Z.X. Chen, H. Dong, S.Y. Wei;  
PRD 110, 056040 (2024)


QCD factorization

**Baryons**

		Unpolarized	L	T
Quarks	Unpolarized	$D_1$		
	L		$G_{1L}$	
	T			$H_{1T}$ quark only

☑  $H_{1T}$ , aka, the transverse spin transfer


Number density of transversely polarized hadrons produced from transversely polarized quarks.





# Transverse spin correlation of B2B dihadron

## Electron-positron annihilation




helicity correlation of  $q\bar{q}$ :

$$|M(\lambda_q = +1, \lambda_{\bar{q}} = -1)|^2 + |M(\lambda_q = -1, \lambda_{\bar{q}} = +1)|^2$$

transverse spin correlation of  $q\bar{q}$ :

$$2\text{Re} \left[ M(\lambda_q = +1, \lambda_{\bar{q}} = -1) M^*(\lambda_q = -1, \lambda_{\bar{q}} = +1) \right]$$



Helicity flip between amplitude and conjugate amplitude

Transverse spin correlation of  $q\bar{q}$  pair


Belle experiment

*L. Yang, Y.K. Song, S.Y. Wei;  
PRD111, 054035 (2025)*

# Transverse spin correlation of B2B dihadron

## Unpolarized pp Collisions



- $g + g \rightarrow q + \bar{q}$   
 $q_i + \bar{q}_i \rightarrow q_j + \bar{q}_j$   
 $q_i + \bar{q}_i \rightarrow q_i + \bar{q}_i$
- negative correlation
- $q_i + q_i \rightarrow q_i + q_i$
- small correlation
- $q_i + q_j \rightarrow q_i + q_j$
- no correlation
- $q + g \rightarrow q + g$   
 $q + \bar{q} \rightarrow g + g$   
 $g + g \rightarrow g + g$
- gluon channels does not contribute



*L. Yang, Y.K. Song, S.Y. Wei;  
PRD111, 054035 (2025)*

## Unpolarized pp Collisions

Partonic transverse spin correlation:  $\mathcal{R} = \frac{d\sigma^T/dt}{d\sigma^U/dt}$





*L. Yang, Y.K. Song, S.Y. Wei;*  
PRD111, 054035 (2025)



## Dihadron Fragmentation Function



$$\mathcal{D}^{h_1 h_2}(z_1, z_2, \lambda_1, \lambda_2, \mu_f^2) = D_1^{h_1 h_2}(z_1, z_2, \mu_f^2) + \lambda_1 \lambda_2 D_{1LL}^{h_1 h_2}(z_1, z_2, \mu_f^2) + \mathbf{S}_{T1} \cdot \mathbf{S}_{T2} D_{1TT}^{h_1 h_2}(z_1, z_2)$$



$$\hat{P}_{jk \leftarrow i}^{LL/U}(\xi) = \frac{1}{2} \sum_{\lambda_i} \left[ \hat{P}_{jk \leftarrow i}(\xi, \lambda_i, +, +) + \hat{P}_{jk \leftarrow i}(\xi, \lambda_i, -, -) - \hat{P}_{jk \leftarrow i}(\xi, \lambda_i, +, -) - \hat{P}_{jk \leftarrow i}(\xi, \lambda_i, -, +) \right]$$

*F. Huang, T. Liu, Y.K. Song, S.Y. Wei*  
PLB 862, 139346 (2025)

## Dihadron Fragmentation Function



*F. Huang, T. Liu, Y.K. Song, S.Y. Wei*  
PLB 862, 139346 (2025)

- ☑ Spin correlations emerge in unpolarized collisions.
- ☑ The unpolarized colliders are also capable of investigating the hadronization of polarized protons.

**Thanks for your attention!**



The End