Contribution ID: 29

Transverse momentum balance of dijets in Xe+Xe collisions at the LHC

Thursday, 21 November 2024 11:00 (20 minutes)

We present a theoretical study of the medium modifications of the $p_{\rm T}$ balance $(x_{\rm J})$ of dijets in Xe+Xe collisions at $\sqrt{s_{\rm NN}} = 5.44$ TeV. The initial production of dijets was carried out using the POWHEG+PYTHIA8 prescription, which matches the next-to-leading-order (NLO) QCD matrix elements with the parton shower (PS) effect. The SHELL model described the in-medium evolution of nucleus-nucleus collisions using a transport approach. The theoretical results of the dijet x_J in the Xe+Xe collisions exhibit more imbalanced distributions than those in the p+p collisions, consistent with recently reported ATLAS data. By utilizing the Interleaved Flavor Neutralisation, an infrared-and-collinear-safe jet flavor algorithm, to identify the flavor of the reconstructed jets, we classify dijets processes into three categories: gluon-gluon (gg), quark-gluon (qg), and quark-quark (qq), and investigated the respective medium modification patterns and fraction changes of the gg, qg, and qq components of the dijet sample in Xe+Xe collisions. It is shown that the increased fraction of qg component at a small x_J contributes to the imbalance of the dijet; in particular, the q_1g_2 (quark-jet-leading) dijets experience more significant asymmetric energy loss than the g_1q_2 (gluon-jet-leading) dijets traversing the QGP. By comparing the $\Delta \langle x_J \rangle = \langle x_J \rangle_{\text{PP}} - \langle x_J \rangle_{\text{AA}}$ of inclusive, $c\bar{c}$ and $b\bar{b}$ dijets in Xe+Xe collisions, we observe $\Delta \langle x_{\rm J} \rangle_{\rm incl.} > \Delta \langle x_{\rm J} \rangle_{\rm cc} > \Delta \langle x_{\rm J} \rangle_{\rm bb}$. Moreover, $\rho_{\rm Xe,Pb}$, the ratios of the nuclear modification factors of dijets in Xe+Xe to those in Pb+Pb, were calculated, which indicates that the yield suppression of dijets in Pb+Pb is more pronounced than that in Xe+Xe owing to the larger radius of the lead nucleus.

Primary authors: ZHANG, Ben-Wei (Central China Normal University); WANG, Sa (China Three Gorges University); Dr SHEN, Shu-Wan (CCNU); LI, Yao (Central China Normal University)

Presenter: LI, Yao (Central China Normal University)

Session Classification: Thursday Morning Second Session