

# Dynamical relaxation behaviors of a critical quench

*Wednesday, 20 November 2024 11:40 (20 minutes)*

We study the universal dynamical relaxation behaviors of a quantum XY chain following a quench, paying special attention to the case in which the initial state is a critical ground state, or the postquenched Hamiltonian is at a critical point of equilibrium quantum phase transition, or both of them are critical. In such a “critical quench,” we find very interesting real-time dynamical scaling behaviors and we find crossover phenomena between them. For a quench from a noncritical point to a critical point, we find that, compared to the noncritical quench, the universal power-law scaling behavior does not change; however, there may be a crossover between the exponential decaying behavior and the power-law scaling. For a quench from a critical point to a noncritical point, the power-law scaling behaviors  $t^{-3/2}$  and  $t^{-3/4}$  in the noncritical quenches may be changed to  $t^{-1}$  and  $t^{-1/2}$ , respectively. If the prequenched Hamiltonian is set to be a point that is close to but not exactly at a critical point, we find interesting crossover phenomena between different power-law scaling behaviors. We also study the quench from the vicinity of a multicritical point, we find crossover behaviors that are induced by a different mechanism, and we find another crossover exponent. All the results are related to the gap-closing properties of the energy spectrum of the critical points.

**Primary author:** 邹, 银涛 (华中师范大学)

**Presenter:** 邹, 银涛 (华中师范大学)

**Session Classification:** Wednesday Morning Second Session