

Next-to-leading order QCD corrections to $B_c^* \rightarrow J/\psi$ form factors

陶伟

河南师范大学

Based on

Qin Chang, Wei Tao, Zhen-Jun Xiao, Ruilin Zhu,
arXiv:2502.19829

2025年3月30日 粒子物理标准模型及新物理精细计算研讨会@保定

Outline

1

Introduction

2

Calculation Procedure

3

Results

4

Summary

B_c^* semileptonic decay and form factors

➤ $c\bar{b}$ meson

- the only meson containing two different heavy flavors
- $B_c(1S)$ discovered in 1998 through $B_c \rightarrow J/\psi + l\nu_l$
- $B_c^*(1S)$ not yet observed due to difficulty in detecting γ of $B_c^* \rightarrow B_c + \gamma$
- $B_c^*(1S)$ weak decays (e.g. $B_c^* \rightarrow J/\psi + l\nu_l$) help search for $B_c^*(1S)$

[CDF, PRL(1998)]
 [CMS, PRL(2019)]
 [LHCb, PRL(2019)]

➤ Form factors (FFs)

- Related to decay widths and branching ratios
- Lattice results available for $B_c \rightarrow J/\psi (\eta_c)$ (axial-)vector form factors, not for B_c^*
- $B_c^* \rightarrow J/\psi (\eta_c)$ form factors known only by light-front quark model (LFQM)

[HPQCD, 1611.01987]
 [HPQCD, PRD(2020)]

[Q.Chang,L.T.Wang,X.N.Li, JHEP(2019)]
 [Q.Chang, et al., AHEP(2020)]
 [S.Y.Wang, et al., CPC(2024)]

Why NRQCD higher-order calculations

[G.T.Bodwin,E.Braaten,G.P.Lepage,PRD(1995)]

➤ Non-Relativistic Quantum Chromodynamics (NRQCD)

- Form factor = short-distance coefficient \times wavefunction at origin
 - perturbative expansion in α_s
 - nonperturbative
 - To test perturbative expansion convergence and renormalization scale dependence in NRQCD
 - To obtain more precise theoretical predictions and test the Standard Model
 - To study new physics by calculating (axial-)tensor form factors

Review NRQCD calculations for $c\bar{b} \rightarrow c\bar{c}$ form factors

- 2007, first one-loop for $B_c \rightarrow \eta_c$ vector and tensor FFs
[G. Bell, 0705.3133]
- Since 2011, next-to-leading order (NLO) QCD corrections to $B_c \rightarrow J/\psi$ (η_c) (axial-)vector and (axial-)tensor FFs
[C.F.Qiao,P.Sun, JHEP(2012)]
[C.F.Qiao,R.Zhu, PRD(2013)]
[W.Tao,Z.J.Xiao,R.Zhu, PRD(2022)]
- From 2017 onward, relativistic corrections for B_c decaying into S(P)-wave charmonium FFs
[R.Zhu,et al., PRD(2017)]
[R.Zhu, NPB(2018)]
[D.Shen,et al., IJMPA(2021)]
- 2024, leading order (LO) for $B_c^* \rightarrow J/\psi$ (axial-)vector FFs
[Y.Geng,M.Cao,R.Zhu, PRD(2024)]

Outline

1

Introduction

2

Calculation Procedure

3

Results

4

Summary

Definition for $B_c^* \rightarrow J/\psi$ form factors

$$\begin{aligned} & \langle J/\psi(\epsilon', p') | \bar{b} \gamma_\mu c | B_c^*(\epsilon, p) \rangle \\ = & -(\epsilon \cdot \epsilon'^*) [P_\mu V_1(q^2) - q_\mu V_2(q^2)] - (\epsilon \cdot q) \epsilon'^*_\mu V_3(q^2) \\ & + (\epsilon'^* \cdot q) \epsilon_\mu V_4(q^2) + (\epsilon \cdot q) (\epsilon'^* \cdot q) \left[\left(\frac{P_\mu}{M^2 - M'^2} \right. \right. \\ & \left. \left. - \frac{q_\mu}{q^2} \right) V_5(q^2) + \frac{q_\mu}{q^2} V_6(q^2) \right], \end{aligned}$$

Independent FFs:
 $V_{1,2,3,4,5,6}, A_{1,2,3,4},$
 $T_{2,3,4,6}, T'_{2,3,4}$







$$\begin{aligned} & \langle J/\psi(\epsilon', p') | \bar{b} \sigma_{\mu\nu} q^\nu c | B_c^*(\epsilon, p) \rangle \\ = & -i(\epsilon \cdot \epsilon'^*) [P_\mu T_1(q^2) - q_\mu T_2(q^2)] (M + M') \\ & - i[(\epsilon \cdot q) \epsilon'^*_\mu T_3(q^2) - (\epsilon'^* \cdot q) \epsilon_\mu T_4(q^2)] (M + M') \\ & + i \frac{(\epsilon \cdot q) (\epsilon'^* \cdot q)}{M + M'} [P_\mu T_5(q^2) + q_\mu T_6(q^2)], \end{aligned}$$

$$\begin{aligned} & \langle J/\psi(\epsilon', p') | \bar{b} \gamma_\mu \gamma_5 c | B_c^*(\epsilon, p) \rangle \\ = & i \varepsilon_{\mu\nu\alpha\beta} \epsilon^\alpha \epsilon'^*\epsilon^\beta [P^\nu A_1(q^2) - q^\nu A_2(q^2)] \\ & + \frac{i \varepsilon_{\mu\nu\alpha\beta} P^\alpha q^\beta}{M^2 - M'^2} [\epsilon'^* \cdot q \epsilon^\nu A_3(q^2) - \epsilon \cdot q \epsilon'^*\epsilon^\nu A_4(q^2)] \\ & - \frac{i \varepsilon_{\rho\nu\alpha\beta} \epsilon^\alpha \epsilon'^*\epsilon^\beta P^\nu q^\rho}{M^2 - M'^2} [P_\mu A_5(q^2) - q_\mu A_6(q^2)]. \end{aligned}$$

$$\begin{aligned} & \langle J/\psi(\epsilon', p') | \bar{b} \sigma_{\mu\nu} \gamma_5 q^\nu c | B_c^*(\epsilon, p) \rangle \\ = & -\varepsilon_{\mu\nu\alpha\beta} \epsilon^\alpha \epsilon'^*\epsilon^\beta [P^\nu T'_1(q^2) + q^\nu T'_2(q^2)] (M + M') \\ & + \frac{\varepsilon_{\mu\nu\alpha\beta} P^\alpha q^\beta}{M + M'} [\epsilon'^* \cdot q \epsilon^\nu T'_3(q^2) + \epsilon \cdot q \epsilon'^*\epsilon^\nu T'_4(q^2)] \\ & - \frac{\varepsilon_{\rho\nu\alpha\beta} \epsilon^\alpha \epsilon'^*\epsilon^\beta P^\nu q^\rho}{M + M'} [P_\mu T'_5(q^2) - q_\mu T'_6(q^2)]. \end{aligned}$$

- $P = p + p'$
- $q = p - p'$: transfer momentum

Step 1: generate Feynman diagrams & amplitudes



Step 2: amplitude simplification

- Dirac matrix simplification, index contraction, color algebra simplification, and trace calculation
- γ_5 scheme for the trace of a fermion chain containing γ_5
 - Naïve γ_5 scheme when containing 0/2 γ_5 [\[V.Shtabovenko,R.Mertig,F.Orellana, CPC\(2025\)\]](#)

$$\gamma_5 \gamma_\mu + \gamma_\mu \gamma_5 = 0, \gamma_5^2 = 1 \text{ and cyclicity}$$

- Fixed reading point γ_5 scheme when containing one γ_5
 - the fermion chain contains current vertex $\Gamma = \gamma_\mu \gamma_5$ or $\sigma_{\mu\nu} \gamma_5$

$$\text{Trace}(a \cdot \Gamma \cdot b) \rightarrow \text{Trace}\left(\frac{\Gamma \cdot b \cdot a + b \cdot a \cdot \Gamma}{2}\right)$$

- otherwise

[\[J.G.Korner,D.Kreimer,K.Schilcher, ZPC\(1992\)\]](#)
[\[S.A.Larin, PLB\(1993\)\]](#)
[\[S.Moch,J.A.M.Vermaseren,A.Vogt, PLB\(2015\)\]](#)

$$\text{Trace}(a \cdot \gamma_5 \cdot b) \rightarrow \text{Trace}(b \cdot a \cdot \gamma_5)$$

Step 3: Express amplitudes as $A_0, B_0, C_{0,1}, D_0$ & calculate them

amplitudes

Consisting of scalar products
of momenta

hierarchical heavy quark limit

FeynCalc TID

$A_0, B_0, C_{0,1}, D_0, E_0$

IBP reduction

Package-X

[K.G.Chetyrkin,F.V.Tkachov, NPB(1981)]
[H.H.Patel, CPC(2017)]

Series: expanding in small m_c and
taking the leading-order terms

PolyLog, Log, Sqrt

Step 4: Renormalization

- One-loop diagrams
- Tree diagrams inserted with one $\mathcal{O}(\alpha_s^1)$ counterterm vertex
 - QCD coupling $\overline{\text{MS}}$ renormalization constant
 - QCD heavy quark field (mass) OS renormalization constant
 - QCD heavy flavor-changing current OS renormalization constant Z_J^{OS}

$$Z_v^{\text{OS}} = Z_a^{\text{OS}} = 1$$

[W.Tao,Z.J.Xiao, JHEP(2023)]
 [W.Tao,Z.J.Xiao, JHEP(2024)]

$$Z_t^{\text{OS}} = Z_{t5}^{\text{OS}} = 1 + \frac{\alpha_s C_F}{4\pi} \left(\frac{1}{\epsilon} - \frac{2x \log x}{1+x} + 2 \log y + \mathcal{O}(\epsilon) \right) + \mathcal{O}(\alpha_s^2)$$

$$x = \frac{m_c}{m_b}, \quad y = \frac{\mu}{m_b}, \quad s = \frac{1}{1 - \frac{q^2}{m_b^2}}$$

Outline

1

Introduction

2

Calculation Procedure

3

Results

4

Summary

Analytical results for $B_c^* \rightarrow J/\psi$ form factors

➤ LO

$$V_1 = \frac{16\sqrt{2}\pi\alpha_s C_F s^2 (1+x)^{\frac{5}{2}} \Psi_{B_c^*}(0) \Psi_{J/\psi}(0)}{m_b^3 x^{\frac{3}{2}} (1+s(x-2)x)^2},$$

$$V_2 = A_2 = T_2 = \frac{1-x}{1+x} V_1 = \frac{2(1+x)}{1+3x} T_6,$$

$$T_3 = \frac{1+x}{2x} T_4 = \frac{-1+s(4+10x+3x^2)}{2s(1+x)(1+3x)} V_1,$$

$$V_1 = \frac{V_3}{2} = A_1 = \frac{1+x}{4x} V_4 = \frac{s(1+x)(1+3x)}{1+4sx+3sx^2} T'_2,$$

$$T'_4 = \frac{1+x}{2} T'_3 = \frac{1+3x}{2(1+x)} V_1,$$

$$V_5 = V_6 = A_3 = A_4 = 0,$$



$\Psi_{B_c^*(J/\psi)}(0)$: $B_c^*(J/\psi)$ wavefunction at origin
 $n_f = n_b + n_c + n_l$

➤ Asymptotic NLO

$$\begin{aligned} \frac{V_1^{\text{NLO}}}{V_1^{\text{LO}}} = & 1 + \frac{\alpha_s}{4\pi} \left\{ \left(\frac{11C_A}{3} - \frac{2}{3}n_f \right) \ln \frac{2sy^2}{x} - \frac{10}{9}n_f + \left(\frac{2\ln s}{3} - \frac{2\ln x}{3} + \frac{10}{9} + \frac{2\ln 2}{3} \right) n_b \right. \\ & - C_A \left[\frac{\ln^2 x}{2} + \left(\ln s + 2\ln 2 + \frac{3}{2} \right) \ln x + \frac{1}{2} \ln^2 s + \left(\frac{3}{2} + 2\ln 2 \right) \ln s + 2\ln^2 2 + \frac{3\ln 2}{2} \right. \\ & - \frac{1}{9} (67 - 3\pi^2) \Big] + C_F \left[2\text{Li}_2(1-s) + \ln^2 x + (2\ln s + 10\ln 2 - 5) \ln x + 2\ln^2 s \right. \\ & \left. \left. + (10\ln 2 - 2) \ln s + 7\ln^2 2 + 9\ln 2 + \frac{1}{3} (\pi^2 - 51) \right] \right\}, \end{aligned}$$

Renormalization scale dependence of form factors



- At maximum recoil point ($q^2 = 0$)
- $\mu_0 = 7 \text{ GeV}$
- $m_b = 4.75 \text{ GeV}$
- $m_c = 1.5 \text{ GeV}$



NLO corrections reduce the renormalization scale dependence

q^2 dependence of NLO to LO form factor ratio

$\mu = 3 \text{ GeV}$



- NLO corrections are both significant and convergent in relatively small q^2 region
- The convergence breaks down in large q^2 region

NRQCD+Lattice predictions for $B_c^* \rightarrow J/\psi$ form factors at $q^2 = 0$

$$F_{i,\text{NRQCD+Lattice}}^{B_c^* \rightarrow J/\psi}(q^2) = \frac{1}{4} \sum_{j=1}^4 \frac{F_{i,\text{NRQCD}}^{B_c^* \rightarrow J/\psi}(q^2)}{F_{j,\text{NRQCD}}^{B_c \rightarrow J/\psi}(q^2)} F_{j,\text{Lattice}}^{B_c \rightarrow J/\psi}(q^2)$$

$$\Psi_{B_c^*}(0) \approx \Psi_{B_c}(0)$$

$$F_j^{B_c \rightarrow J/\psi} \in \{V, A_{0,1,2}\}^{B_c \rightarrow J/\psi}$$

The second uncertainties from lattice data dominate over the first uncertainties from $\mu = 3^{+4}_{-1.5}$ GeV

[HPQCD, PRD(2020)]
 [Q.Chang,L.T.Wang,X.N.Li, JHEP(2019)]
 [Q.Chang,et al., AHEP(2020)]

	NRQCD+Lattice	LFQM [7, 8]
V_1	$0.4320^{+0.0030}_{-0.0048} \pm 0.0448$	$0.56^{+0.01+0.17}_{-0.01-0.17}$
V_2	$0.2295^{+0.0003}_{-0.0004} \pm 0.0238$	$0.33^{+0.01+0.05}_{-0.01-0.04}$
V_3	$0.8865^{+0.0001}_{+0.0003} \pm 0.0919$	$1.17^{+0.02+0.23}_{-0.02-0.29}$
V_4	$0.4294^{-0.0009}_{+0.0018} \pm 0.0445$	$0.65^{+0.01+0.20}_{-0.01-0.19}$
V_5	$0.1303^{-0.0338}_{+0.0569} \pm 0.0135$	$0.20^{+0.00+0.02}_{-0.00-0.02}$
V_6	$0.1303^{-0.0338}_{+0.0569} \pm 0.0135$	$0.20^{+0.00+0.02}_{-0.00-0.02}$
A_1	$0.4458^{-0.0006}_{+0.0013} \pm 0.0462$	$0.54^{+0.01+0.16}_{-0.01-0.17}$
A_2	$0.2510^{-0.0053}_{+0.0090} \pm 0.0260$	$0.35^{+0.00}_{-0.00}$
A_3	$0.0942^{-0.0244}_{+0.0411} \pm 0.0098$	$0.13^{+0.00+0.03}_{-0.00-0.02}$
A_4	$0.1092^{-0.0284}_{+0.0477} \pm 0.0113$	$0.14^{+0.00+0.02}_{-0.00-0.02}$
T_2	$0.2352^{-0.0012}_{+0.0021} \pm 0.0244$	—
T_3	$0.5724^{-0.0035}_{+0.0062} \pm 0.0593$	—
T_4	$0.2790^{-0.0028}_{+0.0049} \pm 0.0289$	—
T_6	$0.2211^{-0.0131}_{+0.0221} \pm 0.0229$	—
T'_2	$0.4387^{+0.0012}_{-0.0018} \pm 0.0455$	—
T'_3	$0.4546^{+0.0115}_{-0.0190} \pm 0.0471$	—
T'_4	$0.3805^{-0.0136}_{+0.0230} \pm 0.0394$	—

NRQCD+Lattice+Z-series predictions in full q^2 range

m_R : masses of low-lying
 $c\bar{b}$ resonances





$$F_i(q^2) = \frac{1}{1 - \frac{q^2}{m_R^2}} \sum_{n=0}^N \alpha_{i,n} z^n(q^2),$$

$$z(q^2) = \frac{\sqrt{t_+ - q^2} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - q^2} + \sqrt{t_+ - t_0}},$$

$$t_0 = t_+ \left(1 - \sqrt{1 - \frac{t_-}{t_+}} \right),$$

$$t_{\pm} = (m_{B_c^*} \pm m_{J/\psi})^2,$$

[C.G.Boyd,B.Grinstein,R.F.Lebed, PRD(1997)]
[D.Leljak,B.Melic,M.Patra, JHEP(2019)]



- Obtain complete and asymptotic analytical results for NLO QCD corrections to $B_c^* \rightarrow J/\psi$ (axial-)vector and (axial-)tensor form factors
- NLO corrections reduce renormalization scale dependence, and are both significant and convergent in relatively small q^2 region
- Provide NRQCD+Lattice+Z-series predictions for $B_c^* \rightarrow J/\psi$ form factors over full q^2 range

Thank you!