

Probing Top-Philic Particles with Boosted Four Top Searches

李浩林

中山大学物理学院



Based on 2404.14482 with Luc Darme, Benjamin Fuks,
Matteo Maltoni, Olivier Mattelaer, Julien Toucheque

New physics and top quark

- **Largest couplings to Higgs**

- Sensitive to Models that modify the EWSB or solving the Hierarchy Problem


- e.g. Composite Higgs Model, SUSY, Extra Dimension...

- Radiative correction to Higgs potential thus sensitive to vacuum stability

- Top-philic Models – New particle preferentially couple to Top

- **LHC is essentially a top factory**

- Top copiously produced, enabling precision study.



Topophilic NP theories: the origin

- › In this talk, Topophilic scalar particle X , $SU(3)$ octet and singlet $y_s \bar{t}tS$
 - Which decays mostly into a pair of top quarks $y_s \bar{t}T^a t S_8^a$
 - Production cannot rely on EW gauge boson or quarks in proton

Topophilic NP theories: the origin

- › In this talk, Topophilic scalar particle X , $SU(3)$ octet and singlet $y_s \bar{t} t S$
 - Which decays mostly into a pair of top quarks $y_s \bar{t} T^a t S_8^a$
 - Production cannot rely on EW gauge boson or quarks in proton
- › Why would a New Physics (NP) boson prefers the top quarks

Because the quark mass
enters into the coupling (e.g.
 $SU(2)$ breaking required)

Extended SUSY (sgluon)

e.g. 2107.13565

N=1 SUSY	N=1 SUSY
O_a	\tilde{g}_a
Spin 0	Spin 1/2

g_a

Topophilic NP theories: the origin

- › In this talk, Topophilic scalar particle X , $SU(3)$ octet and singlet
 - Which decays mostly into a pair of top quarks
 - Production cannot rely on EW gauge boson or quarks in proton
- › Why would a New Physics (NP) boson prefers the top quarks

Because the quark mass enters into the coupling (e.g. $SU(2)$ breaking required)

Extended SUSY (sgluon)
e.g. 2107.13565

N=1 SUSY	N=1 SUSY
O_a	\tilde{g}_a
Spin 0	Spin 1/2
	Spin 1

Because the top quark is made (partially) of NP

Partial top compositeness

e.g. 1507.02283, 1610.06591

Color coset	$SU(3)_c \times U(1)_Y$
$SU(6)/SO(6)$	$\mathbf{8}_0 + \mathbf{6}_{(-2/3 \text{ or } 4/3)} + \bar{\mathbf{6}}_{(2/3 \text{ or } -4/3)}$
$SU(6)/Sp(6)$	$\mathbf{8}_0 + \mathbf{3}_{2/3} + \bar{\mathbf{3}}_{-2/3}$
$SU(3) \times SU(3)' / SU(3)_D$	$\mathbf{8}_0$

Top-philic NP theories: the origin

- In this talk, Top-philic scalar particle X , $SU(3)$ octet and singlet $y_s \bar{t}tS$
 - Which decays mostly into a pair of top quarks $y_s \bar{T}^a t S_8^a$
 - Production cannot rely on EW gauge boson or quarks in proton
- Why would a New Physics (NP) boson prefers the top quarks

Because the quark mass enters into the coupling (e.g. SU(2) breaking required)

Extended SUSY (sgluon)
e.g. 2107.13565

N=1 SUSY	N=1 SUSY
O_a	\tilde{g}_a
Spin 0	Spin 1/2

Because the top quark is made (partially) of NP

Partial top compositeness
e.g. 1507.02283, 1610.06591

Because the NP helps in generating the top quark mass

Extended Higgs sectors
e.g. 2202.02333

Color coset	$SU(3)_c \times U(1)_Y$
$SU(6)/SO(6)$	$\mathbf{8}_0 + \mathbf{6}_{(-2/3 \text{ or } 4/3)} + \bar{\mathbf{6}}_{(2/3 \text{ or } -4/3)}$
$SU(6)/Sp(6)$	$\mathbf{8}_0 + \mathbf{3}_{2/3} + \bar{\mathbf{3}}_{-2/3}$
$SU(3) \times SU(3)' / SU(3)_D$	$\mathbf{8}_0$


$$\mathcal{L}_{\text{Yukawa}}^{\text{2HDM}} = - \sum_{f=u,d,l} \frac{m_f}{v} (\xi_h^f \bar{f} f h + \xi_H^f \bar{f} f H - i \xi_A^f \bar{f} \gamma_5 f A)$$

	Type I	Type II	Lepton-specific
ξ_H^u	$\sin \alpha / \sin \beta$	$\sin \alpha / \sin \beta$	$\sin \alpha / \sin \beta$
ξ_A^u	$\cot \beta$	$\cot \beta$	$\cot \beta$

Heavy top-phobic NP Production @LHC

Final state: 4 tops


Associate production



$$\sigma \propto y_s^2$$

Need energetic gluons
Depend on y_s

pair production




$$\sigma \propto \alpha_s^2$$

Need energetic gluons and
only works for octet

Final state: di-top

single production



$$\sigma \propto \frac{1}{(4\pi)^4} y_s^2$$

Loop-induced,
Good for lighter resonance,
May suffer large background



Four top theory prediction summary

- **SM NLO (QCD+EW) predictions (MadGraph):**
Full NLO correction up to order $\mathcal{O}(\alpha_s^i \alpha^j)$ with $i+j = 4, 5$. Treating top quark as stable particle, i.e. without decay.
Frederix, Pagani, Zaro 1711.02116
- **SM NLO QCD matched to parton shower (POWHEG-BOX):**
NLO QCD + LO EW, LO top quark spin correlation effect are included in the decay.
Ježo, Kraus 2110.15159
- **SM NLO (QCD+EW) + NLL' :**
NLL threshold resummation, without top decay.
Bekkeld, Kulesza, Valero 2212.03259
- **SM NLO QCD with 3/4 leptonic decay channel:**
Higher order QCD effect in the decay are included.
Dimitrakopoulos, Wore 2410.05960 2401.10678
- **SMEFT correction at LO and NLO QCD up to dim-6:**
Aoude et.al, 2208.04962; Degrande, Rosenfeld, Vasquez 2402.06528

SM four top measurements @ LHC

For SM most signal are produced at threshold **very hard to reconstruct tops**
Searches from ATLAS and CMS mostly focus on the **multilepton final states**:




ATLAS[[2303.15061](#)] 2LSS and 3L – 6.1σ
CMS[[2305.13439](#)] 4L 3L and 2LSS – 5.6σ



Boosted 4-top searches for NP

If the Top-philic particle are heavy (>1 TeV) then the four top produced are highly boosted, one can leverage the boosted top tagging developed in recent years to search for signal.

Aim for: four reconstructed tops in fully hadronic decay and 1L channel ($\sim 60\%$ BR)




Top tag mis-tagging rate

Simulation and Event selection

FeynRules UFO model → **MadGraph@NLO** → **Pythia8** → **MadAnalysis5**

Reconstruction:

- AK10-jets: $pT > 350 \text{ GeV}$ and $|\eta| < 2$, top-tag with 80% efficiency if $dR(\text{jet}, \text{top}) \leq 0.75$
- AK4-jets: with $pT > 30 \text{ GeV}$ $|\eta| < 2.5$ b-tag with MV2c10 77% WP
- Isolated lepton, remove AK10-jet if $dR(l, \text{AK10-jet}) < 1.0$
- Remove b-tagged jets if $dR(\text{bjet}, \text{AK10-jet}) < 1.0$



Background study

Process	σ (LO)	Scale	PDF	σ (NLO)	Scale	PDF
$t\bar{t}jj$	354	+62% -35%	$\pm 5.8\%$	352	+3.7% -13%	$\pm 2.6\%$
$t\bar{t}W$	0.376	+23% -17%	$\pm 3.9\%$	0.565	+8.3% -8.3%	$\pm 1.8\%$
$t\bar{t}Wj$	0.329	+39% -26%	$\pm 2.1\%$	0.452	+8.1% -12%	$\pm 1.2\%$
$t\bar{t}Z$	0.563	+31% -22%	$\pm 4.8\%$	0.756	+9.2% -11%	$\pm 2.1\%$
$t\bar{t}Zj$	0.639	+47% -30%	$\pm 6.5\%$	0.672	+2.6% -9%	$\pm 2.5\%$
$t\bar{t}t\bar{t}$	0.00612	+65% -37%	$\pm 13\%$	0.00920	+28% -24%	$\pm 6.0\%$
$t\bar{t}t + t\bar{t}\bar{t}$	0.00155	+22% -17%	$\pm 13\%$	0.00201	+20% -19%	$\pm 7.5\%$


Backgrounds are simulated at leading order and rescaled to NLO cross-section

Background study

Process	σ (LO)	Scale	PDF	σ (NLO)	Scale	PDF
$t\bar{t}jj$	354	+62% -35%	$\pm 5.8\%$	352	+3.7% -13%	$\pm 2.6\%$
$t\bar{t}W$	0.376	+23% -17%	$\pm 3.9\%$	0.565	+8.3% -8.3%	$\pm 1.8\%$
$t\bar{t}Wj$	0.329	+39% -26%	$\pm 2.1\%$	0.452	+1.8% -1.8%	$\pm 1.2\%$
$t\bar{t}Z$	0.563	+31% -22%	$\pm 4.8\%$			
$t\bar{t}Zj$	0.639	+47% -30%	$\pm 6.5\%$			
$t\bar{t}t\bar{t}$	0.00612	+65% -37%	$\pm 13\%$			
$t\bar{t}t + t\bar{t}\bar{t}$	0.00155	+22% -17%	$\pm 13\%$			


After all the selection cuts
the dominant background is
 $t\bar{t}jj$ with like jet false tagged

Backgrounds are simulated at leading order and rescaled to NLO cross-section




BP1: $M=1.3\text{TeV}$, $ys=0.25$, $\Gamma=1.45\text{ GeV}$ pair dominant
BP2: $M=2\text{TeV}$, $ys=1.0$, $\Gamma=38.04\text{ GeV}$ mixed production

Result



Octet exclude $M < 2.1$ TeV for $L = 3000$ fb $^{-1}$
due to superior pair production efficiency.




For singlet it outperform
conventional search at larger mass

NLO signal computation (preliminary)

- We generate NLO UFO model with **FeynRules** + **NLOCT** including UV and R2 counter-terms
- Compute Full NLO QCD correction to LO NP four tops production match to parton shower:

$$\mathcal{O}(\alpha_s y_S^2) \rightarrow \mathcal{O}(\alpha_s^2 y_S^2)$$




Include corrections to both **production** and **decay**

NLO signal computation (preliminary)

- We generate NLO UFO model with **FeynRules** + **NLOCT** including UV and R2 counter-terms
- Compute Full NLO QCD correction to LO NP four tops production match to parton shower:

$$\mathcal{O}(\alpha_s y_S^2) \rightarrow \mathcal{O}(\alpha_s^2 y_S^2)$$




Include corrections to both **production** and **decay**

	Octet M=1TeV, y=1	Octet M=2TeV, y=1	Singlet M=1.5 TeV, y=1
$\sigma_{4t,\text{NLO}}$	$54.27^{+9.4\%}_{-13.1\%} {}^{+9\%}_{-9\%}$	$0.2199^{+15.9\%}_{-16.7\%} {}^{+16.9\%}_{-16.9\%}$	$0.4512^{+19.9\%}_{-17.9\%} {}^{+11\%}_{-11\%}$
K_{4t}	1.74	1.69	1.87
$\sigma_{OO/\text{ttS},\text{NLO}}$	$45.37^{+17.5\%}_{-17.1\%} {}^{+9\%}_{-9\%}$	$0.06859^{+19.3\%}_{-18.2\%} {}^{+27\%}_{-27\%}$	$0.3909^{+19.5\%}_{-18.7\%} {}^{+12.1\%}_{-12.1\%}$
$K_{OO/\text{ttS}}$	1.63	1.56	1.86

NLO signal computation (preliminary)



- We generate NLO UFO model with **FeynRules** + **NLOCT** including UV and R2 counter-terms
- Compute Full NLO QCD correction to LO NP four tops production match to parton shower:

$$\mathcal{O}(\alpha_s y_S^2) \rightarrow \mathcal{O}(\alpha_s^2 y_S^2)$$



Include corrections to both **production** and **decay**

Distribution of reconstructed M_{tt} for the **full Hadronic decay channel**, passing all the cuts while allowing one AK10-jet not top tagged.



Summary

- Top-philic particle present in different types of UV theory.
- Color octet can be pair produced at LO and without suppression from the scalar couplings
- Fully hadronic decay and 1 leptonic decay channels are good for searching top-philic particle in the mass range 1-3 TeV
- The NLO QCD correction to the top-philic particle production is significant for both pair and associative production.

	FullNP_gg	SmintNP_gg	qq4tNP	qq4tSMint
Msig8=1TeV y=1	0.04406 +- 0.0001438	7.206e-05 +- 1.49e-06	0.008611 +- 1.487e-05	4.439e-07 +- 6.706e-08
Msig8=2TeV y=1	0.000241 +- 6.113e-07	1.859e-05 +- 2.699e-07	1.183e-05 +- 1.442e-08	3.439e-08 +- 7.566e-09
Msig8=3TeV y=1	9.917e-06 +- 2.703e-08	8.323e-06 +- 9.265e-08	1.053e-07 +- 2.718e-10	1.833e-08 +- 3.664e-09
Msig8=1TeV y=2	0.04697 +- 0.0001081	0.0002819 +- 5.179e-06	0.003085 +- 7.715e-06	1.222e-06 +- 2.565e-07
Msig8=2TeV y=2	0.000552 +- 1.416e-06	7.576e-05 +- 1.052e-06	1.381e-05 +- 3.831e-08	9.743e-08 +- 4.138e-08
Msig8=3TeV y=2	4.233e-05 +- 1.498e-07	3.387e-05 +- 3.958e-07	3.267e-07 +- 9.956e-10	7.956e-08 +- 1.466e-08

	double only gg	single only gg	t only gg
Msig8=1TeV y=1	0.03322 +- 6.773e-05	0.007346 +- 1.242e-05	2.252e-05 +- 8.959e-08
Msig8=2TeV y=1	5.82e-05 +- 1.227e-07	0.0001617 +- 3.479e-07	3.066e-06 +- 1.151e-08
Msig8=3TeV y=1	1.873e-07 +- 3.784e-10	8.113e-06 +- 1.991e-08	7.735e-07 +- 2.557e-09
Msig8=1TeV y=2	0.0306 +- 7.811e-05	0.02854 +- 6.694e-05	0.0003607 +- 1.383e-06
Msig8=2TeV y=2	6.025e-05 +- 1.436e-07	0.0007248 +- 2.257e-06	4.858e-05 +- 1.842e-07
Msig8=3TeV y=2	3.151e-07 +- 6.268e-10	5.183e-05 +- 1.712e-07	1.236e-05 +- 3.037e-08

Top tag. \mathcal{L} [fb $^{-1}$]	BP1				BP2			
	Optimistic		Conservative		Optimistic		Conservative	
	400	3000	400	3000	400	3000	400	3000
SR1	1.35	0.52	1.69	0.64	0.68	0.24	0.82	0.30
SR2	0.64	0.26	0.75	0.36	0.51	0.14	0.61	0.20
SSL	0.97	0.27	0.97	0.27	1.13	0.29	1.12	0.28