



Institute of Particle Physics
粒子物理研究所

b-hadron FCNC decays as a probe of light Dark Matter

Xing-Bo Yuan (袁兴博)

Central China Normal University (华中师范大学)


Biao-Feng Hou, Xin-Qiang Li, Meng Shen, Ya-Dong Yang, XBY, JHEP06(2024)172

Meng-Chao Gao, Xin-Qiang Li, Ya-Dong Yang, XBY, Xin Zhang, work in progress

Physics beyond the Standard Model

Problems of the SM $\implies$ Physics beyond the SM


- experimental
- Dark Matter (26%)
 - Dark Energy (69%)
 - Matter-antimatter asymmetry
 - Anomalies in experimental data
- conceptual
- Neutrino masses
 - Strong CP problem
 - Vacuum metastability
 - NP flavour puzzle
 - Gravity
 - Hierarchy problem
 - SM flavour puzzle
 - Family replication ...




$$- \text{t-loop} + \dots = \frac{c}{16\pi^2} \Lambda^2$$

fine-tuning

$$m_{h,0}^2 + \frac{c}{16\pi^2} \Lambda^2 = 125 \text{ GeV}^2$$



Theories beyond the SM



Flavour Physics as a Probe of New Physics

► **Flavour universal**

- couplings $\propto \delta_{ij}$ in flavour space
- example: strong and electromagnetic interactions
- consequence of gauge invariance

► **Flavour diagonal**

- couplings $\propto \lambda_i \delta_{ij}$ (diagonal, but not necessarily universal)
- example: Yukawa interactions

► **Flavour violation (changing)**

- couplings involve different quarks
- no flavour violation in lepton sector ($m_\nu = 0$)
- example: $W^\pm$ interactions in quark sector

► **Flavour Changing Neutral Current (FCNC)**


- absent at the tree-level
- arise at the one-loop, but suppressed by GIM mechanism

► **Why flavour physics**


- New physics $\iff \mathcal{O}(10^9)$ $B\bar{B}$ events at BaBar and Belle
- structure of CKM and mass
- CP violation
- strong interaction

experimental status
no evidence of NP $> 5\sigma$
but, anomalies $2 \sim 4\sigma$


flavour universal



flavour diagonal




flavour violation



CKM matrix

► Cabibbo–Kobayashi–Maskawa matrix

$$\begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} \end{bmatrix}$$





► 3 mixing angles and 1 CP phase

► CP violation in the Standard Model

not enough to explain the baryon asymmetry in our universe

new CP violation sources



penguin diagram

Theoretically cleanest processes in heavy flavour physics

$B_{s,d} - \bar{B}_{s,d}$ mixing

$b \rightarrow u(c)\ell\bar{\nu}$: $B \rightarrow D\ell\bar{\nu}$, $B \rightarrow D^*\ell\bar{\nu}$,

$b \rightarrow s(d)\gamma$: $B \rightarrow X_s\gamma$, $B \rightarrow K^*\gamma$,

$b \rightarrow s(d)\ell^+\ell^-$: $B_s \rightarrow \ell^+\ell^-$, $B \rightarrow K^*\ell^+\ell^-$,


$b \rightarrow s(d)\nu\bar{\nu}$: $B \rightarrow K\nu\bar{\nu}$, $K \rightarrow \pi\nu\bar{\nu}$, ...

NP Effects $< \mathcal{O}(20\%)$



Signal: $B \rightarrow K + \text{inv}$ NOT measured yet !

$B \rightarrow K + \nu + \bar{\nu}$
 $B \rightarrow K + \text{DM} + \text{DM}$



Dark Matter Detection





Collider



Direct Detection






Indirect Detection





Dark Matter Detection

Collider





Direct
Detection



Hadron FCNC Decay


$$\begin{aligned} B^+ &\rightarrow K^+ + \text{DM} + \text{DM} \\ \Lambda_b &\rightarrow \Lambda + \text{DM} + \text{DM} \\ K^+ &\rightarrow \pi^+ + \text{DM} + \text{DM} \\ D^0 &\rightarrow \pi^0 + \text{DM} + \text{DM} \end{aligned}$$

Indirect
Detection



Light DM: a bottom-up view

DM is electrically neutral ! $\implies$ DM only have FC or FCNC couplings to fermions !




	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II/CEPC
<i>s</i>		DD	Belle II/CEPC
<i>b</i>			Belle II/LHC
	<i>u</i>	<i>c</i>	<i>t</i>
<i>u</i>	DD	BES/STCF	LHC/CEPC
<i>c</i>		BES/STCF	LHC/CEPC
<i>t</i>			LHC/CEPC

example:
 $B^+ \rightarrow K^+ + \text{DM} + \text{DM}$
 $K^+ \rightarrow \pi^+ + \text{DM} + \text{DM}$
 $D^0 \rightarrow \pi^0 + \text{DM} + \text{DM}$

DD = Direct Detection

 means related to the DM relic density




Experimental Search

	Observable	SM	Exp	Unit
$b \rightarrow s$	$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
	$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
	$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
	$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}
$b \rightarrow d$	$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
	$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}
$s \rightarrow d$	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}
	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}
$c \rightarrow u$	$D^+ \rightarrow \pi^+ + \text{inv}, D^0 \rightarrow \rho^0 + \text{inv}, \dots$			

Belle II
CEPC
FCC-ee

NA62
KOTO

BESIII
STCF



Experimental Search

Observable		SM	Exp	Unit	
$b \rightarrow s$	$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}	Belle II CEPC FCC-ee
	$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}	
	$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}	
	$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}	
	$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}	
	$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}	
$b \rightarrow d$	$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}	NA62 KOTO
	$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}	
	$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}	
	$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}	
	$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}	
$s \rightarrow d$	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}	BESIII STCF
	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}	
$c \rightarrow u \quad D^+ \rightarrow \pi^+ + \text{inv}, D^0 \rightarrow \rho^0 + \text{inv}, \dots \dots$					




figure from Amhis, Kenzie, Reboud, Wiederhold, 2309.11353

Experimental Search


Observable		SM	Exp	Unit
$b \rightarrow s$	$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
	$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
	$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
	$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}
$b \rightarrow d$	$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
	$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}
$s \rightarrow d$	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}
	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}
$c \rightarrow u$ $D^+ \rightarrow \pi^+ + \text{inv}, D^0 \rightarrow \rho^0 + \text{inv}, \dots \dots$				

Belle II
CEPC
FCC-ee


NA62
KOTO

BESIII
STCF

► 2021 Apr



► 2023 Aug: first evidence




Experimental Search

	Observable	SM	Exp	Unit	Expected sensitivities at future Belle II, CEPC, FCC-ee
$b \rightarrow s$	$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5_{-4}^{+5}$	10^{-6}	19%, Belle II, 5 fb^{-1}
	$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}	87%, Belle II, 5 fb^{-1} ; 3.37%, FCC-ee, Tera-Z
	$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}	75%, Belle II, 5 fb^{-1} ; 0.53%, FCC-ee, Tera-Z
	$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}	40%, Belle II, 5 fb^{-1}
	$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}	2%, CEPC, Tera-Z; 1.2% FCC-ee, Tera-Z
	$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}	$< 1.1 \times 10^{-5}$, Belle II, 5 fb^{-1} , $\Upsilon(5S)$; $< 7.2 \times 10^{-9}$, FCC-ee, Tera-Z
$b \rightarrow d$	$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}	Belle II snowmass white paper, 2207.06307 CEPC flavor white paper, 2412.19743 Amhis, Kenzie, Reboud, Wiederhold, 2309.11353 Cartelle, Kenzie, Mangrulkar, Wiederhold, Wood, 2508.04471 see also talks by Lorenzo Calibbi, Xunwu Zuo, Sen Jia
	$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}	
	$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}	
	$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}	
	$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}	
$s \rightarrow d$	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6_{-3.4}^{+4.0} \pm 0.9$	10^{-11}	
	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}	
$c \rightarrow u$	$D^+ \rightarrow \pi^+ + \text{inv}, D^0 \rightarrow \rho^0 + \text{inv}, \dots \dots$				

Light DM: a bottom-up view

DM is electrically neutral ! $\implies$ DM only have FC or FCNC couplings to fermions !



	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II
<i>s</i>		DD	Belle II
<i>b</i>			Belle II/LHC
	<i>u</i>	<i>c</i>	<i>t</i>
<i>u</i>	DD	BES/STCF	LHC/CEPC
<i>c</i>		BES/STCF	LHC/CEPC
<i>t</i>			LHC/CEPC


example:
 $B^+ \rightarrow K^+ + \text{DM} + \text{DM}$
 $K^+ \rightarrow \pi^+ + \text{DM} + \text{DM}$
 $D^0 \rightarrow \pi^0 + \text{DM} + \text{DM}$

future exp uncertainties:
 $\mathcal{O}(20 \sim 40\%)@Belle\ II\ with\ 5ab^{-1}$
 $\mathcal{O}(2 \sim 4\%)@CEPC\ Tera-Z$

future theo uncertainties:
 less than 10 %


DD = Direct Detection

 means related to the DM relic density



Light DM: a bottom-up view

DM is electrically neutral ! $\implies$ DM only have FC or FCNC couplings to fermions !




	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II
<i>s</i>		DD	Belle II
<i>b</i>			Belle II/LHC


example:
 $B^+ \rightarrow K^+ + \text{DM} + \text{DM}$
 $K^+ \rightarrow \pi^+ + \text{DM} + \text{DM}$
 $D^0 \rightarrow \pi^0 + \text{DM} + \text{DM}$

It's natural to ask:

1. By combing the direct detection, flavour measurements, and relic density, is it possible to obtain the flavour structure of interactions between light DM and SM particles ?
2. Compared to DM direct detection, what's the advantage of hadron decays ? which DM scenario is more sensitive to ? Can they provide other information ?




Theoretical Framework



Effective Field Theory approach to combine the various experimental searches

In EFT, DM is just a singlet under the SM gauge group.

for light DM



$H_1 \rightarrow H_2 + \mathbf{DM}$ theoretical calculation and experimental searches

► $d_i \rightarrow d_j + \phi + \phi$

- 2011 Kamenik, Smith
- 2014 Bird, Jackson, Kowalewski, Pospelov
- 2019 G.Li, J.Y. Su, Tandean
- $\Lambda \rightarrow n + \phi\phi, \Sigma^+ \rightarrow p + \phi\phi, \Xi^0 \rightarrow \Lambda + \phi\phi,$
 $\Xi^- \rightarrow \Sigma^- \phi\phi, \Omega^- \rightarrow \Sigma^- + \phi\phi$
- 2020 X.G. He, X.D. Ma, Tandean, Valencia
- 2020 C.Q.Geng, Tandean, $K \rightarrow \pi\pi + \phi\phi$
- 2021 G. Li, T. Wang, J.B. Zhang, G.L. Wang
- 2022 Kling, S. Li, H. Song, S. Su, W. Su

► $d_i \rightarrow d_j + \chi + \chi$

- 2011 Kamenik, Smith
- 2019 J.Y. Su, Tandean
- 2020 G. Li, T. Wang, Y. Jiang, J.B. Zhang, G.L. Wang
- 2021 Felkl, S. L. Li, Schmidt

► $d_i \rightarrow d_j + X + X$

- 2011 Kamenik, Smith
- 2021 G. Li, T. Wang, J.B. Zhang, G.L. Wang
- 2022 X.G. He, X.D. Ma, Valencia

► $d_i \rightarrow d_j + a$

- 2020 Camalich, Pospelov, Vuong, Ziegler, Zupan,
- 2021 Bauer, Neubert, Renner, Schnubel, Thamm
- 2022 Guerrero and S. Rigolin

- theoretically clean: $A \propto C \cdot \langle H_2 | O | H_1 \rangle \cdot \text{DM current}$ ^{form factor}
 - no GIM suppression
 - possibly two-body decay
- enhancement**

Observable
$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$
$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$
$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$
$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$
$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$
$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$
$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$
$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$
$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$
$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$
$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$
$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$
$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$

longitudinal polarization
 F_L

$\Lambda_b \rightarrow \Lambda + \mathbf{DM} + \mathbf{DM}$

► $c \rightarrow u + \mathbf{DM}$

- 2022 C.Q.Geng, G.Li
- 2023 G.Li, Tandean

- $D^0 \rightarrow S \bar{S}'$
- $D^0 \rightarrow \gamma S \bar{S}'$
- $D^0 \rightarrow \pi^0 S \bar{S}'$
- $D^+ \rightarrow \pi^+ S \bar{S}'$
- $D_s^+ \rightarrow K^+ S \bar{S}'$
- $D^0 \rightarrow \rho^0 S \bar{S}'$
- $D^+ \rightarrow \rho^+ S \bar{S}'$
- $D_s^+ \rightarrow K^{*+} S \bar{S}'$
- $\Lambda_c^+ \rightarrow p S \bar{S}'$
- $\Xi_c^+ \rightarrow \Sigma^+ S \bar{S}'$
- $\Xi_c^0 \rightarrow \Sigma^0 S \bar{S}'$
- $\Xi_c^0 \rightarrow \Lambda S \bar{S}'$


$\nu \rightarrow \mathbf{DM}$ **based on complete EFT basis (Dark LEFT)**

HadronToNP: a package to calculate decay of hadron to new particles
 B.F. Hou, X.Q.Li, H.Yan, Y.D.Yang, **XBY** *to be finished*

$b \rightarrow s\nu\bar{\nu}$: DSMEFT

Can DSMEFT operators explain the Belle II excess, while satisfy other $b \rightarrow s$ bounds ?

Observable	SM	Exp	Unit
$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}
$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}
$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}
$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}



Dark SMEFT

$$\begin{aligned} \mathcal{Q}_{d\phi} &= (\bar{q}_p d_r H) \phi + \text{h.c.}, & \mathcal{Q}_{d\phi^2} &= (\bar{q}_p d_r H) \phi^2 + \text{h.c.}, \\ \mathcal{Q}_{\phi q} &= (\bar{q}_p \gamma_\mu q_r) (i \phi_1 \overleftrightarrow{\partial}^\mu \phi_2), & \mathcal{Q}_{\phi d} &= (\bar{d}_p \gamma_\mu d_r) (i \phi_1 \overleftrightarrow{\partial}^\mu \phi_2), \\ \mathcal{Q}_{q\chi} &= (\bar{q}_p \gamma_\mu q_r) (\bar{\chi} \gamma^\mu \chi), & \mathcal{Q}_{d\chi} &= (\bar{d}_p \gamma_\mu d_r) (\bar{\chi} \gamma^\mu \chi), \\ \mathcal{Q}_{dHX} &= (\bar{q}_p \sigma_{\mu\nu} d_r) H X^{\mu\nu} & \mathcal{Q}_{dX^2} &= (\bar{q}_p d_r H) X_\mu X^\mu \\ \mathcal{Q}_{qa} &= (\bar{q}_p \gamma_\mu q_r) \partial^\mu a & \mathcal{Q}_{da} &= (\bar{d}_p \gamma_\mu d_r) \partial^\mu a \end{aligned}$$

scalar: 4

fermion: 2

vector: 1+13

ALP: 2

Dark LEFT

$$\begin{aligned} \mathcal{O}_{d\phi} &= (\bar{d}_{Lp} d_{Rr}) \phi + \text{h.c.}, & \mathcal{O}_{\phi d}^L &= (\bar{d}_{Lp} \gamma_\mu d_{Lr}) (i \phi_1 \overleftrightarrow{\partial}^\mu \phi_2), \\ \mathcal{O}_{d\chi}^{V,LR} &= (\bar{d}_{Lp} \gamma_\mu d_{Lr}) (\bar{\chi}_a \gamma^\mu \chi_b), & \mathcal{O}_{d\chi}^{V,RR} &= (\bar{d}_{Rp} \gamma_\mu d_{Rr}) (\bar{\chi}_a \gamma^\mu \chi_b), \\ \mathcal{O}_{dX}^T &= (\bar{d}_{Lp} \sigma_{\mu\nu} d_{Rr}) X_a^{\mu\nu} & \mathcal{O}_{dXX}^L &= (\bar{d}_{Lp} \gamma_\mu d_{Lr}) X^{\mu\nu} X_\nu \\ \mathcal{O}_{da}^L &= (\bar{d}_{Lp} \gamma_\mu d_{Lr}) \partial^\mu a, & \mathcal{O}_{da}^R &= (\bar{d}_{Rp} \gamma_\mu d_{Rr}) \partial^\mu a. \end{aligned}$$







scalar: 4

fermion: 5












vector: 1+10

ALP: 2



Dark SMEFT: Scalar




Dark SMEFT: Vector



Dark SMEFT: Fermion, ALP



One example





Summary of the numerical analysis

All the operators survive from the constraints of the various FCNC decays.

In the future, all the parameter space to explain the Belle II excess can be covered by combining

- ▶ $B_s \rightarrow \text{inv}$
- ▶ $b \rightarrow s + \text{inv} (B_s \rightarrow \phi + \text{inv}, B^0 \rightarrow K^0 + \text{inv}, \dots)$
- ▶ Belle II, **CEPC, FCC-ee**



Dark SMEFT: $\mathrm{d}B/\mathrm{d}q^2$



Difficult to distinguish the DSMEFT operators by considering only the $B^+ \rightarrow K^+ \nu \bar{\nu}$ decay. However,

Dark SMEFT: $dB/dq^2, F_L$

$m_{\text{DM}} = 1500 \text{ MeV}$




All the operators are **distinguishable** from each other by combining these observables

Effective Field Theory approach to combine the various experimental searches

In EFT, DM is just a singlet under the SM gauge group.

for light DM



Top-flavored DM

► Dark SMEFT with 3rd generation@ μ_{EW}

$$\mathcal{Q}_{u\chi} = (\bar{u}_p \gamma_\mu u_r)(\bar{\chi} \gamma^\mu \chi), \implies (\bar{t}_R \gamma_\mu t_R)(\bar{\chi} \gamma^\mu \chi)$$

$$\mathcal{Q}_{q\chi} = (\bar{q}_p \gamma_\mu q_r)(\bar{\chi} \gamma^\mu \chi),$$

$$\mathcal{Q}_{u\chi^2} = (\bar{q}_p u_r \tilde{H})(\bar{\chi}\chi), \quad \mathcal{C} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathcal{C}_{33}$$

$$\mathcal{O}_{d\phi^2} = (\bar{d}_p P_R d_r) \phi^2$$


$$\mathcal{O}_{dXX}^L = (\bar{d}_p \gamma^\mu P_L d_r) X_{\mu\nu} X^\nu$$

$$\mathcal{O}_{d\chi}^{V,LR}=(\bar{d}_p\gamma_\mu P_L d_r)(\bar{\chi}\gamma^\mu\chi),$$

+ $\mathcal{O}(20)$ operators

$$\langle H_2 | \mathcal{O}_i | H_1 \rangle$$

hadron decay

 μ_{EW} μ_b


1 GeV

Dark SMEFT

one-loop matching

Dark LEFT

direct detection



$$\mathcal{O}_{g\phi} = G_{\mu\nu}^a G^{\mu\nu,a} \phi^2,$$

$$\mathcal{O}_{gX} = G_{\mu\nu}^a G^{\mu\nu,a} X_\rho X^\rho,$$

$$\mathcal{O}_{g\chi} = G_{\mu\nu}^a G^{\mu\nu,a} \bar{\chi}\chi$$

+ operators with u, d, s

$$\langle N | \mathcal{O}_j | N \rangle$$

2013 Tongyan Lin, Kolb, Lian-Tao Wang

2015 Kilic, Klimek, Jiang-Hao Yu

2015 Haisch, Re

2015 Boucheneb, Cacciapaglia, Deandrea, Fuks

2017 Blanke, Kast

2021 Blanke, Pani, Polesello, Rovelli

2021 Haisch, Polesello, Schulte

2021 Hermanna, Worek

2022 Yandong Liu, Bin Yan, Rui Zhang

■ ■ ■ ■ ■

2019 ATLAS [JHEP05(2019)142]


One-Loop Matching between Dark SMEFT and Dark LEFT

► Dark SMEFT

$$\mathcal{L}_{\text{DSMEFT}} \supset \sum_i \mathcal{C}_i \mathcal{Q}_i^{(4)} + \boxed{\frac{1}{\Lambda} \sum_j \mathcal{C}_j \mathcal{Q}_j^{(5)}} + \boxed{\frac{1}{\Lambda^2} \sum_k \mathcal{C}_k \mathcal{Q}_k^{(6)}} + \boxed{\frac{1}{\Lambda^3} \sum_l \mathcal{C}_l \mathcal{Q}_l^{(7)}}$$

$\mathcal{Q}_{u\phi^2} = (\bar{q}_p u_r \tilde{H}) \phi^2 + \text{h.c.}$	$\mathcal{Q}_{\phi u} = (\bar{u}_p \gamma_\mu u_r) (i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$	$\mathcal{Q}_{\phi q} = (\bar{q}_p \gamma_\mu q_r) (i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$	...	scalar
$\mathcal{Q}_{u\chi^2} = (\bar{q}_p u_r \tilde{H}) (\chi^T C \chi)$	$\mathcal{Q}_{u\chi} = (\bar{u}_p \gamma_\mu u_r) (\bar{\chi} \gamma^\mu \chi)$	$\mathcal{Q}_{q\chi} = (\bar{q}_p \gamma_\mu q_r) (\bar{\chi} \gamma^\mu \chi)$	...	fermion
$\mathcal{Q}_{qXX} = (\bar{q}_p \gamma_\mu q_r) X^{\mu\nu} X_\nu$	$\mathcal{Q}_{u\tilde{X}X} = (\bar{u}_p \gamma_\mu u_r) \tilde{X}^{\mu\nu} X_\nu$	$\mathcal{Q}_{uX^2} = (\bar{q}_p u_r \tilde{H}) X_\mu X^\mu + \text{h.c.}$	...	vector

► Matching



► Dark LEFT: **hadron detection**

$$\mathcal{L}_{\text{DLEFT}} \supset \sum_k L_k \mathcal{O}_k^{(4)} + \boxed{\frac{1}{v} \sum_i L_i \mathcal{O}_i^{(5)}} + \boxed{\frac{1}{v^2} \sum_j L_j \mathcal{O}_j^{(6)}} + \boxed{\frac{1}{v^3} \sum_l L_l \mathcal{O}_l^{(7)}}$$

$\mathcal{O}_{\phi d}^R = (\bar{d}_p \gamma_\mu P_R d_r) (i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$	$\mathcal{O}_{d\phi^2} = (\bar{d}_p P_R d_r) \phi^2$	...	scalar
$\mathcal{O}_{d\chi}^{V,RR} = (\bar{d}_p \gamma_\mu P_R d_r) (\bar{\chi} \gamma^\mu \chi)$	$\mathcal{O}_{d\chi}^{V,LR} = (\bar{d}_p \gamma_\mu P_L d_r) (\bar{\chi} \gamma^\mu \chi)$	...	fermion
$\mathcal{O}_{dXX}^R = (\bar{d}_p \gamma^\mu P_R d_r) X_{\mu\nu} X^\nu$	$\mathcal{O}_{dXX}^L = (\bar{d}_p \gamma^\mu P_L d_r) X_{\mu\nu} X^\nu$	...	vector


One-Loop Matching between Dark SMEFT and Dark LEFT

► Dark SMEFT

$$\mathcal{L}_{\text{DSMEFT}} \supset \sum_i c_i \mathcal{Q}_i^{(4)} + \boxed{\frac{1}{\Lambda} \sum_j c_j \mathcal{Q}_j^{(5)}} + \boxed{\frac{1}{\Lambda^2} \sum_k c_k \mathcal{Q}_k^{(6)}} + \boxed{\frac{1}{\Lambda^3} \sum_l c_l \mathcal{Q}_l^{(7)}}$$

$\mathcal{Q}_{u\phi^2} = (\bar{q}_p u_r \tilde{H}) \phi^2 + \text{h.c.}$	$\mathcal{Q}_{\phi u} = (\bar{u}_p \gamma_\mu u_r) (i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$	$\mathcal{Q}_{\phi q} = (\bar{q}_p \gamma_\mu q_r) (i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$	...	scalar
$\mathcal{Q}_{u\chi^2} = (\bar{q}_p u_r \tilde{H}) (\chi^T C \chi)$	$\mathcal{Q}_{u\chi} = (\bar{u}_p \gamma_\mu u_r) (\bar{\chi} \gamma^\mu \chi)$	$\mathcal{Q}_{q\chi} = (\bar{q}_p \gamma_\mu q_r) (\bar{\chi} \gamma^\mu \chi)$	...	fermion
$\mathcal{Q}_{qXX} = (\bar{q}_p \gamma_\mu q_r) X^{\mu\nu} X_\nu$	$\mathcal{Q}_{u\tilde{X}X} = (\bar{u}_p \gamma_\mu u_r) \tilde{X}^{\mu\nu} X_\nu$	$\mathcal{Q}_{uX^2} = (\bar{q}_p u_r \tilde{H}) X_\mu X^\mu + \text{h.c.}$	...	vector

► Matching





► Dark LEFT: **direct detection**



$$\mathcal{L}_{\text{DLEFT}} \supset \sum_k L_k \mathcal{O}_k^{(4)} + \boxed{\frac{1}{v} \sum_i L_i \mathcal{O}_i^{(5)}} + \boxed{\frac{1}{v^2} \sum_j L_j \mathcal{O}_j^{(6)}} + \boxed{\frac{1}{v^3} \sum_l L_l \mathcal{O}_l^{(7)}}$$

$\mathcal{O}_{g\phi} = G_{\mu\nu}^a G^{\mu\nu,a} \phi^2$	$\mathcal{O}_{d\phi^2} = (\bar{d}_p P_R d_r) \phi^2$	...	scalar
$\mathcal{O}_{g\chi} = G_{\mu\nu}^a G^{\mu\nu,a} \bar{\chi} \chi$	$\mathcal{O}_{d\chi}^{V,LR} = (\bar{d}_p \gamma_\mu P_L d_r) (\bar{\chi} \gamma^\mu \chi)$	...	fermion
$\mathcal{O}_{gX} = G_{\mu\nu}^a G^{\mu\nu,a} X_\rho X^\rho$	$\mathcal{O}_{dXX}^L = (\bar{d}_p \gamma^\mu P_L d_r) X_{\mu\nu} X^\nu$	...	vector

Hadron decay vs Direct detection





Hadron decay vs Direct detection



Conclusion

A Possible Flavour Path to Dark Matter

	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II/CEPC
<i>s</i>		DD	Belle II/CEPC
<i>b</i>			Belle II/LHC
	<i>u</i>	<i>c</i>	<i>t</i>
<i>u</i>	DD	BES/STCF	LHC/CEPC
<i>c</i>		BES/STCF	LHC/CEPC
<i>t</i>			LHC/CEPC



b-hadron FCNC decays with invisible final states is a possible path to New Physics, e.g., dark matter.

Many promising decay modes can be searched at **CEPC, FCC-ee** and Belle II

▶ $B_s \rightarrow \text{inv}$

▶ $\Lambda_b \rightarrow \Lambda + \text{inv}$

▶ $B_s \rightarrow \phi + \text{inv}$

▶ $B \rightarrow K^{(*)} + \text{inv}$

... ..

Thank You !

Backup

Summary

It's natural to ask:

1. By combing the direct detection, flavour measurements, and relic density, is it possible to obtain the **flavour structure** of interactions between light DM and SM particles ?

top-flavoured DM

2. Compared to DM direct detection, what's the advantage of hadron decays? which DM scenario is more sensitive to ? Can they provide **other information** ?

angular distribution (F_L) ;


HadronToNP: a package to calculate decay of hadron to new particles

$B \rightarrow K + \text{DM}, B \rightarrow \rho + \text{DM}, \Lambda_b \rightarrow \Lambda + \text{DM}, \Upsilon \rightarrow \text{DM}, \dots$

$D \rightarrow \pi + \text{DM}, D \rightarrow \rho + \text{DM}, \Xi_c \rightarrow \Xi + \text{DM}, J/\psi \rightarrow \text{DM}, \dots$

to be finished

many things to do, e.g., other flavour structures (MFV, $U(2)^5$), simplified model (light mediator), operator basis (SPVA), UV completion, full one-loop matching between DLEFT and DSMEFT, astro-constraints, ...



A Possible Flavour Path to Dark Matter

	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II
<i>s</i>		DD	Belle II
<i>b</i>			Belle II/LHC
	<i>u</i>	<i>c</i>	<i>t</i>
<i>u</i>	DD	BES/STCF	LHC/CEPC
<i>c</i>		BES/STCF	LHC/CEPC
<i>t</i>			LHC/CEPC

Dark SMEFT Operator

operator	definition	tree	loop	eq
$\mathcal{Q}_{u\phi}$	$(\bar{q}_p u_r \tilde{H})\phi + \text{h.c.}$			
$\mathcal{Q}_{u\phi^2}$	$(\bar{q}_p u_r \tilde{H})\phi^2 + \text{h.c.}$			
$\mathcal{Q}_{\phi q}$	$(\bar{q}_p \gamma_\mu q_r)(i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$			
$\mathcal{Q}_{\phi u}$	$(\bar{u}_p \gamma_\mu u_r)(i\phi_1 \overleftrightarrow{\partial}^\mu \phi_2)$			
$\mathcal{Q}_{q\chi}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{\chi} \gamma^\mu \chi)$			
$\mathcal{Q}_{u\chi}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{\chi} \gamma^\mu \chi)$			
$\mathcal{Q}_{u\chi^2}$	$(\bar{q}_p u_r \tilde{H})(\chi^T C \chi)$			

$\mathcal{Q}_{qX}$	$(\bar{q}_p \gamma^\mu q_r) X_\mu$			
$\mathcal{Q}_{uX}$	$(\bar{u}_p \gamma^\mu u_r) X_\mu$			
$\mathcal{Q}_{HqX}^{(1)}$	$(H^\dagger H)(\bar{q}_p \gamma^\mu q_r) X_\mu$			
$\mathcal{Q}_{HqX}^{(3)}$	$(H^\dagger \tau^I H)(\bar{q}_p \tau^I \gamma^\mu q_r) X_\mu$			
$\mathcal{Q}_{uX^2}$	$(\bar{q}_p u_r \tilde{H}) X_\mu X^\mu + \text{h.c.}$			
$\mathcal{Q}_{HuX}$	$(H^\dagger H)(\bar{u}_p \gamma^\mu u_r) X_\mu$			
$\mathcal{Q}_{qXX}$	$(\bar{q}_p \gamma_\mu q_r) X^{\mu\nu} X_\nu$			
$\mathcal{Q}_{q\tilde{X}X}$	$(\bar{q}_p \gamma_\mu q_r) \tilde{X}^{\mu\nu} X_\nu$			
$\mathcal{Q}_{uXX}$	$(\bar{u}_p \gamma_\mu u_r) X^{\mu\nu} X_\nu$			
$\mathcal{Q}_{u\tilde{X}X}$	$(\bar{u}_p \gamma_\mu u_r) \tilde{X}^{\mu\nu} X_\nu$			
$\mathcal{Q}_{DqX^2}$	$(\bar{q}_p \gamma^\nu i D^\mu q_r) X_\mu X_\nu + \text{h.c.}$			
$\mathcal{Q}_{DuX^2}$	$(\bar{u}_p \gamma^\nu i D^\mu u_r) X_\mu X_\nu + \text{h.c.}$			
$\mathcal{Q}_{uHX^2}$	$(\bar{q}_p \sigma_{\mu\nu} u_r \tilde{H}) X_1^\mu X_2^\nu + \text{h.c.}$			
$\mathcal{Q}_{uHX}$	$(\bar{q}_p \sigma_{\mu\nu} u_r) \tilde{H} X^{\mu\nu} + \text{h.c.}$			

Dark LEFT Operator

operator	definition	tree	loop
$\mathcal{O}_{d\phi}$	$(\bar{d}_{Lp}d_{Rr})\phi + \text{h.c.}$	—	$\mathcal{Q}_{u\phi}$
$\mathcal{O}_{d\phi^2}$	$(\bar{d}_{Lp}d_{Rr})\phi^2 + \text{h.c.}$	—	$\mathcal{Q}_{u\phi^2}$
$\mathcal{O}_{\phi d}^L$	$(\bar{d}_{Lp}\gamma_\mu d_{Lr})(i\phi_1\overleftrightarrow{\partial}^\mu\phi_2)$	$\mathcal{Q}_{\phi q}$	$\mathcal{Q}_{\phi u}$
$\mathcal{O}_{\phi d}^R$	$(\bar{d}_{Rp}\gamma_\mu d_{Rr})(i\phi_1\overleftrightarrow{\partial}^\mu\phi_2)$	—	—
$\mathcal{O}_{d\chi}^{S,LR}$	$(\bar{d}_{Rp}d_{Lr})(\chi^TC\chi) + \text{h.c.}$	—	$\mathcal{Q}_{u\chi^2}$
$\mathcal{O}_{d\chi}^{S,RR}$	$(\bar{d}_{Lp}d_{Rr})(\chi^TC\chi) + \text{h.c.}$	—	$\mathcal{Q}_{u\chi^2}$
$\mathcal{O}_{d\chi}^{V,LR}$	$(\bar{d}_{Lp}\gamma_\mu d_{Lr})(\bar{\chi}\gamma^\mu\chi)$	$\mathcal{Q}_{q\chi}$	$\mathcal{Q}_{u\chi}$
$\mathcal{O}_{d\chi}^{V,RR}$	$(\bar{d}_{Rp}\gamma_\mu d_{Rr})(\bar{\chi}\gamma^\mu\chi)$	—	—
$\mathcal{O}_{d\chi}^{T,RR}$	$(\bar{d}_{Lp}\sigma^{\mu\nu}d_{Rr})(\chi^TC\sigma_{\mu\nu}\chi) + \text{h.c.}$	—	—
$\mathcal{O}_{dX}^T$	$(\bar{d}_{Lp}\sigma^{\mu\nu}d_{Rr})X_{\mu\nu} + \text{h.c.}$	—	$\mathcal{Q}_{uX}, \mathcal{Q}_{qX}, \mathcal{Q}_{uHX}, \mathcal{Q}_{HuX}, \mathcal{Q}_{HqX}$
$\mathcal{O}_{dX}^L$	$(\bar{d}_{Lp}\gamma^\mu d_{Lr})X_\mu$	$\mathcal{Q}_{qX}, \mathcal{Q}_{HqX}$	$\mathcal{Q}_{uX}, \mathcal{Q}_{HuX}$
$\mathcal{O}_{dX}^R$	$(\bar{d}_{Rp}\gamma^\mu d_{Rr})X_\mu$	—	—
$\mathcal{O}_{dX^2}$	$(\bar{d}_{Lp}d_{Rr})X_\mu X^\mu + \text{h.c.}$	—	$\mathcal{Q}_{uX^2}, \mathcal{Q}_{DuX^2}, \mathcal{Q}_{DqX^2}$
$\mathcal{O}_{dX^2}^T$	$(\bar{d}_{Lp}\sigma_{\mu\nu}d_{Rr})X_1^\mu X_2^\nu + \text{h.c.}$	—	$\mathcal{Q}_{uHX^2}$
$\mathcal{O}_{dXX}^L$	$(\bar{d}_{Lp}\gamma^\mu d_{Lr})X_{\mu\nu}X^\nu$	$\mathcal{Q}_{qXX}$	$\mathcal{Q}_{uXX}$
$\mathcal{O}_{dXX}^R$	$(\bar{d}_{Rp}\gamma^\mu d_{Rr})X_{\mu\nu}X^\nu$	—	—
$\mathcal{O}_{d\tilde{X}X}^L$	$(\bar{d}_{Lp}\gamma^\mu d_{Lr})\tilde{X}_{\mu\nu}X^\nu$		
$\mathcal{O}_{d\tilde{X}X}^R$	$(\bar{d}_{Rp}\gamma^\mu d_{Rr})\tilde{X}_{\mu\nu}X^\nu$		
$\mathcal{O}_{DdX^2}^L$	$(\bar{d}_{Lp}\gamma^\mu iD^\nu d_{Lr})X_\mu X_\nu + \text{h.c.}$	$\mathcal{Q}_{DqX^2}$	$\mathcal{Q}_{DuX^2}$
$\mathcal{O}_{DdX^2}^R$	$(\bar{d}_{Rp}\gamma^\mu iD^\nu d_{Rr})X_\mu X_\nu + \text{h.c.}$	—	—

$\mathcal{O}_{da}^L$	$(\bar{d}_{Lp}\gamma_\mu d_{Lr})\partial^\mu a$	$\mathcal{Q}_{qa}$	$\mathcal{Q}_{ua}$
$\mathcal{O}_{da}^R$	$(\bar{d}_{Rp}\gamma_\mu d_{Rr})\partial^\mu a$	—	—
$\mathcal{O}_{GG\phi^2}$	$G_{\mu\nu}^a G^{\mu\nu,a}\phi^2 (\mathcal{O}_{g\phi})$	—	$\mathcal{Q}_{u\phi^2}$
$\mathcal{O}_{G\tilde{G}\phi^2}$	$G_{\mu\nu}^a \tilde{G}^{\mu\nu,a}\phi^2 (\mathcal{O}_{g\phi})$	—	$\mathcal{Q}_{u\phi^2}$
$\mathcal{O}_{GG\chi}$	$G_{\mu\nu}^a G^{\mu\nu,a}\chi^TC\chi (\mathcal{O}_{g\chi})$	—	$\mathcal{Q}_{u\chi^2}$
$\mathcal{O}_{G\tilde{G}\chi}$	$G_{\mu\nu}^a \tilde{G}^{\mu\nu,a}\chi^TC\chi (\mathcal{O}_{g\chi})$	—	$\mathcal{Q}_{u\chi^2}$
$\mathcal{O}_{GGXX}$	$G_{\mu\nu}^a G^{\mu\nu,a}X_\rho X^\rho (\mathcal{O}_{gX})$	—	$\mathcal{Q}_{uX^2}$
$\mathcal{O}_{G\tilde{G}XX}$	$G_{\mu\nu}^a \tilde{G}^{\mu\nu,a}X_\rho X^\rho (\mathcal{O}_{gX})$	—	$\mathcal{Q}_{uX^2}$
$\mathcal{O}_{GXGX}$	$G_{\lambda\mu}^a G_{\nu}^{\lambda,a}X^\mu X^\nu (\mathcal{O}_{gXX})$	—	$\mathcal{Q}_{DqX^2}, \mathcal{Q}_{DuX^2}$
$\mathcal{O}_{GX\tilde{G}X}$	$G_{\lambda\mu}^a \tilde{G}_{\nu}^{\lambda,a}X^\mu X^\nu (\mathcal{O}_{gXX})$		
$\mathcal{O}_{FF\phi^2}$	$F_{\mu\nu}F^{\mu\nu}\phi^2 (\mathcal{O}_{\phi\gamma})$	—	$\mathcal{Q}_{u\phi^2}$
$\mathcal{O}_{F\tilde{F}\phi^2}$	$F_{\mu\nu}\tilde{F}^{\mu\nu}\phi^2 (\mathcal{O}_{\phi\gamma})$		
$\mathcal{O}_{FF\chi}$	$F_{\mu\nu}F^{\mu\nu}\chi^TC\chi (\mathcal{O}_{\chi\gamma})$	—	$\mathcal{Q}_{u\chi}$
$\mathcal{O}_{F\tilde{F}\chi}$	$F_{\mu\nu}\tilde{F}^{\mu\nu}\chi^TC\chi (\mathcal{O}_{\chi\gamma})$		
$\mathcal{O}_{FFXX}$	$F_{\mu\nu}F^{\mu\nu}X_\rho X^\rho (\mathcal{O}_{X\gamma})$	—	$\mathcal{Q}_{uX^2}$
$\mathcal{O}_{F\tilde{F}XX}$	$F_{\mu\nu}\tilde{F}^{\mu\nu}X_\rho X^\rho (\mathcal{O}_{X\gamma})$		
$\mathcal{O}_{FXFX}$	$F_{\lambda\mu}F_{\nu}^{\lambda}X^\mu X^\nu (\mathcal{O}_{XX\gamma})$	—	$\mathcal{Q}_{DqX^2}, \mathcal{Q}_{DuX^2}$
$\mathcal{O}_{FX\tilde{F}X}$	$F_{\lambda\mu}\tilde{F}_{\nu}^{\lambda}X^\mu X^\nu (\mathcal{O}_{XX\gamma})$		

$$\mathcal{C} = \begin{cases} \mathcal{C}_t & \text{for } \bar{u}_L \gamma^\mu \mathcal{C} u_L, u_R \gamma^\mu \mathcal{C} u_R, \bar{u}_L \mathcal{C} u_R, \bar{u}_L \sigma^{\mu\nu} \mathcal{C} u_R, \\ V^\dagger \mathcal{C}_t V & \text{for } \bar{d}_L \gamma^\mu \mathcal{C} d_L, \\ V^\dagger \mathcal{C}_t & \text{for } \bar{d}_L \mathcal{C} u_R, \bar{d}_L \sigma^{\mu\nu} \mathcal{C} u_R, \end{cases}$$

$$V^\dagger \mathcal{C}_t V = \mathcal{C}_{33} \begin{pmatrix} V_{td}^* V_{td} & V_{td}^* V_{ts} & V_{td}^* V_{tb} \\ V_{ts}^* V_{td} & V_{ts}^* V_{ts} & V_{ts}^* V_{tb} \\ V_{tb}^* V_{td} & V_{tb}^* V_{ts} & V_{tb}^* V_{tb} \end{pmatrix} \quad \mathcal{C}_t \equiv \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \mathcal{C}_{33} \end{pmatrix}$$



$$= \mathcal{C}_{33} \begin{pmatrix} 0.71 & -3.09 - 1.45i & 76.81 + 34.29i \\ -3.09 + 1.45i & 16.43 & -404.93 + 7.87i \\ 76.81 - 34.29i & -404.93 - 7.87i & 9982.86 \end{pmatrix} \times 10^{-4}$$

$$V^\dagger \mathcal{C}_t = \mathcal{C}_{33} \begin{pmatrix} 0 & 0 & V_{td}^* \\ 0 & 0 & V_{ts}^* \\ 0 & 0 & V_{tb}^* \end{pmatrix} = \mathcal{C}_{33} \begin{pmatrix} 0 & 0 & 0.77 + 0.34i \\ 0 & 0 & -4.05 + 0.08i \\ 0 & 0 & 99.91 \end{pmatrix} \times 10^{-2}$$

$$\mathcal{C}_i = \begin{cases} \mathcal{C}_t^i & \text{for } \mathcal{Q}_i = \mathcal{Q}_{\phi u}, \mathcal{Q}_{u\chi}, \mathcal{Q}_{uX}, \mathcal{Q}_{HuX}, \mathcal{Q}_{uXX}, \mathcal{Q}_{u\tilde{X}X}, \mathcal{Q}_{DuX^2}, \mathcal{Q}_{ua}, \\ (\mathcal{C}_t^i, V^\dagger \mathcal{C}_t^i) & \text{for } \mathcal{Q}_i = \mathcal{Q}_{u\phi}, \mathcal{Q}_{u\phi^2}, \mathcal{Q}_{u\chi^2}, \mathcal{Q}_{qu\chi l}, \mathcal{Q}_{uHX}, \mathcal{Q}_{uX^2}, \mathcal{Q}_{uHX^2}, \\ (\mathcal{C}_t^i, V^\dagger \mathcal{C}_t^i V) & \text{for } \mathcal{Q}_i = \mathcal{Q}_{\phi q}, \mathcal{Q}_{q\chi}, \mathcal{Q}_{qX}, \mathcal{Q}_{HqX}^{(1,3)}, \mathcal{Q}_{qXX}, \mathcal{Q}_{q\tilde{X}X}, \mathcal{Q}_{DqX^2}, \mathcal{Q}_{qa}, \end{cases}$$

Conclusion

HadronToNP: a package to calculate decay of hadron to new particles
 $B \rightarrow K + \text{DM}, B \rightarrow \rho + \text{DM}, \Lambda_b \rightarrow \Lambda + \text{DM}, \Upsilon \rightarrow \text{DM}, \dots$ *to be finished*
 $D \rightarrow \pi + \text{DM}, D \rightarrow \rho + \text{DM}, \Xi_c \rightarrow \Xi + \text{DM}, J/\psi \rightarrow \text{DM}, \dots$



SMEFT

Dark SMEFT

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}}} = 0.46 \pm 0.07$$
$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}}} = 29.7 \pm 5.6$$

- Belle II excess (if confirmed in the future) implies:
- impossible to explain in SMEFT with MFV
 - NP flavour structure is highly non-trivial
 - **NP structure in quark sector is beyond MFV**
 - **flavour violation is beyond Yukawa coupling**

A Possible Flavour Path to Dark Matter

	<i>d</i>	<i>s</i>	<i>b</i>
<i>d</i>	DD	NA62/KOTO	Belle II
<i>s</i>		DD	Belle II
<i>b</i>			Belle II/LHC
<i>u</i>			
<i>c</i>			
<i>t</i>			
<i>u</i>	DD	BES/STCF	LHC/CEPC
<i>c</i>		BES/STCF	LHC/CEPC
<i>t</i>			LHC/CEPC


μ_{EW}
LEFT

Dark LEFT


μ_b

$b \rightarrow s\nu\bar{\nu}$: exp & theory

► 2021 Apr



► 2023 Aug




► Exp vs SM [10⁻⁶]

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}} = 4.16 \pm 0.57$$

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{exp}} = 23 \pm 7$$

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{exp}} \gtrsim 10 \text{ (} 2\sigma \text{ lower bound)}$$

2.7σ difference
NP/SM ≳ 2



► Theoretical prediction

Factorization

$$\mathcal{A} \propto C_L \cdot \langle K | \bar{s} \gamma^\mu b | \bar{B} \rangle \cdot \bar{\nu} \gamma_\mu \nu$$

Wilson coef quark current neutrino current

theoretically, simple and clean
one of the cleanest channels in
flavour physics

$$\mathcal{O}_L = (\bar{s} \gamma_\mu P_L b)(\bar{\nu} \gamma^\mu P_L \nu) \text{ in the SM}$$

$$\mathcal{O}_L = (\bar{s} P_L b)(\bar{\nu} P_L \nu) \times$$

$$\mathcal{O}_R = (\bar{s} \gamma_\mu P_R b)(\bar{\nu} \gamma^\mu P_L \nu) \text{ possible in BSM}$$

$$\mathcal{O}_R = (\bar{s} P_R b)(\bar{\nu} P_R \nu) \times$$

$$\mathcal{O}_T = (\bar{s} \sigma_{\mu\nu} b)(\bar{\nu} \sigma^{\mu\nu} \nu) \times$$

operator structure highly
constrained by LH neutrino

$$\mathcal{O}_{T5} = (\bar{s} \sigma_{\mu\nu} \gamma_5 b)(\bar{\nu} \sigma^{\mu\nu} \nu) \times$$

Evidence for $B^+ \rightarrow K^+ \nu \bar{\nu}$ decays

Belle-II Collaboration • I. Adachi et al. (Nov 24, 2023)

Published in: *Phys.Rev.D* 109 (2024) 11, 112006 • e-Print: [2311.14647](https://arxiv.org/abs/2311.14647) [hep-ex]

pdf DOI cite claim

reference search

83 citations

$b \rightarrow s\nu\bar{\nu}$: exp & theory

$b \rightarrow s$

Observable	SM	Exp	Unit
$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}

$b \rightarrow d$

$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}

$s \rightarrow d$

$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}
$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}

Why such a large NP effect has not shown up
in other $b \rightarrow s$ decays ?
in $b \rightarrow d, s \rightarrow d$ decays ?

► Exp vs SM [10⁻⁶]


$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}} = 4.16 \pm 0.57$$

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{exp}} = 23 \pm 7$$

$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{exp}} \gtrsim 10 \text{ (2}\sigma \text{ lower bound)}$$

}

2.7σ difference
 NP/SM ≳ 2



► Theoretical prediction

Factorization

$$\mathcal{A} \propto C_L \cdot \langle K | \bar{s} \gamma^\mu b | \bar{B} \rangle \cdot \bar{\nu} \gamma_\mu \nu$$

Wilson coef
 quark current
 neutrino current

theoretically, simple and clean
 one of the cleanest channels in
 flavour physics

$$\mathcal{O}_L = (\bar{s} \gamma_\mu P_L b)(\bar{\nu} \gamma^\mu P_L \nu)$$
 in the SM

$$\mathcal{O}_R = (\bar{s} \gamma_\mu P_R b)(\bar{\nu} \gamma^\mu P_L \nu)$$
 possible in BSM

$$\mathcal{O}_L = (\bar{s} P_L b)(\bar{\nu} P_L \nu) \quad \times$$

$$\mathcal{O}_R = (\bar{s} P_R b)(\bar{\nu} P_R \nu) \quad \times$$

$$\mathcal{O}_T = (\bar{s} \sigma_{\mu\nu} b)(\bar{\nu} \sigma^{\mu\nu} \nu) \quad \times$$

$$\mathcal{O}_{T5} = (\bar{s} \sigma_{\mu\nu} \gamma_5 b)(\bar{\nu} \sigma^{\mu\nu} \nu) \quad \times$$

operator structure highly
 constrained by LH neutrino

$b \rightarrow s\nu\bar{\nu}$: exp & theory

	Observable	SM	Exp	Unit
$b \rightarrow s$	$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5^{+5}_{-4}$	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
	$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
	$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
	$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
	$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}
$b \rightarrow d$	$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
	$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
	$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}
$s \rightarrow d$	$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6^{+4.0}_{-3.4} \pm 0.9$	10^{-11}
	$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}

Why such a large NP effect has not shown up in other $b \rightarrow s$ decays ?
in $b \rightarrow d, s \rightarrow d$ decays ? **NP flavour structure**

SMEFT

$$\mathcal{Q}_{Hq}^{(1)} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r) \, ,$$

$$\mathcal{Q}_{Hq}^{(3)} = (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r) \, ,$$

$$\mathcal{Q}_{Hd} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r) \, ,$$

$$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t) \, ,$$

$$\mathcal{Q}_{lq}^{(1)} = (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t) \, ,$$

$$\mathcal{Q}_{lq}^{(3)} = (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t) \, ,$$

μ_{EW}

LEFT

$$\mathcal{O}_L^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

$$\mathcal{O}_R^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

μ_b

operator structure highly constrained by Left-handed neutrino

Minimal Flavour Violation

- ▶ Flavour symmetry without Yukawa

$$G_{\text{QF}} = SU(3)_q \otimes SU(3)_u \otimes SU(3)_d$$

- ▶ Flavour symmetry breaking only from SM Yukawa

$$-\mathcal{L}_Y = \bar{q} Y_d H d + \bar{q} Y_u \tilde{H} u + \text{h.c.}$$

- ▶ Flavour symmetry recovering: Yukawa coupling $\implies$ spurion field

$$Y_u \sim (\mathbf{3}, \bar{\mathbf{3}}, 1) \quad Y_d \sim (\mathbf{3}, 1, \bar{\mathbf{3}})$$

D'Ambrosio, Giudice, Isidori, Strumia, 2009

- ▶ EFT with MFV: operators, constructed from SM and Yukawa spurion fields, are invariant under CP and G_{QF}

$$\mathcal{C}^{\text{MFV}} = \begin{cases} f(A, B) & \text{for } \bar{q} \gamma^\mu \mathcal{C} q, \\ f(A, B) Y_d & \text{for } \bar{q} \mathcal{C} d, \bar{q} \sigma^{\mu\nu} \mathcal{C} d, \\ \epsilon_0 \mathbb{1} + Y_d^\dagger g(A, B) Y_d & \text{for } \bar{d} \gamma^\mu \mathcal{C} d, \end{cases} \quad \begin{aligned} f(A, B) &= \epsilon_0 \mathbb{1} + \epsilon_1 A + \epsilon_2 B + \epsilon_3 A^2 + \epsilon_4 B^2 + \epsilon_5 AB + \dots \\ A &= Y_u Y_u^\dagger \\ B &= Y_d Y_d^\dagger \end{aligned}$$

Minimal Flavour Violation

- ▶ Spurion function

$$f(A, B) = \epsilon_0 \mathbb{1} + \epsilon_1 A + \epsilon_2 B + \epsilon_3 A^2 + \epsilon_4 B^2 + \epsilon_5 AB + \dots$$

- ▶ Cayley-Hamilton identity for 3×3 invertible matrix X

$$X^3 = \text{Det}X \cdot \mathbb{1} + \frac{1}{2}[\text{Tr}X^2 - (\text{Tr}X)^2] \cdot X + \text{Tr}X \cdot X^2$$

- ▶ Spurion function after resummation

Colangelo, Nikolidakis, Smith, 2009
Mercolli, Smith, 2009

$$f(A, B) = \epsilon_0 \mathbb{1} + \epsilon_1 A + \epsilon_3 A^2 + \epsilon_5 AB + \epsilon_7 ABA + \epsilon_{10} AB^2 + \epsilon_{12} A^2 B^2 + \epsilon_{14} B^2 AB + \epsilon_{15} AB^2 A^2 \\ + \epsilon_2 B + \epsilon_4 B^2 + \epsilon_6 BA + \epsilon_9 BAB + \epsilon_8 BA^2 + \epsilon_{13} B^2 A^2 + \epsilon_{11} ABA^2 + \epsilon_{16} B^2 A^2 B.$$

- ▶ assumption #1: neglect tiny imaginary parts of ϵ_i
- ▶ assumption #2: neglect spurion B (suppressed by $\mathcal{O}(\lambda_d^2)$)

$$f(A, B) \approx \epsilon_0 \mathbb{1} + \epsilon_1 A + \epsilon_2 A^2$$

Minimal Flavour Violation

- ▶ MFV coupling FCNC controlled by CKM

$$C^{\text{MFV}} = \begin{cases} \epsilon_0 1 + \epsilon_1 \Delta_q & \text{for } \bar{d}_L \gamma^\mu C d_L \\ \epsilon_0 \hat{\lambda}_d + \epsilon_1 \Delta_q \hat{\lambda}_d & \text{for } \bar{d}_L C d_R, \bar{d}_L \sigma^{\mu\nu} C d_R \\ \epsilon_0 1 & \text{for } \bar{d}_R \gamma^\mu C d_R \end{cases} \quad \Delta_q = V^\dagger \hat{\lambda}_u^2 V$$

No Right-handed down-type FCNC !

- ▶ Numerics

$$\Delta_q = \begin{pmatrix} 0.8 & -3.3 - 1.5i & 79.3 + 35.4i \\ -3.3 + 1.5i & 16.6 & -397.5 + 8.1i \\ 79.3 - 35.4i & -397.5 - 8.1i & 9839.0 \end{pmatrix} \times 10^{-4}$$

$$\Delta_q \hat{\lambda}_d = \begin{pmatrix} 0.0021 & -0.18 - 0.08i & 191.3 + 85.4i \\ -0.009 + 0.004i & 0.88 & -958.7 + 19.6i \\ 0.21 - 0.10i & -21.1 - 0.4i & 23728.1 \end{pmatrix} \times 10^{-6}$$

$b \rightarrow s\nu\bar{\nu}$: SMEFT with MFV

► Prediction

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}}} = 0.46 \pm 0.07$$

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}}} = 29.7 \pm 5.6$$

► prediction

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}} = (9.00 \pm 0.87) \times 10^{-6}$$

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{MFV}} = (50^{+17}_{-16}) \times 10^{-6}$$

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{exp}} < 18 \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} = (1.40 \pm 0.18) \times 10^{-7}$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{MFV}} = (7.8^{+2.8}_{-2.6}) \times 10^{-7}$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{exp}} < 140 \times 10^{-7}$$

SMEFT

$$\mathcal{Q}_{Hq}^{(1)} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r),$$

$$\mathcal{Q}_{Hq}^{(3)} = (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r),$$

$$\mathcal{Q}_{Hd} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r),$$

induce $\bar{s}bZ$ interaction,
Thus, universally affect
 $b \rightarrow se^+e^-, \mu^+\mu^-, \tau^+\tau^-$

$$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t),$$

forbidden by MFV

$$\mathcal{Q}_{lq}^{(1)} = (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t),$$

$$\mathcal{Q}_{lq}^{(3)} = (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t),$$

μ_{EW}

LEFT

μ_b

$$\mathcal{O}_L^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

$$\mathcal{O}_R^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

one LEFT operator !
just the SM operator

$b \rightarrow s\nu\bar{\nu}$: SMEFT with MFV

► Prediction

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}}} = 0.46 \pm 0.07$$

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}}} = 29.7 \pm 5.6$$

► prediction

$$\left. \begin{aligned} \mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}} &= (9.00 \pm 0.87) \times 10^{-6} \\ \mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{MFV}} &= (50^{+17}_{-16}) \times 10^{-6} \\ \mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{exp}} &< 18 \times 10^{-6} \end{aligned} \right\} \text{Inconsistent} \longrightarrow$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} = (1.40 \pm 0.18) \times 10^{-7}$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{MFV}} = (7.8^{+2.8}_{-2.6}) \times 10^{-7}$$

$$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{exp}} < 140 \times 10^{-7}$$

Belle II excess (if confirmed in the future) implies:

- impossible to explain in SMEFT with MFV
- NP flavour structure is highly non-trivial
- **NP structure in quark sector is beyond MFV**
- **flavour violation is beyond Yukawa coupling**

This conclusion only assumes the quark MFV.
No lepton flavour structure is assumed.

$b \rightarrow s\nu\bar{\nu}$: SMEFT

SMEFT

$$\mathcal{Q}_{Hq}^{(1)} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r),$$

$$\mathcal{Q}_{Hq}^{(3)} = (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r),$$

$$\mathcal{Q}_{Hd} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r),$$

$$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t),$$

$$\mathcal{Q}_{lq}^{(1)} = (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t),$$

$$\mathcal{Q}_{lq}^{(3)} = (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t),$$

μ_{EW}


LEFT

$$\mathcal{O}_L^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

$$\mathcal{O}_R^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

μ_b

operator structure highly
constrained by Left-handed neutrino



$$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu}) = A_+^{BK} x^+,$$

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu}) = A_+^{BK^*} x^+ + A_-^{BK^*} x^-,$$

$$x^\pm = \sum_{\nu, \nu'} |C_L^{\nu \nu'} \pm C_R^{\nu \nu'}|^2,$$

Bause, Gisbert, Hiller, 2309.00075

Allwicher, Becirevic, Piazza, Rosauero-Alcaraz, Sumensari, 2309.02246

Chen, Wen, Xu, 2401.11552

$b \rightarrow s\nu\bar{\nu}$: SMEFT

SMEFT	$\mathcal{Q}_{Hq}^{(1)} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r),$
	$\mathcal{Q}_{Hq}^{(3)} = (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r),$
	$\mathcal{Q}_{Hd} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r),$
	$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t),$
	$\mathcal{Q}_{lq}^{(1)} = (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t),$
μ_{EW}	$\mathcal{Q}_{lq}^{(3)} = (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t),$
	$\mathcal{O}_L^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$
	$\mathcal{O}_R^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$
LEFT	
μ_b	operator structure highly constrained by Left-handed neutrino

$b \rightarrow s$

Observable	SM	Exp	Unit
$\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})$	4.16 ± 0.57	$23 \pm 5_{-4}^{+5}$	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^0 \nu \bar{\nu})$	3.85 ± 0.52	< 26	10^{-6}
$\mathcal{B}(B^+ \rightarrow K^{*+} \nu \bar{\nu})$	9.70 ± 0.94	< 61	10^{-6}
$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})$	9.00 ± 0.87	< 18	10^{-6}
$\mathcal{B}(B_s \rightarrow \phi \nu \bar{\nu})$	9.93 ± 0.72	< 5400	10^{-6}
$\mathcal{B}(B_s \rightarrow \nu \bar{\nu})$	≈ 0	< 5.9	10^{-4}
$\mathcal{B}(B^+ \rightarrow \pi^+ \nu \bar{\nu})$	1.40 ± 0.18	< 140	10^{-7}
$\mathcal{B}(B^0 \rightarrow \pi^0 \nu \bar{\nu})$	6.52 ± 0.85	< 900	10^{-8}
$\mathcal{B}(B^+ \rightarrow \rho^+ \nu \bar{\nu})$	4.06 ± 0.79	< 300	10^{-7}
$\mathcal{B}(B^0 \rightarrow \rho^0 \nu \bar{\nu})$	1.89 ± 0.36	< 400	10^{-7}
$\mathcal{B}(B^0 \rightarrow \nu \bar{\nu})$	≈ 0	< 1.4	10^{-4}
$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	8.42 ± 0.61	$10.6_{-3.4}^{+4.0} \pm 0.9$	10^{-11}
$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	3.41 ± 0.45	< 300	10^{-11}

$b \rightarrow d$

$s \rightarrow d$

Why such a large NP effect has not shown up in other $b \rightarrow s$ decays ?
in $b \rightarrow d, s \rightarrow d$ decays ? **NP flavour structure**

$b \rightarrow s\nu\bar{\nu}$ and $b \rightarrow s\ell\ell$

SMEFT notation: $l = \begin{pmatrix} \nu \\ e \end{pmatrix}_L$, $q = \begin{pmatrix} u \\ d \end{pmatrix}_L$, $d = d_R$

B.F.Hou, X.Q.Li, M.Shen, Y.D.Yang, **XB**, 2402.19208

► Prediction

$$\frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})} = \frac{\mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}}{\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}}} = 0.46 \pm 0.07$$

► prediction

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SM}} = (9.00 \pm 0.87) \times 10^{-6}$$

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{SMEFT}} = (50^{+17}_{-16}) \times 10^{-6} \quad \text{conflict}$$

$$\mathcal{B}(B^0 \rightarrow K^{*0} \nu \bar{\nu})_{\text{exp}} < 18 \times 10^{-6}$$

► Only $\mathcal{O}_{lq}^{(3)}$ is relevant with $R_{D^{(*)}}$

► $\mathcal{O}_{ld}$ can explain the $B^+ \rightarrow K^+ \nu \bar{\nu}$ data

► $\mathcal{O}_{ld}$ also induce $O'_{9,ij}$ and $O'_{10,ij}$

► They can't improve the $b \rightarrow s\ell\ell$ fit

► O'_{9e} and $O'_{10\mu}$ worsen the fit. **weird** (LFV, $\tau\tau \gg ee, \mu\mu$)

► $O'_{9,ij}$ and $O'_{10,ij}$ with $i = j = \tau$ has no effect.

► $O'_{9,ij}$ and $O'_{10,ij}$ with $i \neq j$ (i.e. LFV) has no effect.

SMEFT

$$\begin{aligned} \mathcal{Q}_{Hq}^{(1)} &= (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r), \\ \mathcal{Q}_{Hq}^{(3)} &= (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r), \\ \mathcal{Q}_{Hd} &= (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r), \end{aligned}$$

$$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t),$$

$$\begin{aligned} \mathcal{Q}_{lq}^{(1)} &= (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t), \\ \mathcal{Q}_{lq}^{(3)} &= (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t), \end{aligned}$$

induce $\bar{s}bZ$ interaction,
Thus, universally affect
 $b \rightarrow se^+e^-, \mu^+\mu^-, \tau^+\tau^-$

μ_{EW}

LEFT

$$\mathcal{O}_L^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$

$$\mathcal{O}_R^{\nu_i \nu_j} = (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j)$$



one LEFT operator !
just the SM operator

μ_b

$$O'_{9,ij} = (\bar{b} \gamma^\mu P_R s) (\bar{\ell}_i \gamma_\mu \ell_j)$$

$$O'_{10,ij} = (\bar{b} \gamma^\mu P_R s) (\bar{\ell}_i \gamma_\mu \gamma_5 \ell_j)$$

$b \rightarrow s\nu\bar{\nu}$: exp picture



$$\begin{aligned} \mathcal{Q}_{Hq}^{(1)} &= (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{q}_p \gamma^\mu q_r), \\ \mathcal{Q}_{Hq}^{(3)} &= (H^\dagger i \overleftrightarrow{D}_\mu^I H) (\bar{q}_p \tau^I \gamma^\mu q_r), \\ \mathcal{Q}_{Hd} &= (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{d}_p \gamma^\mu d_r), \end{aligned}$$

$$\mathcal{Q}_{ld} = (\bar{l}_p \gamma^\mu l_r) (\bar{d}_s \gamma_\mu d_t),$$

$$\begin{aligned} \mathcal{Q}_{lq}^{(1)} &= (\bar{l}_p \gamma^\mu l_r) (\bar{q}_s \gamma_\mu q_t), \\ \mathcal{Q}_{lq}^{(3)} &= (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{q}_s \tau^I \gamma_\mu q_t), \end{aligned}$$

$$\begin{aligned} \mathcal{O}_L^{\nu_i \nu_j} &= (\bar{s} \gamma_\mu P_L b) (\bar{\nu}_i \gamma^\mu P_L \nu_j) \\ \mathcal{O}_R^{\nu_i \nu_j} &= (\bar{s} \gamma_\mu P_R b) (\bar{\nu}_i \gamma^\mu P_L \nu_j) \end{aligned}$$

SMEFT

Dark SMEFT

example

$$\mathcal{Q}_{d\phi^2} = (\bar{q}_p d_r H) \phi^2$$

$$\mathcal{Q}_{d\chi} = (\bar{d}_p \gamma_\mu d_r) (\bar{\chi} \gamma^\mu \chi)$$

$$\mathcal{Q}_{dX^2} = (\bar{q}_p d_r H) X_\mu X^\mu$$

$$\mathcal{Q}_{qa} = (\bar{q}_p \gamma_\mu q_r) \partial^\mu a$$

2011 Kamenik, Smith

2014 Duch, Grzadkowski, Wudka

2017 Brod, Gootjes-Dreesbach, Tamaro, Zupan

2021 Criado, Djouadi, Perez-Victoria, Santiago

2022 Aebischer, Altmannshofer, Jenkins, Manohar ([basis@dim-6](#))

2023 Song, Sun, Yu ([basis@dim-8](#))

Axion-like particle,

see also H.Y.Cheng, Phys.Rept 1988

2020 Bauer, Neubert, Renner, Schnubel, Thamm

2023 Song, Sun, Yu ([basis@dim-8](#))

μ_{EW}
LEFT

Dark LEFT

$$\mathcal{O}_{d\phi^2} = (\bar{d}_{Lp} d_{Rr}) \phi^2$$

$$\mathcal{O}_{d\chi}^{V,LR} = (\bar{d}_{Lp} \gamma_\mu d_{Lr}) (\bar{\chi}_a \gamma^\mu \chi_b)$$

$$\mathcal{O}_{dXX}^L = (\bar{d}_{Lp} \gamma_\mu d_{Lr}) X^{\mu\nu} X_\nu$$

$$\mathcal{O}_{da}^L = (\bar{d}_{Lp} \gamma_\mu d_{Lr}) \partial^\mu a$$

example

2022 Aebischer, Altmannshofer, Jenkins, Manohar ([basis@dim-6](#))

2022 He, Ma, Valencia ([basis@dim-6](#))

2023 Liang, Liao, Ma, Wang ([basis@dim-8](#))

μ_b

Dark SMEFT with MFV

- ▶ MFV coupling $b \rightarrow s, b \rightarrow d, s \rightarrow d$ are connected with each other.

$$c_i^{\text{MFV}} = \begin{cases} \epsilon_0^i \hat{\lambda}_d + \epsilon_1^i \Delta_q \hat{\lambda}_d & \text{for } \mathcal{Q}_i = \mathcal{Q}_{d\phi}, \mathcal{Q}_{d\phi^2}, \mathcal{Q}_{dHX}, \mathcal{Q}_{dHX^2}, \mathcal{Q}_{dX^2}, \\ \epsilon_0^i \mathbb{1} + \epsilon_1^i \Delta_q & \text{for } \mathcal{Q}_i = \mathcal{Q}_{\phi q}, \mathcal{Q}_{q\chi}, \mathcal{Q}_{qXX}, \mathcal{Q}_{q\tilde{X}X}, \mathcal{Q}_{DqX^2}, \mathcal{Q}_{qX}, \mathcal{Q}_{HqX}^{(1,3)}, \mathcal{Q}_{qa}, \\ \epsilon_0^i \mathbb{1} & \text{for } \mathcal{Q}_i = \mathcal{Q}_{\phi d}, \mathcal{Q}_{d\chi}, \mathcal{Q}_{dXX}, \mathcal{Q}_{d\tilde{X}X}, \mathcal{Q}_{DdX^2}, \mathcal{Q}_{dX}, \mathcal{Q}_{HdX}, \mathcal{Q}_{da}, \end{cases}$$


8 operators are eliminated

- ▶ Numerics

$$\Delta_q = \begin{pmatrix} 0.8 & -3.3 - 1.5i & 79.3 + 35.4i \\ -3.3 + 1.5i & 16.6 & -397.5 + 8.1i \\ 79.3 - 35.4i & -397.5 - 8.1i & 9839.0 \end{pmatrix} \times 10^{-4}$$



$$\Delta_q \hat{\lambda}_d = \begin{pmatrix} 0.0021 & -0.18 - 0.08i & 191.3 + 85.4i \\ -0.009 + 0.004i & 0.88 & -958.7 + 19.6i \\ 0.21 - 0.10i & -21.1 - 0.4i & 23728.1 \end{pmatrix} \times 10^{-6}$$

Dark SMEFT with MFV: Scalar




all the operators survive
some ones highly constrained

Dark SMEFT with MFV: Fermion, ALP



all the operators survive

Dark SMEFT with MFV: Vector



all the operators survive, some ones highly constrained

Backup

$$\begin{aligned}
\mathcal{Q}_{d\phi} &= (\bar{q}_p d_r H) \phi + \text{h.c.}, & \mathcal{Q}_{d\phi^2} &= (\bar{q}_p d_r H) \phi^2 + \text{h.c.}, \\
\mathcal{Q}_{\phi q} &= (\bar{q}_p \gamma_\mu q_r) (i \phi_1 \overleftrightarrow{\partial}^\mu \phi_2), & \mathcal{Q}_{\phi d} &= (\bar{d}_p \gamma_\mu d_r) (i \phi_1 \overleftrightarrow{\partial}^\mu \phi_2),
\end{aligned} \tag{4.2}$$

$$\mathcal{Q}_{q\chi} = (\bar{q}_p \gamma_\mu q_r) (\bar{\chi} \gamma^\mu \chi), \quad \mathcal{Q}_{d\chi} = (\bar{d}_p \gamma_\mu d_r) (\bar{\chi} \gamma^\mu \chi), \tag{4.3}$$

$$\mathcal{Q}_{dHX} = (\bar{q}_p \sigma_{\mu\nu} d_r) H X^{\mu\nu} + \text{h.c.}, \tag{4.4}$$

$$\begin{aligned}
\mathcal{Q}_{dX} &= (\bar{d}_p \gamma_\mu d_r) X^\mu, & \mathcal{Q}_{HdX} &= (H^\dagger H) (\bar{d}_p \gamma^\mu d_r) X_\mu, \\
\mathcal{Q}_{qX} &= (\bar{q}_p \gamma_\mu q_r) X^\mu, & \mathcal{Q}_{HqX}^{(1)} &= (H^\dagger H) (\bar{q}_p \gamma^\mu q_r) X_\mu, \\
\mathcal{Q}_{dX^2} &= (\bar{q}_p d_r H) X_\mu X^\mu + \text{h.c.}, & \mathcal{Q}_{HqX}^{(3)} &= (H^\dagger \tau^I H) (\bar{q}_p \tau^I \gamma^\mu q_r) X_\mu, \\
\mathcal{Q}_{qXX} &= (\bar{q}_p \gamma_\mu q_r) X^{\mu\nu} X_\nu, & \mathcal{Q}_{dXX} &= (\bar{d}_p \gamma_\mu d_r) X^{\mu\nu} X_\nu, \\
\mathcal{Q}_{q\tilde{X}X} &= (\bar{q}_p \gamma_\mu q_r) \tilde{X}^{\mu\nu} X_\nu, & \mathcal{Q}_{d\tilde{X}X} &= (\bar{d}_p \gamma_\mu d_r) \tilde{X}^{\mu\nu} X_\nu, \\
\mathcal{Q}_{DqX^2} &= i (\bar{q}_p \gamma^\mu D^\nu q_r) X_\mu X_\nu + \text{h.c.}, & \mathcal{Q}_{DdX^2} &= i (\bar{d}_p \gamma^\mu D^\nu d_r) X_\mu X_\nu + \text{h.c.}, \\
\mathcal{Q}_{dHX^2} &= (\bar{q}_p \sigma_{\mu\nu} d_r H) X_1^\mu X_2^\nu + \text{h.c.}, & &
\end{aligned} \tag{4.5}$$

$$\mathcal{C}_i = \tilde{\mathcal{C}}_i \cdot \begin{cases} (m_X/\Lambda)^2 & \text{for } \mathcal{Q}_i = \mathcal{Q}_{dX^2}, \mathcal{Q}_{DdX^2}, \mathcal{Q}_{DqX^2}, \mathcal{Q}_{dHX^2}, \\ (m_X/\Lambda) & \text{for } \mathcal{Q}_i = \text{others.} \end{cases}$$

$$\mathcal{Q}_{qa} = (\bar{q}_p \gamma_\mu q_r) \partial^\mu a, \quad \mathcal{Q}_{da} = (\bar{d}_p \gamma_\mu d_r) \partial^\mu a, \tag{4.7}$$

Backup

One can also apply the MFV hypothesis to the lepton sector. However, since the mechanism of neutrino mass generation is still unknown, there are different approaches to formulate the leptonic MFV [73–79]. Here, we consider the realization of leptonic MFV within the so-called minimal field content [73, 74], in which the neutrino masses are generated by the Weinberg operator. In this case, the Yukawa interactions in the lepton sector can be written as

$$-\Delta\mathcal{L} = \bar{e} Y_e H^\dagger l + \frac{1}{2\Lambda_{\text{LN}}} (\bar{l}^c \tau_2 H) Y_\nu (H^T \tau_2 l) + \text{h.c.}, \quad (2.18)$$

where l denotes the left-handed lepton doublet with the charge conjugated field given by $l^c = -i\gamma_2 l^*$, and e is the right-handed charged lepton singlet. Λ_{LN} denotes the breaking scale of the lepton number symmetry $U(1)_{\text{LN}}$. Y_e and Y_ν stand for the 3×3 Yukawa coupling matrices in flavour space. In the absence of these Yukawa couplings, the lepton sector respects the flavour symmetry

$$G_{\text{LF}} = SU(3)_l \otimes SU(3)_e. \quad (2.19)$$

finite polynomial of $\mathbf{A}_\ell$ and $\mathbf{B}_\ell$. After neglecting all the terms involving $\mathbf{B}_\ell$, which are suppressed by the small lepton Yukawa couplings Y_e , we obtain





$$\mathcal{C}_{\text{MFV}} \approx \kappa_0 + \kappa_1 \mathbf{A}_\ell + \kappa_2 \mathbf{A}_\ell^2, \quad (2.21)$$

where the coefficients $\kappa_{0,1,2}$ are free real parameters. In the numerical analysis, we keep only the leading lepton flavour violation term $\mathbf{A}_\ell$ for simplicity, i.e., $\kappa_2 = 0$. Turning to the lepton mass eigenbasis, the current $\bar{l}\gamma^\mu \mathcal{C} l$ gives in the MFV hypothesis the following interactions:

$$\bar{e}_L \gamma^\mu (\kappa_0 \mathbb{1} + \kappa_0 \Delta_\ell) e_L + \bar{\nu}_L \gamma^\mu (\kappa_0 \mathbb{1} + \kappa_0 \hat{\lambda}_\nu^2) \nu_L, \quad (2.22)$$

where the basic LFV coupling Δ_ℓ can be obtained from $\mathbf{A}_\ell$ and takes the form


$$\Delta_\ell = U \hat{\lambda}_\nu^2 U^\dagger, \quad (2.23)$$





$$\Delta_\ell^{\text{NO}} = \begin{pmatrix} -0.19 - 0.01i & -0.25 - 0.02i & 0.31 - 0.04i \\ 0.12 + 0.01i & 0.28 - 0.00i & 0.29 + 0.04i \\ -0.37 - 0.01i & 0.21 - 0.05i & -0.03 + 0.01i \end{pmatrix}, \quad \Delta_\ell^{\text{IO}} = \begin{pmatrix} 0.21 + 0.09i & -0.34 + 0.05i & 0.03 + 0.11i \\ 0.31 + 0.12i & 0.19 + 0.00i & -0.15 - 0.14i \\ 0.12 - 0.02i & 0.04 - 0.19i & 0.34 - 0.10i \end{pmatrix}$$

RECONSTRUCTION AND SELECTION

- Charged particles: $p_T > 100$ MeV/c, close to collision point, in the central part of the detector
- Neutral particles: $E > 100$ MeV, in the central part of the detector
- Signal kaon candidates reconstructed applying kaon-enriching selection



In following, for



Particle	BESIII	STCF (1 ab ⁻¹)	Belle II (50 ab ⁻¹ on $\Upsilon(4S)$)	LHCb (300 fb ⁻¹)	CEPC (TDR)
$B^0, \bar{B}^0$	-	-	5.4×10^{10}	3×10^{13}	4.8×10^{11}
$B^\pm$	-	-	5.7×10^{10}	3×10^{13}	4.8×10^{11}
$B_s^0, \bar{B}_s^0$	-	-	6.0×10^8 (5 ab ⁻¹ on $\Upsilon(5S)$)	1×10^{13}	1.2×10^{11}
$B_c^\pm$	-	-	-	1×10^{11}	7.2×10^8
$\Lambda_b^0, \bar{\Lambda}_b^0$	-	-	-	2×10^{13}	1×10^{11}