Measurement of Global Spin Polarization at RHIC

Subhash Singha

(subhash@impcas.ac.cn)

中国科学院近代物理研究所

Institute of Modern Physics Chinese Academy of Sciences, Huizhou

The 2nd International Workshop on Physics at High Baryon Density PHD 2025

CCNU, Wuhan

October 16 – 18, 2025






Outline

- RHIC Beam Energy Scan Program
- Experimental results:
 - Global spin polarization (Beam-energy, centrality & rapidity dependence)
 - Local spin polarization (Beam-energy, baryonic SHE)
 - Opportunities at high baryon density region
- Summary

RHIC BES program


Phase diagram of QCD



RHIC Beam Energy Scan (BES):

- Find signatures of Phase Transition
- QCD Critical point
- Turn-off of QGP signatures

BES Program:




Collider: 7.7, 9.2, 11.5, 14.6, 17.3, 19.6, 27, 39, 54.4, 62.4, 200 (GeV)

Fixed Target: 3.0, 3.2, 3.5, 3.9, 4.5, 5.2, 6.2, 7.2, 7.7, 9.2, 11.5, 13.7 (GeV)

High statistics data and detector upgrade in BES-II

→most precise data to map the QCD phase diagram

STAR Detector




- Uniform acceptance, full azimuthal coverage, excellent PID capability
- TPC: tracking, centrality and event plane
- EPD, ZDC, BBC: event plane (polarization axis, perpendicular to event plane)
- TPC+TOF: particle identification

Motivation


In non-central heavy-ion collisions

A large orbital angular momentum (OAM) imparted into the system

$$L = r \times p \sim bA\sqrt{s_{\rm NN}} \sim 10^4 \, \hbar$$



Angular momentum



Participants

Vorticity generation

Motivation

In non-central heavy-ion collisions

 A large orbital angular momentum (OAM) imparted into the system

$$L = r \times p \sim bA\sqrt{s_{\rm NN}} \sim 10^4 \, \hbar$$


 Gradient of longitudinal flow velocity → vorticity along orbital angular momentum

$$\omega = \frac{1}{2}\nabla \times v \qquad \omega_y = \frac{1}{2}(\nabla \times v)_y \approx \frac{1}{2}\frac{dv_z}{dy}$$


 Polarize quarks and anti-quarks due to "spin-orbit" interaction.

> Liang et. al., Phys Rev Lett B 94, 102301 (2005) Jiang et al, Phys Rev C 94, 044910 (2016)

QCD matter under rapid global rotation







QGP: femtometer ~ 10⁻¹⁵ m

Water: a few cm

Neutron-star: a few km

Major question:

How angular momentum distributed into the system?

Allows to study properties of the system

Vorticity	(s-1)
Tornado	10-1
Superfluid He	150
Heavy-ion collisions	10 ⁷ - 10 ²¹

Heavy-ion collisions create rapidly rotating QCD matter in laboratory \rightarrow controlled experiment

Measurement global spin polarization

Global polarization is measured from the angular distributions using parity violating weak decay of hyperons ("self-analyzing"):


$$\cdot \frac{d\mathbf{N}}{d\Omega^*} = \frac{1}{4\pi} (1 + \alpha_H \mathbf{P_H^*} \cdot \mathbf{p_d^*})$$

P_H: Hyperon polarization

 α_H : Hyperon decay parameter

 p_d : Daughter momentum direction

* : Measurements in parent's rest frame



Component perpendicular to reaction plane:

$$P_{H} = \frac{8}{\pi \alpha_{H}} \frac{\langle \sin(\Psi_{1} - \phi_{d}^{*}) \rangle}{\text{Res}(\Psi_{1})}$$

Schiling et. al., Nucl Phys B 15, 397 (1970) (STAR Collaboration) Phys Rev C 76, 024915 (2007)

Observation of global polarization of Λ hyperons

First prediction in 2005

Globally Polarized Quark-Gluon Plasma in Noncentral A+A Collisions

Zuo-Tang Liang¹ and Xin-Nian Wang^{2,1}


Show more ~

Phys. Rev. Lett. **94**, 102301 – **Published 14 March, 2005** Erratum Phys. Rev. Lett. **96**, 039901 (2006)

DOI: https://doi.org/10.1103/PhysRevLett.94.102301



Discovery in 2017






- Discover the Evidence of vorticity (~10²¹ s⁻¹) in Quark Gluon Plasma
- Opened new avenue of research: probe spinning QGP matter under extreme rotation

Global polarization: beam energy dependence

Beam energy dependence of global P_{Λ}



- BES-II: uncertainty reduced by a factor ~ 9
- P_∧ smooth increase with decrease in beam energy
- Models can capture the energy dependence trend

STAR:

Phys Rev C 76, 024915 (2007)

Nature 548, 62 (2017)


Phys Rev C 101, 044611 (2020)

Phys Rev C 104, 061901 (2021)


ALICE:


Phys Rev C 101, 044611 (2020)

Beam energy dependence of global P_{Λ}



- \bullet BES-II: uncertainty reduced by a factor ~ 9
- P_∧ smooth increase with decrease in beam energy
- Models can capture the energy dependence trend






Karpenko et. al., Eur Phys J C 77, 213 (2017) Ivanov et. al., Phys Rev C 100, 014908 (2019) Ivanov et. al., Phys Rev C 102, 024916 (2020)

High precision in data can differentiate models


Beam energy dependence of global $P_{\Lambda,\Xi,\Omega}$




- ullet BES-II: First measurements of Ξ and Ω , follow energy dependent trend
- ullet \bullet $P_\Lambda \sim P_\Xi \sim P_\Omega$: precision not sufficient to distinguish ordering

Global polarization: differential measurements (centrality and rapidity dependence)

P_{Λ} : centrality dependence






- ullet P_{Λ} increases from central to peripheral collisions
- Similar pattern followed across beam energies
- Trend consistent with expectation from vorticity

Is there a saturation? No drop in polarization expected from angular momentum?


P_{\Lambda}: rapidity dependence




 No significant rapidity dependence observed within measured acceptance


> Deng and Huang, Phys Rev C 93, 064907 (2016) Wu et al, Phys Rev Research 1, 033058 (2019) Alzhrani et. al., Phys Rev C 106, 014905 (2022)

Rapidity dependence of P_∧ is different among various models






Major question: How angular momentum is distributed into the system, correlation with baryon stopping?



Differences in global spin polarization P_{Λ} and $P_{\overline{\Lambda}}$


Possible constraint on B field by PA



In non-central heavy-ion collisions

- Initial strong magnetic field (B) is expected $eB \sim m_\pi^2 \sim 10^{18} \ {\rm Gauss}$
- Such strong B field can also polarize quarks. Can induce different spin polarization for quarks and antiquarks with different magnetic moments

Magnetic Field





 $|\overline{P_{\Lambda}}||\overline{B}|$ $|\overline{P_{\Lambda}}||-\overline{B}|$ |VS|


$$(P_{\Lambda} - P_{\bar{\Lambda}}) \approx \frac{2\mu_{\Lambda}B}{T}$$


Possible constraint on B field by PA

STAR BES-II









Scenarios can cause Λ and anti- Λ polarization difference:

- Different freeze out for particles and anti-particles
- Different response to mesonic field generated by baryonic current
-

McLerran and Skokov, Nucl Phys A 929, 184 (2014)


- No difference between $\Lambda, \bar{\Lambda}$ is observed with BES-II data
- ullet Results set an upper-limit on late stage magnetic-field ($B \leq 10^{13}~T$)

Local polarization: beam energy dependence


Local vortices via helicity polarization of A



More refine structures of vortices



Violent vortices in Quark-Gluon-Plasma in femtometer scale
$$\omega = \frac{1}{2} \nabla \times v$$
 Polarization driven by hydrodynamic gradients



Local vortices within QGP -> azimuthal angle dependence of Λ polarization (most challenging for theory)

Contributions = Thermal vorticity + Kinematic vorticity + Shear effect + Baryon density gradient (bSHE)

Baryonic Spin Hall effect (SHE)

Condensed matter

Heavy Ion Collisions


 $s \propto \pm p \times E$


 $s \propto \pm p \times \nabla \mu_B$

splitting in spin on opposite boundaries

splitting in local polarization of Λ and $\overline{\Lambda}:P_z^{\Lambda}-P_z^{\overline{\Lambda}}$

"spin-orbit"






Predicted Spin Hall type effect driven by gradient of baryonic density ($\nabla \mu_B$)


Polarization \sim vorticity \oplus ∇T \oplus Shear \oplus $\nabla \mu_B$

Fu et., al., Phys Rev Lett 127, 142301 (2021) Liu and Yin, Phys Rev D 104, 054043 (2021)







Longitudinal-plane (x, η)



Baryonic Spin Hall effect (bSHE)





$$P_{2,z} = \langle P_z \sin(2\Delta\phi) \rangle$$

$$P_{y} = \frac{8}{\pi \alpha_{\Lambda} R_{EP}} \langle \sin(\Psi_{1} - \phi_{p}^{*}) \rangle$$

$$P_z = \frac{3}{\alpha_H} \frac{\langle \cos \theta_{\rm p}^* \rangle}{\langle (\cos \theta_{\rm p}^*)^2 \rangle}$$

Kinematics:

 $0.5 < p_T < 5.0 \text{ GeV/c}$ |y|< 1.0 for 27 GeV


|y| < 1.5 for other energies


Qiang Hu, SQM 2024 (STAR)

Polarization \sim vorticity \oplus ∇T \oplus Shear \oplus $\nabla \mu_B$

ullet $P_{2,z}$: non-zero, no strong energy dependence, (hints of sign change at lower energies ?)

Baryonic Spin Hall effect (bSHE)





 $P_{2,z}^{\text{net}} = P_{2,z}^{\Lambda} - P_{2,z}^{\overline{\Lambda}}$

$$P_{y} = \frac{8}{\pi \alpha_{\Lambda} R_{EP}} \langle \sin(\Psi_{1} - \phi_{p}^{*}) \rangle$$

$$P_{2,y} = -\langle P_y \cos(2\Delta\phi) \rangle$$

$$P_z = \frac{3}{\alpha_H} \frac{\langle \cos \theta_p^* \rangle}{\langle (\cos \theta_p^*)^2 \rangle}$$

$$P_{2,z} = \langle P_z \sin(2\Delta\phi) \rangle$$

Kinematics:

 $0.5 < p_T < 5.0 \text{ GeV/c}$


|y|< 1.0 for 27 GeV

|y|<1.5 for other energies

Qiang Hu, SQM 2024 (STAR)

Not sufficient precision at BES energies

Future opportunities Global polarization at high baryon density region




Global spin polarization of A at high baryon density



Faster/Slower rotation induces anisotropic stresses and redistributes matter internally

→ Equation of State (EoS)

- Beam energy depedence of angular momentum interactions
 - Can induce *non-monotonicity* and affects *QCD-EoS*





$$L \sim \frac{1}{2} Ab\sqrt{s} \sqrt{1 - (2M/\sqrt{s})^2}$$

$$P_{\Lambda} \sim 0$$
 at $\sqrt{s_{\rm NN}} \sim 2 m_{\rm N}$


• Is there a non-monotonic pattern in global polarization?

Global spin polarization of A at high baryon density






- Is there a non-monotonic pattern in global polarization?
- Interplay between angular momentum and baryon stopping?
- Partonic versus Hadronic dominant medium carrying vorticity?
- Dependence on QCD EoS with spin degrees of freedom




- \bullet Measurement of P_{Λ} using STAR fixed-target data is underway
- Unlike higher beam energy, 3 GeV acceptance can reach mid and target rapidity region


Global spin polarization of hyper-triton ³_ΛH (pnΛ)



- Unique loosely bound "pn/\lambda" system under global rotation
- Abundant ³_∧H Production at FXT region



• Measurement using STAR fixed-target data is underway




Sign of ³_ΛH polarization relative to Λ
 What is spin structure of hyper-triton? singlet or triplet?


Next Talk: Kaijia Sun

Summary

- Global spin polarization of hyperons (P_y)
 - monotonic beam energy dependence for Λ ,
 - $P_{\Lambda} \sim P_{\Xi} \sim P_{\Omega}$ follow same energy dependence
 - clear centrality dependence at all beam energies
 - no strong rapidity dependence within acceptance
 - No difference between Λ and $\overline{\Lambda}$
- Local spin polarization of hyperons
 - In-plane $(P_{2,z})$ positive magnitude
 - Baryonic Spin Hall effect: $(P_{2,z,z}^{\Lambda-\overline{\Lambda}})$ hints of sign change, no precision




- Measurement at high baryon density are essential to address important questions:
- Expected non-monotonic pattern induced by angular momentum?
- Correlation with baryon stopping?
- QCD EoS?




Thank you for your attention!

Many Thanks to STAR colleagues!

Local spin polarization of hyperons





ullet $P_{2,z}$ shows significant centrality dependence, expected from elliptic flow


STAR: Phys Rev Lett 131, 202301 (2023)

Spin-spin correlations in BES


Article Published: 18 January 2023

Pattern of global spin alignment of φ and K^{*0} mesons in heavy-ion collisions


ature 614, 244–248 (2023) Cite this article



- Signal larger than expectation from P_{Λ}
- Role of spin fluctuation









Error projection for BES-II

P_{Λ} : rapidity dependence with models






Ivanov et. al., Phys Rev C 105, 034915 (2022)

- Challenge for present models to explain rapidity dependence of P_{Λ} , while they can largely capture its energy dependence at mid-rapidity
- Needed proper modeling of amount of angular momentum deposition along rapidity

The correlation between net-baryon and angular momentum needed to be understood


Local (P₇) spin polarization of hyperons




- ullet Clear centrality dependence of P_{z}
- At mid-central collisions
 Zr+Zr,Ru+Ru > Au+Au ~ Pb+Pb
- Hints of system size dependence, No obvious energy dependence

STAR: Phys Rev Lett 123, 132301 (2019) STAR: Phys Rev Lett 131, 202301 (2023) ALICE: Phys Rev Lett 128, 172005 (2022)

P_{Λ} : centrality dependence with 3FD model





Baryonic Spin Hall effect (SHE)

Condensed matter

Heavy Ion Collisions


 $s \propto \pm p \times E$


$$s \propto \pm p \times \nabla \mu_B$$

splitting in spin on opposite boundaries

splitting in local polarization of Λ and $\overline{\Lambda}:P_z^{\Lambda}-P_z^{\overline{\Lambda}}$

"spin-orbit"





Predicted Spin Hall type effect driven by gradient of baryonic density ($\nabla \mu_B$)

Polarization \sim vorticity \oplus ∇T \oplus Shear \oplus $\nabla \mu_B$

Fu et., al., Phys Rev Lett 127, 142301 (2021) Liu and Yin, Phys Rev D 104, 054043 (2021)

How do we visualize?










Figure 1: An example of ink-made fish animation created with our method. Two fish are swimming in water freely, and the fishtails swing in high speed. Ink flows emit from the fishtails at the same time, then the attractive dynamic ink diffusion effects can be observed.


https://doi.org/10.1145/2087756.20877

Naive picture

- Consider *Ink* as baryons in the medium
- Ink (baryon) density is changing \rightarrow (baryon) chemical potential gradient ($\nabla \mu_R$)
- Heavy-ion collisions (much complicated) → adding ink (baryon) into expanding swirling water → baryonic Spin Hall effect
- How the ink (baryon) density changing in transverse and longitudinal plane?


Local spin polarization






Alzhrani et. al., Phys Rev C 106, 014905 (2022)

Beam energy dependence of global PA





- Similar rising pattern in polarization and directed flow at low energy
- Some models can capture such rising trend in both P_H and v₁