

# Forward Hadron Productions in proton-nucleus collisions at NLO

Bo-Wen Xiao

School of Science and Engineering, CUHK-Shenzhen

[G. A. Chirilli, BX, Feng Yuan, 12]



[A. Stasto, BX, D. Zaslavsky, 14]

[Y. Shi, L. Wang, S.Y. Wei, BX, 22]



# Saturation Physics, Color Glass Condensate

Describe the **emergent property** of high density gluons inside proton and nuclei.




- Gluon density grows rapidly as  $x$  gets small.
- Many gluons with fixed size packed in a confined hadron, gluons **overlap and recombine**  $\Rightarrow$  **Non-linear QCD dynamics** (BK-JIMWLK)  $\Rightarrow$  **ultra-dense gluons** with collective property.



# Forward hadron production in $pA$ collisions

[Dumitru, Jalilian-Marian, 02] Dilute-dense factorization at forward rapidity

$$\frac{d\sigma_{\text{LO}}^{pA \rightarrow hX}}{d^2p_{\perp} dy_h} = \int_{\tau}^1 \frac{dz}{z^2} \left[ x_1 q_f(x_1, \mu) \mathcal{F}_{x_2}(k_{\perp}) D_{h/q}(z, \mu) + x_1 g(x_1, \mu) \tilde{\mathcal{F}}_{x_2}(k_{\perp}) D_{h/g}(z, \mu) \right].$$




- $\mathcal{F}(k_{\perp})$  (dipole gluon distribution) encodes dense gluon info.
- **Early attempts:** [Dumitru, Hayashigaki, Jalilian-Marian, 06; Altinoluk, Kovner 11]  
 [Altinoluk, Armesto, Beuf, Kovner, Lublinsky, 14]
- Full NLO: [Chirilli, BX and Yuan, 12]



## Wilson Lines in Color Glass Condensate Formalism

The Wilson loop (**color singlet dipole**) in McLerran-Venugopalan (MV) model

$$\frac{1}{N_c} \langle \text{Tr} U(x_\perp) U^\dagger(y_\perp) \rangle = e^{-\frac{g_s^2 (x_\perp - y_\perp)^2}{4}}$$




- Dipole amplitude  $S^{(2)}$  then produces the quark  $k_T$  spectrum via Fourier transform

$$\mathcal{F}(k_\perp) \equiv \frac{dN}{d^2k_\perp} = \int \frac{d^2x_\perp d^2y_\perp}{(2\pi)^2} e^{-ik_\perp \cdot (x_\perp - y_\perp)} \frac{1}{N_c} \langle \text{Tr} U(x_\perp) U^\dagger(y_\perp) \rangle.$$



## d+Au collisions at RHIC

$$R_{d+Au} = \frac{1}{\langle N_{\text{coll}} \rangle} \frac{d^2 N_{d+Au} / d^2 p_T d\eta}{d^2 N_{pp} / d^2 p_T d\eta}; \quad R = 1 \quad \text{benchmark for no nuclear effects.}$$



BRAHMS


- Cronin effect at middle rapidity: redistribution of momentum
- Rapidity evolution of the nuclear modification factors  $R_{d+Au}$
- Promising evidence for gluon saturation effects



## New LHCb Results

[R. Aaet al. (LHCb Collaboration), Phys. Rev. Lett. 128 (2022) 142004]

$$R_{pPb} = \frac{1}{\langle N_{\text{coll}} \rangle} \frac{d^2 N_{p+Pb} / d^2 p_T d\eta}{d^2 N_{pp} / d^2 p_T d\eta}$$




- Forward rapidity:  
Nuclear effects  
Small- $x$  suppression
- Backward rapidity:  
Cronin enhancement

- Rapidity evolution of the nuclear modification factors  $R_{pPb}$  similar to RHIC



## NLO diagrams in the $q \rightarrow q$ channel

[Chirilli, BX and Yuan, 12]




- Take into account real (top) and virtual (bottom) diagrams together!
- Non-linear multiple interactions inside the grey blobs!
- Integrate over gluon phase space  $\Rightarrow$  Divergences!.



# Factorization for single inclusive hadron productions

Factorization for the  $p + A \rightarrow H + X$  process [Chirilli, BX and Yuan, 12]



- Include all real and virtual graphs in all channels  $q \rightarrow q$ ,  $q \rightarrow g$ ,  $g \rightarrow q(\bar{q})$  and  $g \rightarrow g$ .
- 1. collinear to target nucleus; rapidity divergence  $\Rightarrow$  BK evolution for UGD  $\mathcal{F}(k_\perp)$ .
- 2. collinear to the initial quark;  $\Rightarrow$  DGLAP evolution for PDFs
- 3. collinear to the final quark.  $\Rightarrow$  DGLAP evolution for FFs.





## Factorization and NLO Calculation

- Factorization is about separation of **short distant physics** (perturbatively calculable **hard factor**) from **large distant physics** (Non perturbative).

$$\sigma \sim xf(x) \otimes \mathcal{H} \otimes D_h(z) \otimes \mathcal{F}(k_\perp)$$

- NLO (1-loop) calculation always contains various kinds of **divergences**.
  - Some divergences can be absorbed into the corresponding **evolution equations**.
  - Renormalization: cutting off infinities and hiding the ignorance.
  - The rest of divergences should be canceled.

- **Hard factor**

$$\mathcal{H} = \mathcal{H}_{\text{LO}}^{(0)} + \frac{\alpha_s}{2\pi} \mathcal{H}_{\text{NLO}}^{(1)} + \dots$$

should always be finite and free of divergence of any kind.



## Hard Factor of the $q \rightarrow q$ channel

$$\frac{d^3 \sigma^{p+A \rightarrow h+X}}{dy d^2 p_{\perp}} = \int \frac{dz}{z^2} \frac{dx}{x} \xi x q(x, \mu) D_{h/q}(z, \mu) \int \frac{d^2 x_{\perp} d^2 y_{\perp}}{(2\pi)^2} \left\{ S_Y^{(2)}(x_{\perp}, y_{\perp}) \left[ \mathcal{H}_{2qq}^{(0)} + \frac{\alpha_s}{2\pi} \mathcal{H}_{2qq}^{(1)} \right] \right. \\ \left. + \int \frac{d^2 b_{\perp}}{(2\pi)^2} S_Y^{(4)}(x_{\perp}, b_{\perp}, y_{\perp}) \frac{\alpha_s}{2\pi} \mathcal{H}_{4qq}^{(1)} \right\}$$

$$\mathcal{H}_{2qq}^{(1)} = C_F \mathcal{P}_{qq}(\xi) \ln \frac{c_0^2}{r_{\perp}^2 \mu^2} \left( e^{-ik_{\perp} \cdot r_{\perp}} + \frac{1}{\xi^2} e^{-i \frac{k_{\perp}}{\xi} \cdot r_{\perp}} \right) - 3C_F \delta(1-\xi) \ln \frac{c_0^2}{r_{\perp}^2 k_{\perp}^2} e^{-ik_{\perp} \cdot r_{\perp}} \\ - (2C_F - N_c) e^{-ik_{\perp} \cdot r_{\perp}} \left[ \frac{1 + \xi^2}{(1-\xi)_+} \tilde{I}_{21} - \left( \frac{(1 + \xi^2) \ln(1-\xi)^2}{1-\xi} \right)_+ \right] \\ \mathcal{H}_{4qq}^{(1)} = -4\pi N_c e^{-ik_{\perp} \cdot r_{\perp}} \left\{ e^{-i \frac{1-\xi}{\xi} k_{\perp} \cdot (x_{\perp} - b_{\perp})} \frac{1 + \xi^2}{(1-\xi)_+} \frac{1}{\xi} \frac{x_{\perp} - b_{\perp}}{(x_{\perp} - b_{\perp})^2} \cdot \frac{y_{\perp} - b_{\perp}}{(y_{\perp} - b_{\perp})^2} \right. \\ \left. - \delta(1-\xi) \int_0^1 d\xi' \frac{1 + \xi'^2}{(1-\xi')_+} \left[ \frac{e^{-i(1-\xi') k_{\perp} \cdot (y_{\perp} - b_{\perp})}}{(b_{\perp} - y_{\perp})^2} - \delta^{(2)}(b_{\perp} - y_{\perp}) \int d^2 r'_{\perp} \frac{e^{ik_{\perp} \cdot r'_{\perp}}}{r'_{\perp}{}^2} \right] \right\},$$

where

$$\tilde{I}_{21} = \int \frac{d^2 b_{\perp}}{\pi} \left\{ e^{-i(1-\xi) k_{\perp} \cdot b_{\perp}} \left[ \frac{b_{\perp} \cdot (\xi b_{\perp} - r_{\perp})}{b_{\perp}^2 (\xi b_{\perp} - r_{\perp})^2} - \frac{1}{b_{\perp}^2} \right] + e^{-ik_{\perp} \cdot b_{\perp}} \frac{1}{b_{\perp}^2} \right\}.$$



## Numerical implementation of the NLO result

Single inclusive hadron production up to NLO

$$d\sigma = \int xf_a(x) \otimes D_a(z) \otimes \mathcal{F}_a^{xg}(k_\perp) \otimes \mathcal{H}^{(0)} + \frac{\alpha_s}{2\pi} \int xf_a(x) \otimes D_b(z) \otimes \mathcal{F}_{(N)ab}^{xg} \otimes \mathcal{H}_{ab}^{(1)}.$$




Consistent implementation should include all the NLO  $\alpha_s$  corrections.

- **NLO parton distributions.** (MSTW or CTEQ)
- **NLO fragmentation function.** (DSS or others.)
- **Use NLO hard factors.** [Chirilli, BX and Yuan, 12]
- **Use the one-loop approximation for the running coupling**
- **rcBK evolution equation for the dipole gluon distribution** [Balitsky, Chirilli, 08; Kovchegov, Weigert, 07]. Full NLO BK evolution not available.
- **Saturation physics at One Loop Order (SOLO).** [Stasto, Xiao, Zaslavsky, 13]



# Numerical implementation of the NLO result


Saturation physics at One Loop Order (**SOLO**). [Stasto, Xiao, Zaslavsky, 13]



- Reduced factorization scale dependence!
- **Catastrophe:** Negative NLO cross-sections at high  $p_T$ .
- Fixed order calculation in field theories is not **guaranteed to be positive**.



## The cross-section at high $k_{\perp}$




- In the dilute limit  $k_{\perp} \gg Q_s$ , partonic cross-sections follow the power law

$$\sigma(k_{\perp}) \sim \mathcal{F}(k_{\perp}) \sim \frac{Q_s^2}{4} \int d^2 r_{\perp} e^{-ik_{\perp} \cdot r_{\perp}} r_{\perp}^2 \ln(r_{\perp} \Lambda) \sim \frac{Q_s^2}{k_{\perp}^4}.$$

- **NLO**  $\sigma_{NLO} \sim \mathcal{CF}(k_{\perp}) \sim \mathcal{C} \frac{Q_s^2}{k_{\perp}^4}$  with  $\mathcal{C}$  containing logarithms such as  $\ln k_{\perp}^2 / Q_s^2$ .



# Large Logarithms



- NLO vs NLL Naive  $\alpha_s$  expansion sometimes is not sufficient!

	LO	NLO	NNLO	...
LL	1	$\alpha_s L$	$(\alpha_s L)^2$	...
NLL		$\alpha_s$	$\alpha_s (\alpha_s L)$	...
...			...	...

- Evolution  $\rightarrow$  Resummation of large logs.  
 LO evolution resums LL; NLO  $\Rightarrow$  NLL.



## Extending the applicability of CGC calculation


- Goal: find a solution within our **current factorization** (exactly resum  $\alpha_s \ln 1/x_g$ ) to extend the applicability of CGC. **Other scheme choices** certainly is possible.
- A lot of logs **arise** in pQCD loop-calculations: **DGLAP, small- $x$ , threshold, Sudakov**.
- **Breakdown** of  $\alpha_s$  expansion occurs due to the appearance of logs in certain PS.
- Demonstrate **onset of saturation** and visualize **smooth transition to dilute regime**.
- Add'l consideration: numerically challenging due to **limited computing resources**.
- Towards a more complete framework. [Altinoluk, Armesto, Beuf, Kovner, Lublinsky, 14; Kang, Vitev, Xing, 14; Ducloue, Lappi and Zhu, 16, 17; Iancu, Mueller, Triantafyllopoulos, 16; Liu, Ma, Chao, 19; Kang, Liu, 19; Kang, Liu, Liu, 20; Altinoluk, Armesto, Kovner, Lublinsky, 23]. Implication for other NLO calculations in CGC. [Tael, Altinoluk, Beuf, Marquet, 22; Iancu, Mulian, 22; Caucal, Salazar, Schenke, Stebel, Venugopalan, 23; Bergabo, Jalilian-Marian, 22, 23; Altinoluk, Armesto, Beuf, 23; Beuf, Lappi, Mäntysaari, Paatelainen, Penttala, 24; etc]



## NLO hadron productions in $pA$ collisions with kinematic constraints

[Watanabe, Xiao, Yuan, Zaslavsky, 15] **Rapidity subtraction!** with kinematic constraints

- Originally assume the limit  $s \rightarrow \infty$



$$\int_0^{1-\frac{q_\perp^2}{x_p s}} \frac{d\xi}{1-\xi} = \underbrace{\ln \frac{1}{x_g}}_{1-\xi < \frac{q_\perp^2}{k_\perp^2}} + \underbrace{\ln \frac{k_\perp^2}{q_\perp^2}}_{\text{missed earlier}} \Rightarrow$$

New terms:  $L_q + L_g$  from  $q_\perp^2 \leq (1-\xi)k_\perp^2$ .


- Corrections related to threshold double logs. **Negative** when  $p_T \gg Q_s$  at forward  $y$  ( $x_p \rightarrow 1$ )! Approach **threshold** at high  $k_\perp$ .
- Ioffe time cutoff [Altinoluk, Armesto, Beuf, Kovner and Lublinsky, 14]





# Numerical results with kinematic constraint

SOLO results [Stasto, Xiao, Zaslavsky, 13; Watanabe, Xiao, Yuan, Zaslavsky, 15]




- SOLO still breaks down in the large  $p_{\perp}$  region with the new term.



## Gluon Radiation at the Threshold

Near threshold:

 radiated gluon has to be soft!  $\tau = \frac{p_{\perp} e^y}{\sqrt{s}}$  density ( $\tau = x_p \xi z \leq 1$ )


- If  $q_{\perp} \sim k_{\perp}$ , then  $q^{-} \rightarrow \infty$ , this is part of the small- $x$  evolution.
- If  $q_{\perp}^2 \leq (1 - \xi)k_{\perp}^2$ , then  $q^{-}$  is finite, this is part of the Sudakov!  $\Rightarrow \ln \frac{k_{\perp}^2}{q_{\perp}^2}$ .
- KLN  $\Rightarrow$  cancellation between real and virtual.  $-\int_{\Lambda^2}^{k_{\perp}^2} \frac{dq_{\perp}^2}{q_{\perp}^2} \ln \frac{k_{\perp}^2}{q_{\perp}^2} = -\frac{1}{2} \ln^2 \frac{k_{\perp}^2}{\Lambda^2}$
- Introduce an additional semi-hard scale  $\Lambda^2$  as the typical  $q_{\perp}$ .



## Threshold resummation in the CGC formalism

Threshold logarithms: **Collinear (plus-distribution)** and **Sudakov soft gluon** part

- Two equivalent methods to resum the collinear part ( $P_{ab}(\xi) \ln \Lambda^2/\mu^2$ ):
  1. Reverse DGLAP evolution; 2. RGE method (threshold limit  $\xi \rightarrow 1$ ).
- Introduce forward threshold quark jet function  $\Delta^q(\Lambda^2, \mu^2, \omega)$ , which satisfies

$$\frac{d\Delta^q(\omega)}{d \ln \mu^2} = -\frac{d\Delta^q(\omega)}{d \ln \Lambda^2} = -\frac{\alpha_s C_F}{\pi} \left[ \ln \omega + \frac{3}{4} \right] \Delta^q(\omega) + \frac{\alpha_s C_F}{\pi} \int_0^\omega d\omega' \frac{\Delta^q(\omega) - \Delta^q(\omega')}{\omega - \omega'}.$$

- Consistent with the threshold resummation in SCET[Becher, Neubert, 06]!



## Threshold resummation in the CGC formalism

Threshold logarithms: **Sudakov soft gluon** part and **Collinear (plus-distribution)** part.

- Soft single and double logs ( $\ln k_{\perp}^2/\Lambda^2, \ln^2 k_{\perp}^2/\Lambda^2$ ) are resummed via Sudakov factor.
- Performing Fourier transformations

$$\begin{aligned} \int \frac{d^2 r_{\perp}}{(2\pi)^2} \mathcal{S}(r_{\perp}) \ln \frac{\mu^2}{\mu_r^2} e^{-ik_{\perp} \cdot r_{\perp}} &= - \int \frac{d^2 l_{\perp}}{\pi l_{\perp}^2} \left[ F(k_{\perp} + l_{\perp}) - J_0\left(\frac{c_0}{\mu} l_{\perp}\right) F(k_{\perp}) \right] \\ &= -\frac{1}{\pi} \int \frac{d^2 l_{\perp}}{(l_{\perp} - k_{\perp})^2} \left[ F(l_{\perp}) - \frac{\Lambda^2}{\Lambda^2 + (l_{\perp} - k_{\perp})^2} F(k_{\perp}) \right] + F(k_{\perp}) \ln \frac{\mu^2}{\Lambda^2}. \end{aligned}$$

- Similar technique is used to extract the double log. At one loop  $\Lambda$  is **arbitrary**.
- After resumming all logs,  $\mu^2$  and  $\Lambda^2$  dependences only cancel up to one-loop order.
- Choose proper  $\mu^2$  (hard scale) and  $\Lambda^2$  (Saddle point) in resummation to **minimize hard factors**.
- Goal of resummation: 1. Estimate  $\Lambda$  according to Sudakov and Saturation effects;  
 2. make sure the un-resummed contribution is small, **restore perturbative expansion**.



## Threshold Logarithms

[Watanabe, Xiao, Yuan, Zaslavsky, 15; Shi, Wang, Wei, Xiao, 21] ▶ 2112.06975 [hep-ph]

- **Threshold enhancement for  $\sigma$ :**  $e^{-x} = 1 - x + \frac{x^2}{2} + \dots$
- In the coordinate space, we can identify two types of logarithms


$$\text{single log: } \ln \frac{k_{\perp}^2}{\mu_r^2} \rightarrow \ln \frac{k_{\perp}^2}{\Lambda^2}, \quad \ln \frac{\mu^2}{\mu_r^2} \rightarrow \ln \frac{\mu^2}{\Lambda^2}; \quad \text{double log: } \ln^2 \frac{k_{\perp}^2}{\mu_r^2} \rightarrow \ln^2 \frac{k_{\perp}^2}{\Lambda^2},$$

with  $\mu_r \equiv c_0/r_{\perp}$  with  $c_0 = 2e^{-\gamma_E}$ .

- Introduce a semi-hard **auxiliary scale**  $\Lambda^2 \sim \mu_r^2 \gg \Lambda_{QCD}^2$ . **Identify dominant  $r_{\perp}$ !**
- Dependence on  $\mu^2$ ,  $\Lambda^2$  cancel **order by order**. Choose “natural” values at fixed order.
- For running coupling,  $\Lambda^2 = \Lambda_{QCD}^2 \left[ \frac{(1-\xi)k_{\perp}^2}{\Lambda_{QCD}^2} \right]^{C_R/[C_R+\beta_1]}$ . **Akin to CSS & Catani *et al.***




## Numerical Results for $pA$ spectra



- Excellent agreement with data across many orders of magnitudes for different energies and  $p_T$  ranges measured from both RHIC and the LHC!
- Explain the rapidity dependence: threshold (Sudakov) logs are less important in the forward regime. In middle rapidity, large phase space gives large  $\alpha_s \ln^2 \frac{k_{\perp}^2}{\Lambda^2}$






## Comparison with the new LHCb data



- LHCb data: 2108.13115
- [Data Link](#) [DIS2021](#)
- Threshold effect is not important at low  $p_T$  for LHCb data. Saturation effects are still dominant.
- Predictions are improved from LO to NLO.
- Solve the negativity problem at both RHIC and LHC.



# Summary



- **Ten-Year Odyssey** in **NLO hadron productions** in  $pA$  collisions in CGC.
- Towards the **precision** test of saturation physics (CGC) at RHIC and LHC.
- Impact: Similar technique used in DIS processes, e.g. [Caucal, et al 22]
- Exciting time of NLO CGC phenomenology with **the upcoming EIC**.

