


Recent flow results from ATLAS

Qipeng Hu (胡启鹏), <u>qipeng.hu@cern.ch</u>

University of Science and Technology of China

October 26, 2025






The 16th Workshop on QCD Phase Transition and Relativistic Heavy-Ion Physics (QPT 2025)

Oct. 24-28, 2025, Guilin

Why light ions collisions at LHC


- Bridge the gap between pp and Pb+Pb/Xe+Xe to test QGP formation in light-ion collisions
- If QGP is formed, use most-central ²⁰Ne (deformed) vs ¹⁶O (spherical) collisions to probe nuclear structure





Giuliano Giacalone et al. PRL 135 (2025) 012302 Jiangyong Jia, arXiv:2501.16071

ATLAS heavy-ion physics program




System	Year	$\sqrt{ m s_{NN}}$ [TeV]	L _{int} [nb ⁻¹]
Pb+Pb	2010	2.76	0.007
Pb+Pb	2011	2.76	0.14
рр	2013	2.76	4000
p+Pb	2013	5.02	29
рр	2015	5.02	28000
Pb+Pb	2015	5.02	0.49
p+Pb	2016	5.02	0.5
p+Pb	2016	8.16	160
Xe+Xe	2017	5.44	0.003
рр	2017	5.02	270000
Pb+Pb	2018	5.02	1.76
Pb+Pb	2023	5.36	1.71
рр	2024	5.36	425000
Pb+Pb	2024	5.36	1.73
p+O	2025	9.62	12.48
0+0	2025	5.36	8.63
Ne+Ne	2025	5.36	1.05
Pb+Pb	2025	5.36	To be taken
??	2026	??	??


All ATLAS HI public results: https://twiki.cern.ch/twiki/bin/view/AtlasPublic/HeavylonsPublicResults

T S

Trigger strategy




- Minimum-bias event triggering with TRT
- Unbiased down to very low multiplicities
- Significantly better performance than other ATLAS minimum-bias triggers (MBTS, ZDC, etc.)






O+O and Ne+Ne Centrality




- Centrality from Forward Calorimeter energy sum (0–80%)
- Glauber analysis with improved light-ion geometry and fluctuations (Loizides, arXiv:2507.05853)
- · Event-by-event fluctuations play a larger role in light-ion collisions in most central collisions (long tails)



Centrality definitions: forward energy vs. multiplicity











Two particle correlation



Shifted peripheral reference (>80%) for visibility




Flow modulation around pedestal

$$v_n(p_{\mathrm{T}}^b) = \frac{v_{n,n}(p_{\mathrm{T}}^a, p_{\mathrm{T}}^b)}{v_n(p_{\mathrm{T}}^a)} = \frac{v_{n,n}(p_{\mathrm{T}}^a, p_{\mathrm{T}}^b)}{\sqrt{v_{n,n}(p_{\mathrm{T}}^a, p_{\mathrm{T}}^a)}}$$

$$\begin{split} C(\Delta\phi) &= FC^{\text{periph}}(\Delta\phi) + G\bigg\{1 + 2\sum_{n=2}^{5} v_{n,n}(p_{\text{T}}^{\text{a}}, p_{\text{T}}^{\text{b}})\cos(n\Delta\phi)\bigg\} \\ &= FC^{\text{periph}}(\Delta\phi) + Y^{\text{ridge}}(\Delta\phi). \end{split}$$

p_T dependence




- Direct Fourier decomposition: $v_n^{\text{2PC}}\{2\}$
- Template fit: $v_n^{\text{sub}}\{2\}$

Effect of subtraction (strongly p_T dependent):

$$v_2\{2\}\downarrow$$
, $v_3\{2\}\uparrow$, $v_4\{2\}\downarrow$


Centrality dependence



- $v_n^{\text{sub}}\{2\}$ is chosen as the central results
- When integrated over wide p_T range (0.5 ~ 5 GeV), the impact of non-flow is small



Non-flow at high p_T




IPGlasma+Hydro+PGCM:

- "Theory-1" vs "Theory-2" correspond to how much anisotropy from the glasma stage is retained in the hydro initialization
- ATLAS data provide input to further improve models




Other aux figures of the paper can be found at https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PAPERS/HION-2025-02/



- Flow signals in O+O and Ne+Ne resemble those seen in heavy-ion collisions
- Characteristic ordering observed: $v_2 > v_3$
- · Comparison with **Hydro+IPGlasma+PGCM** models clearly indicates a collective (hydrodynamical) response





Comparison of different ion collisions



- Flow in Ne+Ne consistently larger than in O+O
- · Light ions vs. heavy ions in centralities:
 - Light ion v_2 is much flatter
 - Light ion v_3 decreases from central to peripheral
 - Stronger fluctuation contributions in light ions








Compare to PGCM (initial conditions) + different pre-equilibrium evolution + Hydro (QGP phase):

- Trento: provides a better description of centrality dependence in central events
- IPGlasma: fails to reproduce the observed central-event trend



Giacolone et al, PRL 135 (2025) 012302 Mantysaari et al, PRL 135 (2025) 022302

Cross-experiment comparison




- Overall good consistency across experiments
- Small difference between experiments could arise from different selections

ATLAS: arXiv:2509.05171 (submit to PRC) ALICE: arXiv:2509.06428 (submit to PRL) CMS: arXiv:2510.02580 (submit to PRL)



Cross-experiment comparison



LHC HI working group:

- · Bridge experimental and theoretical HI communities ...
- Improve the uniformity of analysis choice ...
-



Conveners:

ALICE: Alice Ohlson, Alexander Kalweit


ATLAS: Qipeng Hu, Martin Rybar **CMS**: Austin Baty, Andre Stahl

LHCb: Giulia Manca, Saverio Mariani

Theory: Liliana Apolinario, Urs Wiedemann Reach all through ihc-hiwg-admin@cern.ch




Dijet asymmetry in O+O collisions




- Leading jet: $63 < p_T < 251$ GeV
- Sub-leading jet: $p_T > 20$ GeV
- Jet rapidities: |y| < 2.1
- Dijet alignment: $|\Delta \phi| > 7\pi/8$
- Unfolded in (p_T^1, p_T^2)





Intermediate leading p_T





- pp and peripheral O+O (60–80%) show similar distributions
- Central O+O (0–10%) events exhibit clear differences compared to pp and peripheral O+O



Centrality dependence

- In each x_J interval, data points are laterally shifted by centrality for visibility
- A smooth, systematic centrality dependence is observed from low $p_{\rm T}$ to high $p_{\rm T}$





Central




High leading p_T

Centrality dependence


• In each x_J interval, data points are laterally shifted by centrality for visibility

A smooth, systematic centrality dependence is observed Qualitatively consistent with hydrodynamical response, confirmed also by dijet measurements; v_n ratios reveal sensitivity to nuclear geometry (20Ne vs 16O)

While we cannot yet conclude that QGP is definitively formed in O+O and Ne+Ne collisions, the medium produced in these light ion systems exhibits behavior eripheral strikingly similar to that in heavy ion collisions




Low leading p_T



High leading p_T

Central

Future ATLAS and runs



LHC Run3 will end in 2026. An upgraded ATLAS will be used in Run4 (~ 2030s)

- High-granularity, high-coverage tracker (2.5 → 4.0)
- New ZDC (same as CMS Phase-II ZDC)
- High-granularity timing detector
- Replaced muon chambers
- New and upgraded forward and luminosity detector
- Improved trigger, high-performance software & computing, deeply embedded machine learning

ATLAS (maybe also CMS) has a strong preference to place p+Pb run in Run4 and dedicate 2026 to lighter ion runs (Ar, Kr, O, Ne)

First respond from LHC program committee: "this is not impossible"



Light ion workshop 2025

Light Ion workshop 2025

Dec 1 - 3, 2025, CERN

https://indico.cern.ch/event/1597414/

Organization:

Reyes Alemany Fernandez (CERN)

Giuliano Giacalone (CERN)

Qipeng Hu (USTC Hefei)

Gian Michele Innocenti (MIT)

Georgios Krintiras (University of Kansas)

Saverio Mariani (CERN)

Aleksas Mazeliauskas (ITP Heidelberg)


Dennis Perepelitsa (CU Boulder)

Anthony Timmins (University of Houston)

Wilke van der Schee (CERN)

Urs Wiedemann (CERN)

You Zhou (NBI Copenhagen)




Workshop topics:

- Experimental overviews of collectivity and energy loss
- Hydrodynamics and non-equilibrium dynamics
- Jet quenching in small systems
- nPDFs of light ions
- Synergies with nuclear structure physics and other areas
- Accelerator and experiments perspectives for future light-ion runs



Backup Slides

Comparison of different ion collisions — multiplicity bins



- Light ions vs. heavy ions in multiplicities:
 - O+O v₂ converges with p+Pb
 - Light ion v_3 align with heavy ion




Flow systematics

Source	harmonic order	0-40%	40–70%
		[%]	[%]
1. MC closure	v ₂ -v ₄	1	1
	v_2	0.5	0.5
2. Track selection	v_3	0.75	0.75
	v_4	1.5	1.5
3. Tracking efficiency	v ₂ -v ₄	0.25	0.25
	v_2	0.2	0.2–0.6
4. Centrality definition	v_3	0.2 - 1.0	1–2
	v_4	0.2	0.2–0.6
5. Residual pileup	v ₂ -v ₄	0.2	0.2
	v_2	0.25	0.25
6. Event-mixing	v_3	0.5	0.5
	v_4	1	1
	v_2	0.5	0.5–3
7. Peripheral reference	v_3	0.75–3.5	3.5–12
	v_4	1.0–4.5	4.5–20
8. Flattening procedure	<i>v</i> ₂ – <i>v</i> ₃	0.25	0.25



LHC heavy ion run schedule in 2026


2026-LHC-v0.5



Another heavy ion run (~3 weeks) at LHC scheduled in summer 2026 Pb+Pb or p+Pb



Non-flow in 4 particle cumulant — subevent method




$$c_n\{2\} = \langle\langle\{2\}_n\rangle\rangle,$$


$$c_n\{4\} = \langle\langle\{4\}_n\rangle\rangle - 2\langle\langle\{2\}_n\rangle\rangle^2$$


$$v_n\{4\} = \sqrt[4]{-c_n\{4\}}$$




- Additional event division further reduce non-flow, but with only small impact for p_{T} -integrated results
- Two subevent results are taken as nominal









Two particle correlation vs. four particle cumulant









 $v_2 (p + Pb) > v_2 (\gamma + Pb)$








- Could be understood as different longitudinal decorrelation and similar radial flow in the hydro picture (Zhao et al PRL 129 (2022) 252302)
- However, direct and resolved processes should be studied separately and it becomes possible





"Intrinsic Component"


Understand roles of fluctuations in initial conditions:

- Geometric fluctuations
- Intrinsic fluctuations


Moments of event-wise average p_T distribution in **ultra-central** Pb+Pb and Xe+Xe via n-particle momentum correlators:

$$c_n = \frac{\sum_{i_1 \neq \neq i_n} w_{i_1} ..w_{i_n} (p_{T,i_1} - \langle [p_T] \rangle) ... (p_{T,i_n} - \langle [p_T] \rangle)}{\sum_{i_1 \neq \neq i_n} w_{i_1} ... w_{i_n}}$$

$$k_2 = \frac{\langle c_2 \rangle}{\langle [p_{\rm T}] \rangle^2}, \quad k_3 = \frac{\langle c_3 \rangle}{\langle [p_{\rm T}] \rangle^3}, \quad \gamma = \frac{\langle c_3 \rangle}{\langle c_2 \rangle^{3/2}}, \quad \Gamma = \frac{\langle c_3 \rangle \langle [p_{\rm T}] \rangle}{\langle c_2 \rangle^2}.$$







 A phenomenological 2D Gaussian fluctuations predicts the trends well (R. Samanta et al. Phys. Rev. C 109 (2024) L051902)

