



粲偶素和奇特强子的蒙特卡洛模拟探究

Jin-Peng Zhang (张金鹏), Jian Cao (曹健), Wen-Chao Zhang*(张文超)

陕西师范大学

*wenchao.zhang@snnu.edu.cn

in collaboration with

Qian Yang, An-Ke Lei, Zhi-Lei She, Hua Zheng, Dai-Mei Zhou, Yu-Liang Yan, Ben-Hao Sa

Phys. Rev. C 109, 035201 (2024), Phys. Rev. C 110, 014910 (2024)


Phys. Rev. D 110, 054046 (2024), Phys. Rev. D 112, 014033 (2025)

@桂林, 26/10/2025

















Outline

- 1. J/ψ production
- 2. Exotic hadrons
- 3. Summary





- J/ψ is the lightest vector charmonium meson.
- The suppression of J/ψ production was proposed as a signal of QGP formation in HI collisions.
- The J/ψ production could also be suppressed due to the CNM effects, such as modifications of nuclear PDFs.
- In order to disentangle the hot and cold medium effects, it is necessary to understand the J/ψ production in pp collisions where the initial state effects are absent.





- The J/ψ production was extensively investigated at colliders such as the Tevatron, RHIC and LHC.
- The ALICE collaboration had published the inclusive J/ψ production in the fwd- and mid-rapidity regions in pp collisions at

 \sqrt{s} =2.76, 5.02, 7, 8 and 13 TeV.

Inclusive J/ψ

feed-down from heavier

charmonium decay

direct J/ψ

beauty hadrons weak decay

Phys. Lett. B 718, 295 (2012), JHEP 10, 084 (2019),

Phys. Lett. B 704, 442 (2011),

Eur. Phys. J. C 74, 2974 (2014),

Eur. Phys. J. C 76, 184 (2016),

Eur. Phys. J. C 81, 1121 (2021),

Eur. Phys. J. C 77, 392 (2017).

4





In order to investigate the interplay between the hard and soft mechanisms in particle productions, the ALICE collaboration had also published multiplicity dependence of J/ψ production in pp collisions at $\sqrt{s} = 5.02$, 7 and 13 TeV.

Phys. Lett. B 712, 165–175 (2012), Phys. Lett. B 810, 135758 (2020), JHEP 06, 015 (2022)


- Several theoretical approaches, such as CSM, COM and CEM have been utilized to describe the experimental data.
- They differ mostly in the treatment of non-perturbative evolution of the $c\bar{c}$ pair into the bound state J/ψ .






• The J/ψ production was also investigated by Monte Carlo

simulations Phys. Lett. B 712 (2012) 165–175










• The J/ψ production was also investigated by Monte Carlo

simulations

Different treatment of MPI in PYTHIA 6 and PYTHIA 8








- In our first work, we use a parton and hadron cascade model PACIAE 4.0 to investigate the J/ψ production in pp collisions at $\sqrt{s} = 5.02$, 7, and 13 TeV.
- We have also investigated the multiplicity dependence of J/ψ production in pp collisions at $\sqrt{s} = 5.02$, 7 and 13 TeV.





The PACIAE Model





PACIAE4.0 is based on PYTHIA 8.3 but further considers the partonic rescattering before hadronization and the hadronic rescattering after hadronization.





- In PACIAE 4.0, different charmonium production mechanisms are implemented.
- 1. In the perturbative scattering processes, leading order NRQCD channels via color-singlet and color-octet pre-resonant states are included.









- In PACIAE 4.0, different production mechanisms for the charmonium productions are implemented.
- 2. Charmonium can be produced from the cluster collapse mechanism.
- If c and \bar{c} are close in phase space the potential energy in the string is sufficient to create a light quark-antiquark pair, and instead the two heavy quarks bind into a charmonium bound state.
- 3. In the so-called non-prompt charmonium production, charmonia can be produced from the weak decay of a hadron containing a beauty quark.






- For the J/ψ hadronic rescattering, besides the elastic scattering, so far the following inelastic processes are considered.
- So far, the scattering of J/ψ with partons is not considered in PACIAE4.0 .

$$\begin{split} J/\psi + n &\to \Lambda_c^+ + D^-, \quad J/\psi + n \to \Sigma_c^+ + D^-, \\ J/\psi + n &\to \Sigma_c^0 + \bar{D}^0, \quad J/\psi + p \to \Lambda_c^+ + \bar{D}^0, \\ J/\psi + p &\to \Sigma_c^+ + \bar{D}^0, \quad J/\psi + p \to \Sigma_c^{++} + D^-, \\ J/\psi + \pi^+ \to D^+ + \bar{D}^{*0}, \quad J/\psi + \pi^- \to D^0 + D^{*-}, \\ J/\psi + \pi^0 \to D^0 + \bar{D}^{*0}, \quad J/\psi + \pi^0 \to D^+ + D^{*-}, \\ J/\psi + \rho^+ \to D^+ + \bar{D}^0, \quad J/\psi + \rho^- \to D^0 + D^-, \\ J/\psi + \rho^0 \to D^0 + \bar{D}^0, \quad J/\psi + \rho^0 \to D^+ + D^-. \end{split}$$




































 $< dN_{J/\psi} >$ and $< dN_{ch}/d\eta >$ are the average J/ψ yield and the charged particle multiplicity in minimum bias events.











is the differential cross section responsible for generating the J/ψ

the J/ψ mass

 $\cos \chi < 0$ hard radiation P.T process


 $cos \chi > 0$ soft radiation N.P.T process



















部分子强子化会弱化初始能量关联性


















• The naive constituent quark model has been the basic framework within which most of the hadronic states could be understood.

Volume 8, number 3


PHYSICS LETTERS

1 February 1964

Phys. Lett. 8, 214 (1964)

A SCHEMATIC MODEL OF BARYONS AND MESONS *

M. GELL-MANN
California Institute of Technology, Pasadena, California




Received 4 January 1964







• The non-Abelian property of quantum chromodynamics (QCD) permits the existence of new types of hadrons.



tetraquark

pentaquark

glueball

hybrid state


Phys. Rep. 389, 61 (2004); Phys. Rep. 454, 1 (2007); Prog. Part. Nucl. Phys. 63, 74 (2009).





• First exotic candidate X(3872)

```
PRL 110, 222001 (2013);
JHEP 04, 154 (2013);
JHEP 01, 117 (2017);
PRL 126, 092001 (2021);
PRL 128, 032001 (2022) ...
```



PRL 91, 262001 (2003)

Later, more experiments have confirmed its existence and have measured its properties in both pp and Pb-Pb collisions.






• However, the exact nature of X(3872) is still unclear.


The loose molecular state and compact tetraquark state are of

most interest.














- In our work, we will investigate the X(3872) production in pp collisions at $\sqrt{s} = 2.76$ TeV with PACIAE model.
- Two sicarios are considered: the tetraquark state and the molecular state containing $D\overline{D}^*$.







• In the quantum statistical mechanics inspired dynamically constrained phase-space coalescence (DCPC) model, the yield of *N*-particle cluster in six-dimensional phase space can be estimated by Phys. Rev. C 85, 024907 (2012).

$$Y_{cluster} = \int \cdots \int \frac{d\vec{q}_1 d\vec{p}_1 \cdots d\vec{q}_N d\vec{p}_N}{h^{3N}}$$

• For the $c\bar{c}qq$ tetraquark, its yield reads as

$$Y_{tetraquark} = \int \cdots \int \, \delta_{1234} \, \frac{d\vec{q}_1 d\vec{p}_1 \cdots d\vec{q}_4 d\vec{p}_4}{h^{12}}$$

$$\delta_{1234} = \begin{cases} if \ 1 = c, 2 = \bar{c}, 3 = q, 4 = \bar{q} \\ 1. \ m_{X(3872)} - \Delta m < m_{inv} < m_{X(3872)} - \Delta m, R \le 1 fm \\ 0. \ otherwise \end{cases}$$

$$\Delta m = \frac{\Gamma}{2} = 1.95 \text{MeV}/c^2$$
, R is the radius of the cluster. 28





• In the quantum statistical mechanics inspired dynamically constrained phase-space coalescence (DCPC) model, the yield of *N*-particle cluster in six-dimensional phase space can be estimated by Phys. Rev. C 85, 024907 (2012).

$$Y_{cluster} = \int \cdots \int \frac{d\vec{q}_1 d\vec{p}_1 \cdots d\vec{q}_N d\vec{p}_N}{h^{3N}}$$

• For the $D\overline{D}^*$ molecular state, its yield reads

as
$$Y_{molecule} = \int \cdots \int \delta_{12} \frac{d\vec{q}_1 d\vec{p}_1 d\vec{q}_2 d\vec{p}_2}{h^6}$$


$$\delta_{12} = \begin{cases} if \ 1 = D, 2 = \overline{D}^* \\ m_{X(3872)} - \Delta m < m_{inv} < m_{X(3872)} - \Delta m, 1 \le R \le 2fm \\ 0. \end{cases}$$
 otherwise

$$\Delta m = \frac{\Gamma}{2} = 1.95 \text{MeV}/c^2$$
, R is the radius of the cluster. 29






• In this work, tetraquark- and molecule-state of X(3872) are assumed to be coalesced in PFS and HFS, respectively.








Phys. Rev. C 110, 014910 (2024)









Lattice QCD predictions

 0^{++} ground state: 1.5 - 1.7 GeV/c²


 2^{++} ground state: 2.3 - 2.4GeV/c²

 0^{-+} ground state: 2.3 - 2.6GeV/c²

Phys. Lett. B **309**, 378 (1993); Phys. Rev. D **60**, 034509 (1999); Phys. Rev. D **73**, 014516 (2006); J. High Energy Phys. 10 (2012) 170; Phys. Rev. D **100**, 054511 (2019)








X(2370) was observed in $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$ decays with a statistical significance of 6.4 σ by the BESIII experiment






Exotic hadrons-X(2370) Eur. Phys. J. C (2020) 80:746



It was further observed from the combined measurement of $J/\psi \to \gamma K^+ K^- \eta'$ and $J/\psi \to \gamma K_s^0 K_s^0 \eta'$ with a statistical significance of 8.3 σ by BESIII







PRL 132, 181901 (2024)

The statistical significance of the X(2370) is greater than 6.4σ and the spin parity is determined to be 0^{-+} for the first time





- Some theoretical studies interpret X(2370) states as the fourth radial excitation of η/η' . PRD 83, 114007 (2011); PRD 102, 114034 (2020)
- Some theoretical studies also suggest that the X(2370) states are the P-wave $ss\bar{s}\bar{s}$ tetraquark states of $J^{PC}=0^{-+}$. PRD 106, 014023 (2022)
- Some theoretical studies indicate that X(2370) states can be explained as light baryonium states $\Lambda \overline{\Lambda}$ and $\Sigma \overline{\Sigma}$ with quantum numbers of 0^{-+} and 1^{--} . PRD 105, 014016 (2022)





• Among the theoretical studies, one of the intriguing explanations is a pseudoscalar glueball




Phys. Rev. D **100**, 054511 (2019); PRD 82, 074026 (2010); PRD 87, 054036 (2013); Phys. Lett. B **827**, 136960 (2022); Nucl. Phys. A 728 (2003) 165–181; Phys. Lett. B **3** 7 (2006) 53–61






- Glueball productions: Int J Mod Phys E, 18, 1-49 (2009)
- J/ψ decays
- Double pomeron exchange in pp collisions
- $p\bar{p}$ annihilation










- In our work, we will investigate the glueball-like particle (X(2370)) production in pp collisions at $\sqrt{s} = 13$ TeV with PACIAE model.
- Three sicarios are considered: the glueball states containing two valence gluons, the tetraquark state $ss\bar{s}\bar{s}$, the molecular state $\Lambda \bar{\Lambda} / \Sigma \bar{\Sigma}$, or $\pi^+ \pi^- \eta' / K^+ K^- \eta' / K^0_S K^0_S \eta'$.















In this work, glueball-like state, tetraquark- and molecule-state of X(2370) are assumed to be coalesced in PFS and HFS, respectively.
















Exotic hadrons-G(3900)



在BESIII能区下利用PACIAE模型研究类粲偶素奇特强子的产生

Phys. Rev. D 112, 014033 (2025)





(3+1)-D nonextensive hydrodynamics

PHYSICAL REVIEW D 111, 036010 (2025)

Nonextensive (3+1)-dimensional hydrodynamics for relativistic heavy-ion collisions

Jia-Hao Shi[®], ¹ Zhi-Ying Qin, ¹ Jin-Peng Zhang[®], ¹ Jian Cao[®], ¹ Ze-Fang Jiang[®], ^{2,3} Wen-Chao Zhang[®], ^{1,*} and Hua Zheng[®]

¹School of Physics and Information Technology, Shaanxi Normal University, Xi'an 710119, China ²Department of Physics and Electronic-Information Engineering, Hubei Engineering University, Xiaogan, Hubei 432000, China

³Institute of Particle Physics and Key Laboratory of Quark and Lepton Physics (MOE), Central China Normal University, Wuhan, Hubei 430079, China



(Received 22 August 2024; accepted 5 January 2025; published 6 February 2025)

Phys. Rev. D 111, 036010 (2025)





QCD phase transition

QCD phase transition at finite temperature and chemical potential with the non-extensive statistics

Zhi-Ying Qin,^{1,2} Jia-Hao Shi,¹ Jin-Peng Zhang,¹ Jian Cao,¹ Bo Feng,¹ Wen-Chao Zhang,^{1,*} Hua Zheng,¹ and Shi-Jun Mao³

¹School of Physics and Information Technology, Shaanxi Normal University, Xi'an 710119, China ²School of Physics Science and Engineering, Tongji University, Shanghai 200092, China ³School of Physics, Xi'an Jiaotong University, Xi'an, Shaanxi 710049, China (Dated: September 30, 2025)

Accepted by Phys. Rev. D, DOI: 10.1103/4gzt-46t5






Conclusions

- We have investigated the J/ψ productions in pp collisions at $\sqrt{s} = 5.02$, 7 and 13 TeV with PACIAE 4.0.
- We have also investigated the multiplicity dependence of J/ψ production in pp collisions at $\sqrt{s} = 5.02$, 7 and 13 TeV.
- The contributions of the J/ψ productions from the NRQCD, cluster collapse and the weak decay from the B hadrons are analyzed in detail.
- The effect of the partonic and hadronic rescattering effects on the J/ψ production has been presented.
- The hadronization process has an obvious effect on the J/ψ energy correlator.
- Discrepancy between the glueball-, tetraquark- or molecule-state of the exotic hadrons in the rapidity and p_T distributions can be served as a criterion for identifying the configuration.





Back up







Back up

$$\frac{dN_{J/\psi}}{\langle dN_{J/\psi} \rangle} = \frac{N_{J/\psi}^{i}}{N_{J/\psi}^{tot}} \cdot \frac{N_{evt}^{tot}}{N_{evt}^{i}}$$

 $N_{J/\psi}^{i}$ and N_{evt}^{i} are the number of J/ ψ and number of events in ith multiplicity bin, respectively. $N_{J/\psi}^{tot}$ and N_{evt}^{tot} are the total number of J/ ψ produced and total number of minimum bias events, respectively.