



Flow measurements at LHCb experiment

Jianqiao Wang on behalf of the LHCb collaboration

October 26, 2025




- Physics background
- 2 Flow in 5.02 TeV PbPb
- 3 Flow in small systems
- 4 Flow in fixed-target PbNe and PbAr
- **1** Light-ion data taking
- 6 Summary and Prospects

2/16

Standard Model for heavy-ion collisions


• Time evolution of heavy-ion collisions



3/16

Standard Model for heavy-ion collisions


• Time evolution of heavy-ion collisions



• Fluid-like QGP created in PbPb collisions

Flow

- Hydrodynamic and thermal expansion of medium can lead to particle collectivity
- Flow considered as one of the key signatures of QGP formation



• Characterized by Fourier coefficients v_n :

$$E\frac{\mathrm{d}^3 N}{\mathrm{d}p^3} = \frac{1}{2\pi} \frac{\mathrm{d}^2 N}{p_{\mathrm{T}} \mathrm{d}p_{\mathrm{T}} \mathrm{d}y} \left(1 + 2 \sum_{1}^{\infty} v_n \cos[n(\phi - \Psi_{\mathrm{R}})] \right)$$

- \triangleright v_1 : direct flow
- \triangleright v_2 : elliptic flow
- \triangleright v_3 : triangular flow






4/16


LHCb detector in Run2

- A single-arm general-purpose forward spectrometer, covering the pseudo-rapidity range of $2 < \eta < 5$.
- Excellent tracking and particle identification capabilities
- Complementary to other LHC experiments: forward rapidity, unique fixed-target mode, high-purity heavy flavor signals for flow measurements



5/16


LHCb detector in Run3



- Upgraded tracking and trigger system to operate at higher luminosities
- More central data accessible for heavy-ion collisions
 - ▶ Up to 30% for PbPb, full-centrality for light-ion

6/16

LHCb detector in Run3




- Upgraded tracking and trigger system to operate at higher luminosities
- More central data accessible for heavy-ion collisions
 - ▶ Up to 30% for PbPb, full-centrality for light-ion
- New SMOG2 system for fixed-target program
 - Running simultaneously with collider mode
 - ▶ Unique energy range $\sqrt{s_{\rm NN}} \sim 30\text{--}110\,\text{GeV}$ and very backward rapdity
 - ▶ Wider choice of gas (H₂, He, Ne, Ar)

6/16


Flow at forward rapidity

• Flow measurement with two-particle correlation with Run2 PbPb data at $\sqrt{s_{\mathrm{NN}}} = 5.02 \,\mathrm{TeV}$



- Near-side ridge observed in (peripheral) PbPb at forward rapidity, similar to midrapidity results
- Lower ridge for more peripheral collisions
- Non-flow subtracted by excluding jet region


Flow in 5.02 TeV PbPb



- Smaller $v_2(p_T)$ and $v_3(p_T)$ compared to ALICE and ATLAS results at midrapidity, possibly due to the dominant freeze-out phase at forward rapidity
- AMPT overestimates v_n at forward rapidity

8/16

Flow in 5.02 TeV PbPb




- Smaller $v_2(p_T)$ and $v_3(p_T)$ compared to ALICE and ATLAS results at midrapidity, possibly due to the dominant freeze-out phase at forward rapidity
- AMPT overestimates v_n at forward rapidity
- Validation with cumulant method
 - Good agreement between two methods
 - Better non-flow subtraction with cumulant method


8/16

Flow in pPb

- Explanation for the QGP signature in high-multiplicity small systems
 - ▶ QGP droplet formation? Transport?
 - ► Glasma?

• Unique coverage to investigate possible gluon saturation






9/16


Flow in pPb

- Explanation for the QGP signature in high-multiplicity small systems
 - ▶ QGP droplet formation? Transport?
 - ► Glasma?

• Long-range correlations observed in 5 TeV pPb

Phys.Lett.B762(2016)473-483






• No flow extraction unfortunately

9/16

Flow in $8.16 \,\mathrm{TeV}\ p\mathrm{Pb}$


- \bullet More detailed analysis and higher statistics with Run2 pPb
- Improved non-flow suppression compared to PbPb: large η gap, jet-yield subtracted method




- Rising v_2 and flat v_3 trends as a function of charged particle density
- Consistent result for both rapidities. No Bjorken-x dependence observed
- Final state effects dominant?

10 / 16

Flow in $8.16 \,\mathrm{TeV}\ p\mathrm{Pb}$






- \bullet Compared with (3+1)D hydrodynamic model, which calibrated with pp results at different energies
- Clearly overestimates v_2
 - ► Issue of fluid parametrization or hydrodynamics itself in small systems?

Jianqiao Wang Flow at LHCb October 26, 2025 11/16

Initial geometry

• Other knowledge from flow measurements




Nuclear shape

• Anisotropy initial shape can be converted to collevtive motion in momentum space



- Different nuclear structure predictions
 - ► Ne: strongly deformed bowling pin shape
 - ▶ O, Ar: nearly spherical
- Large v_2 difference expected between the most central AA collisions
- Even more pronounced differences in fixed-target systems!





Phys. Rev. Lett. 135,



Flow in fixed-target PbNe and PbAr

- Flatter v_2 for PbNe compared with PbAr and similar decreasing v_3
- Significant enhanced ($\sim 40\%$) v_2 ratios between PbNe and PbAr as collision goes central






14 / 16

Flow in fixed-target PbNe and PbAr


- Flatter v_2 for PbNe compared with PbAr and similar decreasing v_3
- Significant enhanced ($\sim 40\%$) v_2 ratios between PbNe and PbAr as collision goes central
- Trend well described by 3+1D hydro prediction with ab-initio nuclear-structure (PGCM and NLEFT), suggesting the elongated Neon shape and the validity of hydrodynamic description in these systems



14/16

Flow in fixed-target PbNe and PbAr


- Flatter v_2 for PbNe compared with PbAr and similar decreasing v_3
- Significant enhanced ($\sim 40\%$) v_2 ratios between PbNe and PbAr as collision goes central
- Trend well described by 3+1D hydro prediction with ab-initio nuclear-structure (PGCM and NLEFT), suggesting the elongated Neon shape and the validity of hydrodynamic description in these systems
- Improved centrality calibration and updated models will be implemented for final results




14/16

Light-ion data taking

• Both collider and fixed-target data from light-ion runs



- Excellent data collected in OO and NeNe run
- Search QGP signature in small systems and study the *transition* from large system
- Fixed-target OH₂ and NeNe data simultaneously with collider mode
- Unique opportunity for cosmic ray and nuclear imaging



Summary and Prospects


- Growing contributions on flow results from LHCb experiment
 - \triangleright Significant charged particle v_2 in pPb and PbPb collisions at forward rapidity, smaller than that observed at midrapidity
 - \blacktriangleright Hydrodynamic model overestimates v_2 in $p{\rm Pb}$ collisions, more studies needed to understand the descrepancy
 - ► First preliminary result with upgraded SMOG2 system, suggesting the ²⁰Ne nucleus deformation
- Outlook
 - Further measurements (e.g. $v_n p_T$ correlation) for studying cluster structures of light nuclei
 - ▶ Excellent data from light-ion run both for colliding and fixed-target modes

16 / 16


Thanks for your attention!

Cumulant method

• Cumulant method: nth order cumulant $c_n\{m\}$ can be derived from Q_n vectors



$$v_2{2} = (c_2{2})^{1/2}, \quad v_2{4} = (-c_2{4})^{1/4}$$





- Non-flow effects with suppressed with more particles correlated
- Results directly come from the event-by-event measurement

4 □ ト 4 □ ト 4 亘 ト 4 亘 ト 9 Q ○

2/3

Two-particle correlation





- Two-particle correlation: $(\Delta \eta, \Delta \phi)$ correlation function
- Obtained by comparing distributions same-event and mixed-event pairs

$$\frac{1}{N_{\text{trig}}} \frac{\mathrm{d}^2 N^{\text{pair}}}{\mathrm{d}\Delta \eta \ \mathrm{d}\Delta \phi} = B(0,0) \times \frac{S(\Delta \eta, \Delta \phi)}{B(\Delta \eta, \Delta \phi)}$$

- Long-range correlation ridge symbolises the positive v_2
- Significant ridge and jet (non-flow) structures
- What about forward rapidity?