



北京航空航天大学
BEIHANG UNIVERSITY

When **inverse seesaw** meets **inverse EWPT**: a novel path to leptogenesis

Ke-Pan Xie (谢柯盼)

Beihang University

2025.9.21@Dalian, 第四届高能物理理论与实验融合发展研讨会

With Wen-Yuan Ai and Peisi Huang, 2509.xxxxx

A mystery and an elegant solution

The baryon-matter asymmetry of the Universe


$$Y_B = \frac{n_B}{s} \approx 8.58 \times 10^{-11}$$

Calling for physics beyond the SM



Leptogenesis [Yanagida et al, Phys.Lett.B 174 \(1986\) 45-47](#)

- Right-handed neutrino ν_R , explaining SM neutrino mass
- CP-violating decay, $\Gamma_{\nu_R \rightarrow \ell H} > \Gamma_{\nu_R \rightarrow \bar{\ell} H^*}$




Lepton asymmetry Y_L generated
SM EW sphaleron process partly
converts $Y_L \rightarrow Y_B$


Solving the **baryon** mystery via the **lepton** dynamics!

Initial condition

Thermal leptogenesis




Typically described by a set of Boltzmann equations




Buchmuller *et al*,
Annals Phys. 315 (2005) 305-351

Initial condition

Thermal leptogenesis



Nonthermal leptogenesis




Scenarios:


1. Inflaton decay $\phi \rightarrow \nu_R \nu_R$ during reheating or preheating
2. Cosmic first-order phase transitions (FOPTs) [this talk]

What is a cosmic FOPT?

Decay of the Universe between two vacuums




Bubble expansion velocity v_w




Why can it produce heavy particles with $m \gg T_*$?

Plasma frame



Wall-rest frame




If $v_w \approx 1$, meaning $\gamma_w \gg 1$:

The bubble can touch a scale up to $\gamma_w T_* \gg T_*$!

Bubble-assisted leptogenesis


Previous work:

Mass-gain: $\phi \bar{\nu}_R^c \nu_R$




Huang and KPX, JHEP 09 (2022) 052

Light-to-heavy: $\phi \bar{\chi}_L \nu_R$



Azatov et al, JHEP 10 (2021) 043

Bubble collisions




Cataldi et al, JCAP 11 (2024) 047

- New physics scalar ϕ with $T_* \gtrsim 10^7$ GeV
- Supercooling $T_* \ll w_*$, and $\gamma_w \gg 1$ driven by large vacuum energy

Bubble-assisted leptogenesis


Previous work:

Mass-gain: $\phi \bar{\nu}_R^c \nu_R$




Huang and KPX, JHEP 09 (2022) 052

Light-to-heavy: $\phi \bar{\chi}_L \nu_R$



Azatov et al, JHEP 10 (2021) 043

Bubble collisions



Cataldi et al, JCAP 11 (2024) 047

- New physics scalar ϕ with $T_* \gtrsim 10^7$ GeV
- Supercooling $T_* \ll w_*$, and $\gamma_w \gg 1$ driven by large vacuum energy



This work:

- The SM Higgs h with an EW phase transition at $T_* \sim 10^2$ GeV
- NOT a supercooled transition, but with $\gamma_w \gg 1$

Leptogenesis from an EWPT


A first-order EWPT: triggered by physics beyond the SM

See for example [Wang et al, Phys.Rev.D 87 \(2013\) 2, 023509](#)



Fatal difficulty


- EW sphaleron $\mathcal{O}_{\text{eff}} = \prod_i [\bar{q}_L^{c,i} q_L^i] [\bar{\ell}_L^{c,i} \ell_L^i]$ [Kuzmin et al, Phys.Lett.B 155 \(1985\) 36](#)
- Reaction rate $\propto e^{-8\pi\langle h \rangle / (g_W T_*)}$
- Inside the bubble $\langle h \rangle \sim T_*$
- Suppression $\propto e^{-38.8} \approx 10^{-17} \ll 1$
- $Y_L \rightarrow Y_B$ forbidden!!



Leptogenesis from an EWPT

A first-order EWPT: triggered by physics beyond the SM

See for example [Wang et al, Phys.Rev.D 87 \(2013\) 2, 023509](#)




To resolve this, we consider an **inverse** EWPT:

From $\langle h \rangle \neq 0$ to $\langle h \rangle = 0$!

Fatal difficulty




- EW sphaleron $\mathcal{O}_{\text{eff}} = \prod_i [\bar{q}_L^{c,i} q_L^i] [\bar{\ell}_L^{c,i} \ell_L^i]$ [Kuzmin et al, Phys.Lett.B 155 \(1985\) 36](#)
- Reaction rate $\propto e^{-8\pi\langle h \rangle / (g_W T_*)}$
- Inside the bubble $\langle h \rangle \sim T_*$
- Suppression $\propto e^{-38.8} \approx 10^{-17} \ll 1$
- $Y_L \rightarrow Y_B$ forbidden!!




The thermal history

Vectorlike leptons (VLLs) $F = (F^0, F^-)^T \sim \mathbf{2}_{-1/2}; N \sim \mathbf{1}_0; E \sim \mathbf{1}_{-1}$


$$-y_{N_L} \bar{F}_R \tilde{H} N_L - y_{N_R} \bar{F}_L \tilde{H} N_R - y_{E_L} \bar{F}_R H E_L - y_{E_R} \bar{F}_L H E_R$$



(Transition duration)/(Hubble time)




(Latent heat)/(Radiation energy)



Inverse EWPT

The inverse EWPT

Not supercooled ($T_{n1} \approx v_{n1}$), but features relativistic bubble walls



SM particles *aspirated* by the plasma, $v_w \approx 1$

Quantitatively, $\gamma_w \sim 300$ [Azatov et al, JHEP 12 \(2024\) 056](#)

The 2nd-generation: VLL'

Higgs portal: $-y'_{N_L} \bar{F}'_R \tilde{H} N'_L - y'_{N_R} \bar{F}'_L \tilde{H} N'_R - y'_{E_L} \bar{F}'_R H E'_L - y'_{E_R} \bar{F}'_L H E'_R$

$U(1)_{B-L}$ -breaking: $-\frac{\mu_N}{2} \bar{N}'_L N'^c_L - \lambda_N \bar{F}'_L \tilde{H} N'^c_L$

SM lepton interactions: $-\sum_{\alpha} y_D^{\alpha} \bar{\ell}_L^{\alpha} \tilde{H} N'_R - \sum_{\alpha} \mu_D^{\alpha} \bar{\ell}_L^{\alpha} F'_R - \sum_{\alpha} \lambda_D^{\alpha} \bar{\ell}_L^{\alpha} H E'_R$



SM lepton doublet

$\alpha = e, \mu, \tau$

Reminder:

$$F' = \begin{pmatrix} F'^0 \\ F'^- \end{pmatrix} \sim \mathbf{2}_{-1/2}; \quad N' \sim \mathbf{1}_0; \quad E' \sim \mathbf{1}_{-1}$$

Same charge with ℓ_L

Pure singlet

Same charge with e_R

The 2nd-generation: VLL'

Higgs portal: $-y'_{N_L} \bar{F}'_R \tilde{H} N'_L - y'_{N_R} \bar{F}'_L \tilde{H} N'_R - y'_{E_L} \bar{F}'_R H E'_L - y'_{E_R} \bar{F}'_L H E'_R$

$U(1)_{B-L}$ -breaking: $-\frac{\mu_N}{2} \bar{N}'_L N'^c_L - \lambda_N \bar{F}'_L \tilde{H} N'^c_L$

SM lepton interactions: $-\sum_{\alpha} y_D^{\alpha} \bar{\ell}_L^{\alpha} \tilde{H} N'_R - \sum_{\alpha} \mu_D^{\alpha} \bar{\ell}_L^{\alpha} F'_R - \sum_{\alpha} \lambda_D^{\alpha} \bar{\ell}_L^{\alpha} H E'_R$

After EWSB, let $N'_R \rightarrow \nu_R^1$, $F_R'^0 \rightarrow \nu_R^2$, and $N'_L \rightarrow S_L^1$, $F_L'^0 \rightarrow S_L^2$

$$-\frac{1}{2} \begin{pmatrix} \bar{\nu}_L & \bar{\nu}_R^c & \bar{S}_L \end{pmatrix} \begin{pmatrix} 0_{3 \times 3} & m_D & 0_{3 \times 2} \\ m_D^T & 0_{2 \times 2} & M_R \\ 0_{2 \times 3} & M_R^T & \mu \end{pmatrix} \begin{pmatrix} \nu_L^c \\ \nu_R \\ S_L^c \end{pmatrix} \quad \begin{array}{l} \nu_L^{\alpha}, \text{ with } \alpha = e, \mu, \tau \\ \nu_R^i \text{ and } S_L^i, \text{ with } i = 1, 2 \end{array}$$

Mass matrices $m_D^{\alpha i} = \left(\frac{y_D^{\alpha} v}{\sqrt{2}} \quad \mu_D^{\alpha} \right)$, $M_R^{ij} = \begin{pmatrix} m'_N & \frac{y'_{N_R} v}{\sqrt{2}} \\ \frac{y'_{N_L} v}{\sqrt{2}} & m'_F \end{pmatrix}$, and $\mu_{ij} = \begin{pmatrix} \mu_N & \frac{\lambda_N v}{\sqrt{2}} \\ \frac{\lambda_N v}{\sqrt{2}} & 0 \end{pmatrix}$

The 2nd-generation: VLL'

Higgs portal: $-y'_{N_L} \bar{F}'_R \tilde{H} N'_L - y'_{N_R} \bar{F}'_L \tilde{H} N'_R - y'_{E_L} \bar{F}'_R H E'_L - y'_{E_R} \bar{F}'_L H E'_R$

$U(1)_{B-L}$ -breaking: $-\frac{\mu_N}{2} \bar{N}'_L N'^c_L - \lambda_N \bar{F}'_L \tilde{H} N'^c_L$

SM lepton interactions: $-\sum_{\alpha} y_D^{\alpha} \bar{\ell}_L^{\alpha} \tilde{H} N'_R - \sum_{\alpha} \mu_D^{\alpha} \bar{\ell}_L^{\alpha} F'_R - \sum_{\alpha} \lambda_D^{\alpha} \bar{\ell}_L^{\alpha} H E'_R$

After EWSB, let $N'_R \rightarrow \nu_R^1$, $F_R'^0 \rightarrow \nu_R^2$, and $N'_L \rightarrow S_L^1$, $F_L'^0 \rightarrow S_L^2$

$$-\frac{1}{2} \begin{pmatrix} \bar{\nu}_L & \bar{\nu}_R^c & \bar{S}_L \end{pmatrix} \begin{pmatrix} 0_{3 \times 3} & m_D & 0_{3 \times 2} \\ m_D^T & 0_{2 \times 2} & M_R \\ 0_{2 \times 3} & M_R^T & \mu \end{pmatrix} \begin{pmatrix} \nu_L^c \\ \nu_R \\ S_L^c \end{pmatrix} \quad \begin{array}{l} \nu_L^{\alpha}, \text{ with } \alpha = e, \mu, \tau \\ \nu_R^i \text{ and } S_L^i, \text{ with } i = 1, 2 \end{array}$$

Mass matrices $m_D^{\alpha i} = \left(\frac{y_D^{\alpha} v}{\sqrt{2}} \quad \mu_D^{\alpha} \right)$, $M_R^{ij} = \begin{pmatrix} m'_N & \frac{y'_{N_R} v}{\sqrt{2}} \\ \frac{y'_{N_L} v}{\sqrt{2}} & m'_F \end{pmatrix}$, and $\mu_{ij} = \begin{pmatrix} \mu_N & \frac{\lambda_N v}{\sqrt{2}} \\ \frac{\lambda_N v}{\sqrt{2}} & 0 \end{pmatrix}$

Minimal inverse seesaw [Xing et al, Phys. Lett. B 679 \(2009\) 242–248](#)

- Lightest SM ν massless, and $m_{\nu} \sim 0.1 \text{ eV} \times \left(\frac{m_D}{\text{GeV}} \right)^2 \left(\frac{\mu}{0.1 \text{ MeV}} \right) \left(\frac{\text{TeV}}{M_R} \right)^2$
- **Near-degenerate** heavy Majorana neutrinos χ with $m_{\chi} \sim M_R \pm \frac{\mu}{2}$

The 2nd-generation: VLL'

Higgs portal: $-y'_{N_L} \bar{F}'_R \tilde{H} N'_L - y'_{N_R} \bar{F}'_L \tilde{H} N'_R - y'_{E_L} \bar{F}'_R H E'_L - y'_{E_R} \bar{F}'_L H E'_R$

$U(1)_{B-L}$ -breaking: $-\frac{\mu_N}{2} \bar{N}'_L N'^c_L - \lambda_N \bar{F}'_L \tilde{H} N'^c_L$

SM lepton interactions: $-\sum_{\alpha} y_D^{\alpha} \bar{\ell}_L^{\alpha} \tilde{H} N'_R - \sum_{\alpha} \mu_D^{\alpha} \bar{\ell}_L^{\alpha} F'_R - \sum_{\alpha} \lambda_D^{\alpha} \bar{\ell}_L^{\alpha} H E'_R$

Neutrino mass origin


Crucial for RHN production



Bubble-assisted leptogenesis

Light-to-heavy transition of $\ell_L \rightarrow E'_R$ on the wall

$\gamma_w \sim 300$, ℓ_L energy in wall frame ~ 60 TeV




Calculated by the Higgs background field theory [Bodeker et al, JCAP 05 \(2017\) 025](#)

Bubble-assisted leptogenesis

Assuming a mass hierarchy $m'_E > m'_F > m'_N$

Cascade decays among the vectorlike leptons




End of the vectorlike lepton decay chain: lightest Majorana $\chi_{1,2}$

Bubble-assisted leptogenesis

$m_\chi \gg T_{n1}$, instant decay of $\chi \rightarrow \ell H / \bar{\ell} H^*$ via seesaw vertices

CP violation, generating Y_L Adhikary et al, Phys. Rev. D 93 (2016) 113001



Active EW sphaleron inside the bubble: $Y_L \rightarrow Y_B$

χ far away from equilibrium, **no thermal washout effects**

Parameter space

Baryon asymmetry generated

Decay asymmetry χ produced via EWPT (inherits from E')

$$Y_B \approx -\frac{28}{79} \epsilon f_{\text{dec}} \frac{n_{E'}^{\text{trans.}}}{s(T_{n1})} \exp \left\{ - \int_{m'_N/T_{n1}}^{m'_N/T_{n2}} \frac{z'^4 dz'}{s(m'_N) H(m'_N)} \frac{\sum_i \gamma_{\chi_i}(z')}{2Y_\ell^{\text{eq}}} \right\}$$

$Y_L \rightarrow Y_B$ factor Fraction of decay Thermal washout, suppressed by $m'_N \gtrsim 20T_{n1}$

Parameter space

Baryon asymmetry generated


Decay asymmetry χ produced via EWPT (inherits from E')

$$Y_B \approx -\frac{28}{79} \epsilon f_{\text{dec}} \frac{n_{E'}^{\text{trans.}}}{s(T_{n1})} \exp \left\{ - \int_{m'_N/T_{n1}}^{m'_N/T_{n2}} \frac{z'^4 dz'}{s(m'_N) H(m'_N)} \frac{\sum_i \gamma_{\chi_i}(z')}{2Y_\ell^{\text{eq}}} \right\}$$

$Y_L \rightarrow Y_B$ factor Fraction of decay Thermal washout, suppressed by $m'_N \gtrsim 20T_{n1}$

- Casas-Ibarra: $m_D = U_{\text{PMNS}} m_n^{1/2} R \mu^{-1/2} M_R^T$, yields oscillation data
- CP violation dominated by the new physics phase from $\mu^{-1/2}$

Mass eigenstates after EWSB



Conclusion

A novel leptogenesis mechanism realized by the vectorlike leptons via the **Higgs field** *electroweak phase transition*

1st-generation: F , N , and E

- Lighter mass $m_{F,N,E} \sim \text{TeV}$
- Stronger Yukawa $y_{N_{L,R},E_{L,R}} \sim 3$
- **Inverse EWPT**: not supercooled, but $\gamma_w \gg 1$

2nd-generation: F' , N' , and E'

- Heavier mass $m'_{F,N,E} \sim 4 \text{ TeV}$
- Weaker Yukawa $y'_{N_{L,R},E_{L,R}} \sim 1$
- **Inverse seesaw**: neutrino mass origin and nonthermal **leptogenesis**


Phenomenology: Higgs measurement (e.g., $h \rightarrow \gamma\gamma$, $h \rightarrow \tau^+\tau^-$), heavy lepton searches, gravitational waves




Thank you!

Backup: check list


After production, we want this



Not this




For the Majorana χ , we want this



CP-violating decay (generating Y_L)
from seesaw vertices $\sim \mathcal{O}(10^{-5})$

Not this



Annihilation (no Y_L generated)
from $y'_{N_{L,R}} \sim \mathcal{O}(1)$

Decay products ℓ and H should thermalize with the plasma rapidly to prevent **nonthermal washout** from $\ell H \rightarrow \chi \rightarrow \bar{\ell} H^*$

Thermal washout from $\ell H \rightarrow \chi \rightarrow \bar{\ell} H^*$ is suppressed by $m_\chi \gtrsim 20T_{n1}$