



The low-lying hadron states in Λ_c^+ decays

谢聚军

中国科学院近代物理研究所

2025年8月12-16日 @ “粲强子衰变和标准模型的精确检验” 2025年夏季年会，贵州贵阳

Outline

Introduction

Possible $\Sigma_{1/2-}^$ state in $\Lambda_c^+ \rightarrow \eta\pi^+\Lambda$ decay*


Study of $N^(1535)$ in the $\Lambda_c^+ \rightarrow \bar{K}^0\eta p$ decay*

Possible ϕp state in $\Lambda_c^+ \rightarrow \pi^0\phi p$ decay


Summary

$$\Lambda_c^+$$

The lightest charmed baryon






Strange baryons ($\Lambda[uds]$)
 $m_u, m_d \approx m_s \rightarrow (qqq)$ uniform



Charmed baryon ($\Lambda_c[udc]$)
 $m_u, m_d \ll m_c \rightarrow$ **diquark + quark**
 (qq) (Q)

Weak decay



CERN COURIER

International Journal of High-Energy Physics

Sign in | Forgotten your password? | Sign up

CERN COURIER

brightrecruits.com/job/8901/senior-research-positions-ihep-cas-china

Latest Issue | [Archive](#) | [Jobs](#) | [Links](#) | [Buyer's guide](#) | [White papers](#) | [Events](#) | [Contact us](#)

Search

REGISTER NOW

Register as a member of cerncourier.com and get full access to all features of the site. Registration is free.

LATEST CERN COURIER ARTICLES

- ▶ Sneeze dynamics
- ▶ The longest proof
- ▶ Electron-hole collider
- ▶ Imaging with muons
- ▶ Towards a nuclear clock

SHARE THIS




CERN COURIER

Mar 18, 2016

BESIII makes first direct measurement of the Λ_c at threshold

The charmed baryon, Λ_c , was first observed at Fermilab in 1976. Now, 40 years later, the Beijing Spectrometer (BESIII) experiment at the Beijing Electron-Positron Collider II (BEPCII) has measured the absolute branching fraction of $\Lambda_c^+ \rightarrow pK^-\pi^+$ at threshold for the first time.



Beam-constrained mass distribution

Because the decays of the Λ_c^+ to hadrons proceed only through the weak interaction, their branching fractions are key probes for understanding weak interactions inside of a baryon. In particular, precise measurements of the decays of the Λ_c^+ will provide important information on the final-state strong interaction in the charm sector, thereby improving the understanding of quantum chromodynamics in the non-perturbative energy region. In addition, because most of the

DIGITAL EDITION

CERN Courier is now available as a regular digital edition. Click [here](#) to read the digital edition.

KEY SUPPLIERS



[More companies ▶](#)

FEATURED COMPANIES




Theory

Λ_c^+ two body decays

Non-leptonic two-body weak decays of $\Lambda_c(2286)$

Physics Letters B 776 (2018) 265–269

C.Q. Geng^{a,b,*}, Y.K. Hsiao^{a,b}, Yu-Heng Lin^b, Liang-Liang Liu^a



[11] H.Y. Cheng, B. Tseng, Phys. Rev. D 48 (1993) 4188.

[23] K.K. Sharma, R.C. Verma, Phys. Rev. D 55 (1997) 7067.

[24] K.K. Sharma, R.C. Verma, Eur. Phys. J. C 7 (1999) 217.

Λ_c^+ three body decays






FIG. 1. The dominant diagram for the $\Lambda_c^+ \rightarrow \pi^+ M B$ decay. The solid lines and the wiggly line show the quarks and the W boson, respectively.




Weak decay of Λ_c^+ for the study of $\Lambda(1405)$ and $\Lambda(1670)$

$$\begin{aligned}
 |MB\rangle &= \frac{1}{\sqrt{2}} |s(\bar{u}u + \bar{d}d + \bar{s}s)(ud - du)\rangle \\
 &= \frac{1}{\sqrt{2}} \sum_{i=1}^3 |P_{3i} q_i (ud - du)\rangle,
 \end{aligned}
 \quad
 q \equiv \begin{pmatrix} u \\ d \\ s \end{pmatrix}, \quad
 P \equiv q\bar{q} = \begin{pmatrix} u\bar{u} & u\bar{d} & u\bar{s} \\ d\bar{u} & d\bar{d} & d\bar{s} \\ s\bar{u} & s\bar{d} & s\bar{s} \end{pmatrix}.$$



$$|MB\rangle = |K^- p\rangle + |\bar{K}^0 n\rangle - \frac{\sqrt{2}}{3} |\eta \Lambda\rangle.$$




$$\mathcal{M}_j = V_P \left[h_j + \sum_i h_i G_i(M_{\text{inv}}) t_{ij}(M_{\text{inv}}) \right],$$

$$h_{\pi^0 \Sigma^0} = h_{\pi^- \Sigma^+} = h_{\pi^+ \Sigma^-} = h_{\pi^0 \Lambda} = 0,$$

$$h_{K^- p} = h_{\bar{K}^0 n} = 1, \quad h_{\eta \Lambda} = -\frac{\sqrt{2}}{3},$$

$$\frac{d\Gamma_j}{dM_{\text{inv}}} = \frac{1}{(2\pi)^3} \frac{p_{\pi^+} \tilde{p}_j M_{\Lambda_c^+} M_j}{M_{\Lambda_c^+}^2} |\mathcal{M}_j|^2,$$

$$h_{\eta \Sigma^0} = h_{K^+ \Xi^-} = h_{K^0 \Xi^0} = 0,$$



A possible Σ^* state with spin-parity $J^P = \frac{1}{2}^-$

	(Y, I)	I_3	Flavor wave functions	Masses (MeV)
p_8	$(1, \frac{1}{2})$	$\frac{1}{2}$	$[su][ud] - \bar{s}$	1460
n_8		$-\frac{1}{2}$	$[ds][ud] - \bar{s}$	1460
Σ_8^+	$(0, 1)$	1	$[su][ud] - \bar{d}$	1360
Σ_8^0		0	$\frac{1}{\sqrt{2}}([su][ud] - \bar{u} + [ds][ud] - \bar{d})$	1360
Σ_8^-		-1	$[ds][ud] - \bar{u}$	1360
Λ_8	$(0, 0)$	0	$\frac{[ud][su] - \bar{u} + [ds][ud] - \bar{d} - 2[su][ds] - \bar{s}}{\sqrt{6}}$	1533
Ξ_8^0	$(-1, \frac{1}{2})$	$\frac{1}{2}$	$[ds][su] - \bar{d}$	1520
Ξ_8^-		$-\frac{1}{2}$	$[ds][su] - \bar{u}$	1520
Λ_1	$(0, 0)$	0	$\frac{[ud][su] - \bar{u} + [ds][ud] - \bar{d} + [su][ds] - \bar{s}}{\sqrt{3}}$	1447

TABLE II: Flavor wave functions and masses of the $\frac{1}{2}^-$ pentaquark octet and singlet.

Ao Zhang, Y. R. Liu, P.Z. Huang, W.Z. Deng, X.L. chen and S.L. Zhu,
High Energy Phys. Nucl. Phys. 29, 250 (2005).

Other Model Predictions

Chiral dynamics in the presence of bound states:
kaon–nucleon interactions revisited

Physics Letters B 500 (2001) 263–272

J.A. Oller, Ulf-G. Meißner

z_R	1401 + 40i	
	g_i	$ g_i $
(I = 1)		
$\pi \Lambda$	0.60 + 0.47 i	0.76
$\pi \Sigma$	1.27 + 0.71 i	1.5
$\bar{K} N$	−1.24 − 0.73 i	1.4
$\eta \Sigma$	0.56 + 0.41 i	0.69
$K \Xi$	0.12 + 0.05 i	0.13

Chiral dynamics of the two $\Lambda(1405)$ states

D. Jido ^{a,c}, J.A. Oller ^{b,*}, E. Oset ^c, A. Ramos ^d, U.-G. Meißner ^e

Nuclear Physics A 725 (2003) 181–200

Cusp or “resonance” around the $\bar{K} N$ threshold

J.A. Oller, Eur. Phys. J. A 28, 63–82 (2006). Zhi-Hui Guo and J. A. Oller, PRC 87, 035202 (2013) .


L. Roca and E. Oset, PRC 88, 055206 (2013).

K. P. Khemchandani, A. Martinez Torres, and J. A. Oller, Phys.Rev. C100, 015208 (2019).

in the fit. Two type of fits are found as a result. In both cases, the properties of $\Lambda(1405)$ are well reproduced. In addition to this, a Σ state is also found with mass around 1400 MeV. Cross sections,

Evidence for a new Σ^* resonance with $J^P = 1/2^-$ in the old data of the $K^- p \rightarrow \Lambda \pi^+ \pi^-$ reaction

Jia-Jun Wu,¹ S. Dulat,^{2,3} and B. S. Zou^{1,3}



	$M_{\Sigma^*(3/2)}$	$\Gamma_{\Sigma^*(3/2)}$
Fit1	1385.3 ± 0.7	46.9 ± 2.5
Fit2	$1386.1^{+1.1}_{-0.9}$	$34.9^{+5.1}_{-4.9}$
	$M_{\Sigma^*(1/2)}$	$\Gamma_{\Sigma^*(1/2)}$
	$1381.3^{+4.9}_{-8.3}$	$118.6^{+55.2}_{-35.1}$

PHYSICAL REVIEW C **81**, 045210 (2010)

Possible evidence for the Σ^* resonance with $J^P = 1/2^-$ around 1380 MeV

PHYSICAL REVIEW C **81**, 055203 (2010)

Jia-Jun Wu,¹ S. Dulat,^{2,3} and B. S. Zou^{1,3}




Possible $\Sigma(\frac{1}{2}^-)$ under the $\Sigma^*(1385)$ peak in $K \Sigma^*$ photoproduction

Puze Gao, Jia-Jun Wu, and B. S. Zou

Yun-Hua Chen and B. S. Zou, PRC **88**, 024304 (2013).

Ju-Jun Xie, Jia-Jun Wu, and Bing-Song Zou, PRC **90**, 055204 (2014).



$\Sigma(1380)$ in $\Lambda_c^+ \rightarrow \eta \pi^+ \Lambda$ decay



$$\mathcal{M}_1 = i g_{\pi \Lambda \Sigma_1^*} \bar{u}(p_3) G^{\Sigma_1^*}(q) (A_1 + B_1 \gamma_5) u(p),$$

$$G^{\Sigma_1^*}(q) = i \frac{\not{q} + M_{\Sigma_1^*}}{q^2 - M_{\Sigma_1^*}^2 + i M_{\Sigma_1^*} \Gamma_{\Sigma_1^*}},$$

$\Sigma(1385)$ in $\Lambda_c^+ \rightarrow \eta \pi^+ \Lambda$ decay



$$\mathcal{M}_2 = \frac{i g_{\pi \Lambda \Sigma_2^*}}{m_\eta m_\pi} \bar{u}(p_3) p_2^\mu G_{\mu\nu}^{\Sigma_2^*}(q) p_1^\nu (A_2 + B_2 \gamma_5) u(p),$$

$$G_{\mu\nu}^{\Sigma_2^*}(q) = i \frac{\not{q} + M_{\Sigma_2^*}}{q^2 - M_{\Sigma_2^*}^2 + i M_{\Sigma_2^*} \Gamma_{\Sigma_2^*}} P_{\mu\nu},$$

with

$$P^{\mu\nu} = -g^{\mu\nu} + \frac{1}{3} \gamma^\mu \gamma^\nu + \frac{2q^\mu q^\nu}{3M_{\Sigma_2^*}^2} + \frac{\gamma^\mu q^\nu - \gamma^\nu q^\mu}{3M_{\Sigma_2^*}},$$

Invariant mass, decay angle and energy distributions

$$\frac{d\Gamma}{dM_{\pi+\Lambda}} = \frac{m_{\Lambda}}{32\pi^3 M_{\Lambda_c^+}} \int \sum |\mathcal{M}|^2 |\vec{p}_1| |\vec{p}^*| d\cos\theta^*$$

$$\frac{d\Gamma}{d\cos\theta^*} = \frac{m_{\Lambda}}{32\pi^3 M_{\Lambda_c^+}} \int \sum |\mathcal{M}|^2 |\vec{p}_1| |\vec{p}^*| dM_{\pi+\Lambda}.$$

$$\frac{d\Gamma}{dE_{\pi+}} = \frac{m_{\Lambda}}{32\pi^3} \int \sum |\mathcal{M}|^2 dE_{\Lambda};$$

Invariant mass distributions




FIG. 5: Invariant mass distributions $d\Gamma/dM_{\pi+\Lambda}$ as a function of $M_{\pi+\Lambda}$.

Decay angle and energy distributions




FIG. 6: Angle distributions $d\Gamma/d\cos\theta^*$ in the c.m. frame of $\pi^+\Lambda$ system as a function of $\cos\theta^*$.






FIG. 7: Energy distributions $d\Gamma/dE_{\pi^+}$ in the rest frame of Λ_c^+ as a function of E_{π^+} .

Measurement of the absolute branching fractions of $\Lambda_c^+ \rightarrow \Lambda \eta \pi^+$ and $\Sigma(1385)^+ \eta$ (BESIII Collaboration)

PRD 99, 032010 (2019).



Observation of $\Lambda_c^+ \rightarrow \Lambda a_0(980)^+$ and Evidence for $\Sigma(1380)^+$ in $\Lambda_c^+ \rightarrow \Lambda \pi^+ \eta$

M. Ablikim *et al.*^{*}
(BESIII Collaboration)

 (Received 17 July 2024; revised 19 November 2024; accepted 4 December 2024; published 17 January 2025)





FIG. 2. Projections of the fit results in the $M_{\pi^+\eta}$, $M_{\Lambda\pi^+}$, and $M_{\Lambda\eta}$ spectra. Points with error bars are sWeighted data at all energy points. The curves in different colors are different components.

$N^*(1535)$: strangeness component

Couples strongly to strangeness channel

$$uud (L=1) 1/2^- \sim N^*(1535) \sim [ud][us] \bar{s}$$



	Mode	Fraction (Γ_i/Γ)
Γ_1	$N\pi$	32-52 %
Γ_2	$N\eta$	30-55 %

Larger $[ud][us] \bar{s}$ component in $N^*(1535)$ makes it coupling strong to $N\eta$ & $K\Lambda$.

$$J/\psi \rightarrow \bar{p}N^* \rightarrow \bar{p} (K\Lambda) / \bar{p} (p\eta) \rightarrow \text{large } g_{N^*K\Lambda}$$

Liu&Zou, PRL96 (2006) 042002; Geng,Oset,Zou&Doring, PRC79 (2009) 025203

$$\gamma p \rightarrow p\eta' \text{ \& } pp \rightarrow pp\eta' \rightarrow \text{large } g_{N^*N\eta'}$$

M.Dugger et al., PRL96 (2006) 062001; Cao&Lee, PRC78(2008) 035207

$$\pi^- p \rightarrow n\phi \text{ \& } pp \rightarrow pp\phi \text{ \& } pn \rightarrow d\phi \rightarrow \text{large } g_{N^*N\phi}$$

Xie, Zou & Chiang, PRC77(2008)015206; Cao, Xie, Zou & Xu, PRC80(2009)025203

$N^*(1535)$: dynamically generated state

- Pole position

PDG 2024  $Z_R = [(1500 \sim 1520) - i(40 \sim 65)] \text{ MeV}$

PHYSICAL REVIEW C, VOLUME 65, 035204

Chiral unitary approach to S -wave meson baryon scattering in the strangeness $S=0$ sector

T. Inoue,^{*} E. Oset, and M. J. Vicente Vacas

*Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC, Institutos de Investigación de Paterna,
Apartado Correos 22085, E-46071 Valencia, Spain*

(Received 31 October 2001; published 14 February 2002)


Chiral dynamics of the $S_{11}(1535)$ and $S_{11}(1650)$ resonances revisited

Peter C. Bruns^a, Maxim Mai^{b,*}, Ulf-G. Meißner^{b,c}

Physics Letters B 697 (2011) 254–259

- Breit-Wigner parameterization

$$(M_R, \Gamma_R) = (1525 \sim 1545, 125 \sim 175) \text{ MeV} = (\simeq 1535, \simeq 150) \text{ MeV}$$




PDG 2024

The $N^*(1535)$ as a dynamically generated state



$$|uud\rangle \rightarrow \frac{1}{\sqrt{2}}|u(ud - du)\rangle$$

$$\bar{u}u + \bar{d}d + \bar{s}s$$




$$|MB\rangle = \frac{\sqrt{3}}{3}|\eta p\rangle + \frac{\sqrt{2}}{2}|\pi^0 p\rangle + |\pi^+ n\rangle - \frac{\sqrt{6}}{3}|K^+ \Lambda\rangle,$$

$$T^{MB} = V_P \left(\frac{\sqrt{3}}{3} + \frac{\sqrt{3}}{3} G_{\eta p}(M_{\eta p}) t_{\eta p \rightarrow \eta p}(M_{\eta p}) \right. \\ \left. + \frac{\sqrt{2}}{2} G_{\pi^0 p}(M_{\eta p}) t_{\pi^0 p \rightarrow \eta p}(M_{\eta p}) \right. \\ \left. + G_{\pi^+ n}(M_{\eta p}) t_{\pi^+ n \rightarrow \eta p}(M_{\eta p}) \right. \\ \left. - \frac{\sqrt{6}}{3} G_{K^+ \Lambda}(M_{\eta p}) t_{K^+ \Lambda \rightarrow \eta p}(M_{\eta p}) \right),$$



Effective Lagrangian approach and the $N^*(1535)$ resonance as a Breit-Wigner resonance




$$T^{N^*} = ig_{N^*N\eta} \bar{u}(p_3, s_p) G_{N^*}(q) (A + B\gamma_5) u(p, s_{\Lambda_c^+}),$$

$$G_{N^*}(q) = i \frac{\not{q} + M_{N^*}}{q^2 - M_{N^*}^2 + iM_{N^*}\Gamma_{N^*}(q^2)},$$


$$\Gamma_{N^*}(q^2) = \Gamma_{N^* \rightarrow \pi N}(q^2) + \Gamma_{N^* \rightarrow \eta N}(q^2) + \Gamma_0,$$

$$\Gamma_0 = 19.5 \text{ MeV} \quad \text{for} \quad \Gamma_{N^*}(\sqrt{q^2} = 1535 \text{ MeV}) = 150 \text{ MeV}.$$



Ju-Jun Xie and Li-Sheng Geng, PRD 96, 054009 (2017).

Invariant ηp mass distributions



Model I : $T = T^{MB}$

Model II : $T = T^{N^*}$, $M_{N^*} = 1535$ MeV, $\Gamma_{N^*} = \Gamma_{N^*}(q^2)$

Other contributions

$$N^*(1650) \rightarrow \eta p$$

Σ^* resonances

$$\rightarrow \bar{K}^0 p$$




FIG. 7. Dalitz plot for $M_{\eta p}^2$ and $M_{\bar{K}^0 p}^2$ in the $\Lambda_c^+ \rightarrow \bar{K}^0 \eta p$ decay. The $N^*(1535)$ energy is shown by the vertical dotted line, and the horizontal band represents the masses of Σ^* states from 1600 to 1800 MeV.

Production of $N^*(1535)$ and $N^*(1650)$ in $\Lambda_c \rightarrow \bar{K}^0 \eta p$ (πN) decay



Measurement of branching fractions of $\Lambda_c^+ \rightarrow pK_S^0 K_S^0$ and $\Lambda_c^+ \rightarrow pK_S^0 \eta$ at Belle


FIG. 4. For $\Lambda_c^+ \rightarrow pK_S^0 \eta$, the Dalitz plot after background subtraction and efficiency correction bin-by-bin and its projections superimposing with signal MC produced by phase space mode (blue histograms). A significant structure of $N^*(1535)$ near the $p\eta$ threshold is found.



 Theoretical study of $N(1535)$ and $\Sigma^*(1/2^-)$ in the Cabibbo-favored process $\Lambda_c^+ \rightarrow p\bar{K}^0 \eta$

Ying Li,¹ Si-Wei Liu^{2,3}, En Wang^{4,*}, De-Min Li,^{1,†} Li-Sheng Geng,^{5,6,7,8,‡} and Ju-Jun Xie^{2,3,8,§}



Possible ϕp state in $\Lambda_c^+ \rightarrow \pi^0 p \phi$ decay

R. Lebed, PRD92(2015)114030



$\Sigma^+ \rightarrow p \pi^0$ vetoed

From Cheng-Ping Shen

- **No significant P_s signal**
- Best fit yields a peak at $M = (2025 \pm 5) \text{ MeV}/c^2$ and $\Gamma = (22 \pm 12) \text{ MeV}$


[PRD96, 051102\(R\) \(2017\); 915fb⁻¹](#)

Number of candidate $\Lambda_c \rightarrow P_s \pi^0 \rightarrow \phi p \pi^0$ events: 77.6 ± 28.1


$B(\Lambda_c \rightarrow P_s \pi^0) \times B(P_s \rightarrow \phi p) < 8.3 \times 10^{-5} @ 90\% \text{ C.L.}$

Possible ϕp state in $\Lambda_c^+ \rightarrow \pi^0 p \phi$ decay





$N^*(2120)$ in $\gamma p \rightarrow \phi p$ and $\gamma p \rightarrow K^+ \Lambda(1520)$ reactions



(a)



(b)



Alvin Kiswandhi, Ju-Jun Xie, Shin Nan Yang, PLB, 691, 214 (2010).
Ju-Jun Xie, and Juan Nieves, Phys. Rev. C 82, 045205 (2010).





Fig. 1. Triangle diagrams for the $\Lambda_c^+ \rightarrow \pi^0 p \phi$ decay. (A): Σ^+ -exchange. (B): Λ -exchange. The definitions of the kinematical variables (P, q, k) are also shown.

$$\begin{aligned}
 t = & \frac{g_{\Lambda_c \Sigma^* K^*} g}{m_\pi} \vec{\epsilon}_\phi \cdot \vec{k} \sum_{i=\Sigma, \Lambda} c_i \int \frac{d^4 q}{(2\pi)^4} \\
 & \times \frac{i 2m_{\Sigma^*}}{(P-q)^2 - m_{\Sigma^*}^2 + i m_{\Sigma^*} \Gamma_{\Sigma^*}} \frac{i}{q^2 - m_{K^*}^2 + i m_{K^*} \Gamma_{K^*}} \\
 & \times \frac{i 2m_i}{(P-q-k)^2 - m_i^2 + i\epsilon},
 \end{aligned} \tag{4}$$

where we have defined $\mathcal{C}_\Sigma = \frac{\sqrt{6}}{3} t_{K^{*0} \Sigma^+ \rightarrow \phi p}$ and $\mathcal{C}_\Lambda = -t_{K^{*+} \Lambda \rightarrow \phi p}$,




Fig. 3. The squared norm of the T -matrix elements for $K^{*+} \Lambda \rightarrow \phi p$ and $K^{*0} \Sigma^+ \rightarrow \phi p$ as a function of the meson-baryon invariant mass E_{cm} in the model of Ref. [72].




Fig. 2. Invariant mass distribution of the $\Lambda_c^+ \rightarrow \pi^0 p \phi$ decay. The experimental data are taken from Ref. [47].

[47] B. Pal, et al., Belle Collaboration, arXiv:1707.00089 [hep-ex].

$$t_I = t,$$

$$t_{II} = t, \text{ but with } t_{K^{*+} \Lambda \rightarrow \phi p} = \frac{\sqrt{6}}{2} t_{K^{*0} \Sigma^+ \rightarrow \phi p}$$

$$= c_1 \frac{\sqrt{6}}{2} \frac{E_{K^*} + E_\phi}{4F_\pi^2},$$

$$t_{III} = c_2,$$

Summary

The $\Lambda_c^+ \rightarrow \eta\pi^+\Lambda$ decay can be used to study the Σ^ and Λ^* resonances*

The $\Lambda_c^+ \rightarrow \bar{K}^0\eta p$ decay can be used to study the $N^(1535)$ resonance*

Possible ϕp state, P_s , in the $\Lambda_c^+ \rightarrow \pi^0\phi p$ decay

TS produces a bump at around 2.02 GeV

P_s , if exists, could distort the line shape, but difficult to be distinguished from TS in this process

Hence, we need more efforts, both on theoretical and experimental sides.

Thank you very much for your attention! ₃₀