
第十一届中国LHC物理会议(CLHCP2025)

Selected highlights of correlations and flow at LHC-ALICE


Qiye Shou 寿齐烨 Fudan University 复旦大学 2025.11.01 新乡







Evolution of relativistic heavy-ion collisions



Quark-gluon plasma (QGP)

Strongly interacting matter, in the extremely hot and dense environment microseconds after the Big Bang

Collective motion (Collectivity)

One of the most important features and probes of the QGP


LHC Run 3 data-taking (2023–2026): high statistics e.g. 13 nb⁻¹ Pb−Pb collisions, ≈100 times the MB statistics of Run 2


Azimuthal correlations and anisotropic flow

Initial eccentricity + QGP evolution → Final momentum anisotropy


$$E\frac{\mathrm{d}^3 N}{\mathrm{d}^3 \mathbf{p}} = \frac{1}{2\pi} \frac{\mathrm{d}^2 N}{p_{\mathrm{t}} \mathrm{d} p_{\mathrm{t}} \mathrm{d} y} \left(1 + 2 \sum_{n=1}^{\infty} v_n \cos[n(\varphi - \Psi_{\mathrm{RP}})] \right)$$


$$v_n(p_t, y) = \langle \cos[n(\varphi - \Psi_{RP})] \rangle$$





Anisotropic flow in OO and Ne-Ne collisions








Observation of hydrodynamic response to initial geometry in OO & Ne-Ne Unique inputs to collectivity in small systems and nuclear structure

Anisotropic flow in OO and Ne-Ne collisions

arXiv:2510.02580






Observation of hydrodynamic response to initial geometry in OO & Ne-Ne Unique inputs to collectivity in small systems and nuclear structure

Anisotropic flow in OO and Ne-Ne collisions


arXiv:2509.05171





Observation of hydrodynamic response to initial geometry in OO & Ne-Ne Unique inputs to collectivity in small systems and nuclear structure


Collectivity in p-Pb and pp collisions



- > Do we expect the collectivity in small collision systems?
- ➤ Where is the "smallest" (p-A, pp, ee...) and "dilutest" (lower multiplicity) limit of collectivity onset?

Collectivity in p-Pb and pp collisions


PRL 123, 142301 (2019)



The magnitudes of v_n in pp and p-Pb are similar as in Pb-Pb at low multiplicities


Elliptic flow of identified particles in small systems


arXiv:2411.09323 Nature Commu. accepted



Mass ordering and baryon-meson grouping for all Pb-Pb, p-Pb and pp collisions, indicating the partonic collectivity


Elliptic flow of identified particles in small systems





Evidence of hadronization via coalescence of flowing quarks in small systems

Elliptic flow of identified particles in small systems



Where is the "smallest" and "dilutest" limit of collectivity onset?

Parameterizing the shape of nuclei

nucleon density described by Woods-Saxon profile


$$\rho(r,\theta,\phi) = \frac{\rho_0}{1 + e^{[r-R(\theta,\phi)]/a_0}},$$

$$R(\theta,\phi) = R_0(1+\beta_2[\cos\gamma Y_{2,0} + \sin\gamma Y_{2,2}] + \beta_3 \sum_{m=-3}^{3} \alpha_{3,m} Y_{3,m} + \beta_4 \sum_{m=-4}^{4} \alpha_{4,m} Y_{4,m})$$

 β_2 : overall deformation parameter


 a_0 : diffuseness parameter

y: triaxiality parameter




RHIC: Au+Au (relatively)


LHC: Pb+Pb (double magic nucleus)




RHIC: U+U (Ru, Zr)


LHC: Xe+Xe








Imagining the nuclear structure



Measurement of Xe at LHC-ALICE


arXiv:2409.04343



Measurement of Xe at LHC-ALICE


$$\chi^2/N_{\text{dof}} = \frac{1}{N_{\text{dof}}} \sum \frac{(y_i - f_i)^2}{\sigma_i^2}$$


arXiv:2409.04343v1




Data favor $a \approx 0.492$ and $\beta_2 \approx 0.207$

Testing γ-soft of Xe at LHC-ALICE









6-particle cumulant needed

Testing γ-soft of Xe at LHC-ALICE






Comprehensive Bayesian analysis to simultaneously pin down all parameters of Xe The extracted γ value with a narrow σ indicates weak/no γ -soft effect.


→ Xe has a rigid, triaxial deforamtion


Collective effects under strong magnetic and vortical fields




B $\approx 10^{18} - 10^{19}$ Gauss $\omega \approx 10^{21}$ s⁻¹


- Anomalous chiral effects Search for the CP violations in the strong interactions and disentangle the background mechanisms
- Polarization phenomena Reveal how global/local polarization observables respond to the QGP collectivity






Charge dependent azimuthal correlations





Significant separations are observed for K and p $\Delta\gamma$ follow p-p > p- Λ > K-K > h-h indicating various global/local charge conservations

Hyperon local polarization at ALICE




Second and third order local polarization are clearly observed with Run 3 data

Hyperon local polarization at CMS


PRL 135 (2025) 132301






First observation of hyperon local polarization in small collisions Post challenges to theoretical framework of spin in heavy ion

Momentum correlations (Femtoscopy)



π-nuclei femto correlation



Summary

- > CF is one of the most essential branches in the QGP physics
- Highlight results across azimuthal correlations, flow, femtoscopy are presented
- Chinese members have made and will keep on making big contributions

Thank you for your attention!