

First measurement of Λ_c^+ elliptic flow in 30-50% Pb-Pb collisions at $\sqrt{s_{NN}} = 5.36$ TeV with ALICE

Thursday, 30 October 2025 17:20 (20 minutes)

Heavy quarks (charm and beauty) are useful probes for investigating the properties of the quark-gluon plasma (QGP) generated in ultra-relativistic heavy-ion collisions. Their participation in the collective motion of the medium can be assessed by measuring the charm-hadron elliptic-flow coefficient, originating from the initial-state spatial asymmetry in non-central heavy-ion collisions. These measurements provide fundamental inputs to constrain theoretical models describing the charm-quark transport in the QGP, as well as its possible thermalization in the medium. In addition, the comparison between meson and baryon can provide further insights into medium-induced phenomena, such as the radial flow and the charm-quark hadronization via coalescence. In this contribution, the first measurements of charm baryon Λ_c^+ elliptic flow in 30-50% centrality intervals of Pb-Pb collisions at $\sqrt{s_{NN}} = 5.36$ TeV collected by the ALICE experiment during the LHC Run 3 are shown. The measurements are compared to model predictions that incorporate various implementations of heavy-quark interaction and hadronization with the QGP constituents.

Primary authors: XUE, Xufei (China University of Geosciences (CN), WuHan); PENG, Xinye (China University of Geosciences)

Presenter: XUE, Xufei (China University of Geosciences (CN), WuHan)

Session Classification: Parallel 3

Track Classification: ALICE