

Recent results at Belle II

李龙科（湖南师范大学）




第七届粒子物理天问论坛@武汉


2025年9月20日于中国地质大学



Outline





- Luminosity
- Recent **B** results
- Recent **charm** results
- Recent **$q\bar{q}$** and **exotic**
- Recent **tau** and **dark** sector
- Summary



Luminosity




In Dec. 2024, SuperKEKB achieved WR: $5.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$




“Belle II has been designed to make precise measurements of weak interaction parameters, study exotic hadrons, and search for new phenomena beyond the Standard Model of particle physics.” -- from Belle II website homepage [\[link\]](#).

Physics & Publications



Belle II Journal Publications per Year



➤ 目前文章数 (含arXiv等): >100

➤ PRL占比: 24% < Belle均值33%





奥利给!

The Belle II Physics Book [PTEP 2019 (2019) 12, 123C01]

Outline



- Luminosity
- Recent B results
 - B CPV
 - B rare/forbidden
 - B (semi-)leptonic
- Recent charm results
- Recent $q\bar{q}$ and exotic
- Recent tau and dark sector
- Summary



$$B^0 \rightarrow J/\psi K_S^0$$

Belle+BABAR final result:

$$S = \sin 2\phi_1 = 0.677 \pm 0.020$$

LHCb ([PRL 132, 021801 \(2024\)](#)):

$$S = 0.722 \pm 0.014 \pm 0.007$$

its precision is dominated by LHCb.

$\sin 2\phi_1: B^0 \rightarrow J/\psi \pi^0$

$$\phi_1 = \arg(-V_{cd}V_{cb}^*/V_{td}V_{tb}^*)$$

PRD 111 (2025) 012011






$S = -\sin 2\phi_1$, $C = 0$ if there is only tree amplitude


Tree is color and CKM suppressed

→ can be used to understand the loop contribution in $B^0 \rightarrow J/\psi K_S^0$

- Improved sensitivity by the better π^0 selection and GflaT
- $\Delta E - m(l\bar{l})$ fit to extract signal



$$(\Delta E = E_B - E_{\text{beam}})$$



$$S = -0.88 \pm 0.17 \pm 0.03$$

$$C = 0.13 \pm 0.12 \pm 0.03$$

$$\mathcal{B} = (2.02 \pm 0.12 \pm 0.10) \times 10^{-5}$$

Most precise,

- first observation of non-zero S parameter (mixing-induced CPV) in this mode



$$\phi_2: B^0 \rightarrow \rho^+ \rho^- \quad \phi_2 = \arg(-V_{td}V_{tb}^*/V_{ud}V_{ub}^*)$$

PRD 111 (2025) 092001



- $B^0 \rightarrow \rho^+ \rho^-$: much smaller loop contribution $\rightarrow$ dominates ϕ_2 precision
- two π^0 's reconstruction needed $\rightarrow$ a channel suitable for Belle II

➤ Extract ϕ_2 using this new result



$B \rightarrow \rho\rho$ world average

$$\phi_2 = (91.5^{+4.5}_{-5.4})^\circ$$

$B \rightarrow \rho\rho$ world average

+ Belle II $\rho^+ \rho^-$ results

$$\rightarrow \phi_2 = (92.6^{+4.5}_{-4.8})^\circ$$

6% ↑

	$\mathcal{B}(10^{-6})$	f_L	S	$\mathcal{C}$
Belle II	$29.0^{+2.3}_{-2.2} {}^{+3.1}_{-3.0}$	$0.921^{+0.024}_{-0.025} {}^{+0.017}_{-0.015}$	$-0.26 \pm 0.19 \pm 0.08$	$-0.02 \pm 0.12 {}^{+0.06}_{-0.05}$
Belle	$28.3 \pm 1.5 \pm 1.5$	$0.988 \pm 0.012 \pm 0.006$	$-0.13 \pm 0.15 \pm 0.05$	$0.00 \pm 0.10 \pm 0.06$
BABAR	$25.5 \pm 2.1 {}^{+3.6}_{-3.9}$	$0.992 \pm 0.024 {}^{+0.026}_{-0.013}$	$-0.17 \pm 0.20 {}^{+0.05}_{-0.06}$	$0.01 \pm 0.15 \pm 0.06$


$B \rightarrow K \nu \bar{\nu}$: first evidence

PRD 109 (2024) 112006




Penguin:

FCNC



Box:



High accuracy in the SM [PRD 107, 014511 (2023)]

$$\mathcal{B}(B \rightarrow K \nu \bar{\nu}) = (5.6 \pm 0.4) \times 10^{-6}$$


Extensions beyond SM may lead to significant rate increase.

$$\mathcal{B}(B \rightarrow K \nu \bar{\nu}) = (2.3 \pm 0.7) \times 10^{-5} (3.5\sigma)$$

Inclusive tag

$\eta(\text{BDT}_2)$

0.92 0.94 0.96 0.98 1.0




3.5 σ

Belle II preliminary
 $\int \mathcal{L} dt = (362 + 42) \text{ fb}^{-1}$

$B^+ \rightarrow K^+ \nu \bar{\nu}$
 $B^0 \bar{B}^0$
 $B^+ B^-$
Continuum
Data

Hadronic tag


Candidates/0.1



1.1 σ

Belle II preliminary
 $\int \mathcal{L} dt = 362 \text{ fb}^{-1}$

$B^+ \rightarrow K^+ \nu \bar{\nu}$
 $B \bar{B}$
 $c \bar{c}$
 $u \bar{u}, d \bar{d}, s \bar{s}$
Data




$B \rightarrow X_s \nu \bar{\nu}$ Search

Preliminary result




- Complimentary measurement to the exclusive $B \rightarrow K(*) \nu \bar{\nu}$ searches.



$M_{X_s} [\text{GeV}/c^2]$	ϵ	N_{sig}	$\mathcal{B} [10^{-5}]$		
			Central value	UL_{obs}	UL_{exp}
*[0, 0.6]	0.26%	10^{+18+18}_{-17-16}	$0.5^{+0.9+0.9}_{-0.8-0.8}$	2.5	2.4
[0.6, 1.0]	0.12%	37^{+27+31}_{-25-26}	$3.8^{+2.8+3.2}_{-2.6-2.7}$	10.1	7.3
[1.0, m_B]	0.06%	33^{+44+63}_{-42-53}	$7.3^{+9.6+13.7}_{-9.3-11.5}$	34.4	27.4

*Compatible with the hadronically-tagged Belle II $B^+ \rightarrow K^+ \nu \bar{\nu}$ result

➤ Main syst. error: MC statistics, background normalization



Full M_{X_s} range result:

$$\mathcal{B}(B \rightarrow X_s \nu \bar{\nu}) = [11.6^{+8.9}_{-8.6}(\text{stat})^{+13.5}_{-11.4}(\text{syst})] \times 10^{-5}$$

$$\mathcal{B}(B \rightarrow X_s \nu \bar{\nu}) < 3.6 \times 10^{-4} \text{ (90 \% CL)}$$

Most stringent upper limit on the inclusive rate


Outline



- Luminosity
- Recent B results
- Recent charm results
- Recent $q\bar{q}$ and exotic
- Recent tau and dark sector
- Summary

charm CPV
charmed baryon
charm spectrum

Belle (II): charm super-factory



PS: no Λ_b^0 sample

New charm wave: Charm CPV



- Charm CPV effect is very small (10^{-3} level or smaller); a sensitive probe for New Physics.
- 2019, LHCb: $\Delta A_{CP}(D^0 \rightarrow K^+ K^-, \pi^+ \pi^-) = (-15.4 \pm 2.9) \times 10^{-4} (5.3\sigma)^{[1]}$;
2023, LHCb: $A_{CP}^{\text{dir}}(D^0 \rightarrow \pi^+ \pi^-) = (2.32 \pm 0.61) \times 10^{-3} (3.8\sigma)^{[2]}$.
→ to understand this CPV, more results and more precise measurements are desired.
- CPV <2025: observed in all open-flavor meson sector, not yet in the baryon sector.
2025, LHCb: $A_{CP}(\Lambda_b^0 \rightarrow p K^- \pi^+ \pi^-) = (2.45 \pm 0.46 \pm 0.10)\% (5.2\sigma)^{[3]}$.
→ charmed baryon CPV, the last piece for heavy-flavor hadron CPV, yet to be observed.

^[1]LHCb, [PRL 122, 211803 \(2019\)](#)

^[2]LHCb, [PRL 131, 091802 \(2023\)](#)

^[3]LHCb, [Nature 643 \(2025\) 8074](#)




New charm wave.

- $A_{CP}(D^0 \rightarrow K_S^0 K_S^0)$: [PRD 111, 012015 \(2025\)](#) + [PRD 112, 012017 \(2025\)](#)
- $A_{CP}^X(D^+ \rightarrow K_S^0 K^- \pi^+ \pi^+)$: [JHEP 04 \(2025\) 036](#)
- $A_{CP}(D^0 \rightarrow \pi^0 \pi^0)$: [PRD 112, 012006 \(2025\)](#)
- $A_{CP}(D^+ \rightarrow \pi^+ \pi^0)$: [PRD 112, L031101 \(2025\)](#)
- $A_{CP}(D^0 \rightarrow \pi^+ \pi^- \pi^0)$: preliminary result
- $A_{CP}(\Lambda_c^+ \rightarrow p h^+ h^-), A_{CP}(\Xi_c^+ \rightarrow \Sigma^+ h^+ h^-)$: preliminary result


2/3/4-body decay
meson/baryon

Charm CPV: $A_{CP}(D^0 \rightarrow K_S^0 K_S^0)$

PRD 111, 012015 (2025)



(one golden channel) $A_{CP}(D^0 \rightarrow K_S^0 K_S^0)$ may be **enhanced to 1% level with the SM**, by the interference of $c \rightarrow us\bar{s}$ and $c \rightarrow ds\bar{s}$ amplitudes. [PRD 99, 113001 (2019), PRD 92, 054036 (2015)]



- Belle: $A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (-1.1 \pm 1.6 \pm 0.1)\%$
- Belle II: $A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (-2.2 \pm 2.3 \pm 0.1)\%$
- Combined $A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (-1.4 \pm 1.3 \pm 0.1)\%$: comparable to the world-best result: $(-3.1 \pm 1.3)\%$ [PRD 104 \(2021\) L031102](#)
- Belle(II)+LHCb average: $(-2.3 \pm 0.9)\%$ vs. CMS: $(6.2 \pm 3.1)\%$: 2.6σ diff. $\Rightarrow$ more precise result desired. [EPJC 84 \(2024\) 1264](#)

Charm CPV: $A_{CP}(D^0 \rightarrow K_S^0 K_S^0)$


PRD 112, 012017 (2025)



➤ An independent sample:

using opposite-side flavor tagging for $e^+e^- \rightarrow c\bar{c}$

[(Belle II) PRD 107, 112010 (2023)]



- such new tagged sample at Belle+Belle II:

$$A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (+1.3 \pm 2.0 \pm 0.2)\%$$

- combining it with that from D^{*+} -tagged sample:


$$A_{CP}(D^0 \rightarrow K_S^0 K_S^0) = (-0.6 \pm 1.1 \pm 0.1)\%$$

most precise

➤ LHCb 2025: $A_{CP} = (1.86 \pm 1.04 \pm 0.38)\%$

➤ Now W.A. $A_{CP} = (-0.17 \pm 0.62 \pm 0.18)\%$

For details, see [Xinchen's talk](#) at HQL2025



- The following sum-rule for $D \rightarrow \pi\pi$ decays helps **determine the source of CPV**:

$$R = \frac{A_{CP}^{\text{dir}}(D^0 \rightarrow \pi^+ \pi^-)}{1 + \frac{\tau_{D^0}}{\mathcal{B}_{+-}} \left(\frac{\mathcal{B}_{00}}{\tau_{D^0}} - \frac{2}{3} \frac{\mathcal{B}_{+0}}{\tau_{D^+}} \right)} + \frac{A_{CP}^{\text{dir}}(D^0 \rightarrow \pi^0 \pi^0)}{1 + \frac{\tau_{D^0}}{\mathcal{B}_{00}} \left(\frac{\mathcal{B}_{+-}}{\tau_{D^0}} - \frac{2}{3} \frac{\mathcal{B}_{+0}}{\tau_{D^+}} \right)} + \frac{A_{CP}^{\text{dir}}(D^+ \rightarrow \pi^+ \pi^0)}{1 - \frac{3}{2} \frac{\tau_{D^+}}{\mathcal{B}_{+0}} \left(\frac{\mathcal{B}_{00}}{\tau_{D^0}} + \frac{\mathcal{B}_{+-}}{\tau_{D^0}} \right)}$$

- if $R \neq 0$, CPV: from $\Delta I=1/2$ amplitude; if $R=0$ & a $A_{CP}^{\text{dir}} \neq 0$, CPV: from a beyond-SM $\Delta I=3/2$ amplitude.
- $A_{CP}(D^0 \rightarrow \pi^+ \pi^-)$ precision: leading by LHCb; first evidence of direct CPV in a specific D decay.

- Raw asymmetry of the tagged $D^0 \rightarrow \pi^0 \pi^0$ sample:

$$A_{\text{raw}} = A_{CP}(D^0 \rightarrow \pi^0 \pi^0) + A_P^{D^*} + A_\varepsilon^{\pi_s} \quad \text{specificity}$$

- $A_P^{D^*}$: being **an odd function** of $\cos \theta^*$ i.e. the cosine of the charmed-meson polar angle in e^+e^- c.m.s
- $A_\varepsilon^{\pi_s}$: using tagged and untagged $D^0 \rightarrow K^- \pi^+$ samples

- Time-integrated CP asymmetry:

$$A_{CP}(D^0 \rightarrow \pi^0 \pi^0) = A'^{\pi^0 \pi^0} - A'^{K\pi} + A'^{K\pi, \text{untag}}; \text{ where } A'^f = \frac{A^f(\cos \theta^* < 0) + A^f(\cos \theta^* > 0)}{2}$$


$f = \pi^0 \pi^0; K\pi; K\pi, \text{untag}.$

Charm CPV: $A_{CP}(D^0 \rightarrow \pi^0 \pi^0)$

PRD 112, 012006 (2025)



$\cos \theta^*(D^{*+}) > 0$ vs. $\cos \theta^*(D^{*+}) < 0$



- Result at Belle II (428/fb):

$$A_{CP}^{\pi^0 \pi^0} = (+0.30 \pm 0.72 \pm 0.20)\%$$

→ only 15% less precise than Belle's result ($\sigma=0.65\%$)

based on 980/fb data [PRL 112, 211601 (2014)]

- Using our result, $A_{CP}^{\pi^+ \pi^-}$ (LHCb), W.A. $A_{CP}^{\pi^+ \pi^0}$,

ΔY (LHCb), W.A. BR, and D lifetimes,

$$R = (1.5 \pm 2.5) \times 10^{-3} \rightarrow 20\% \text{ improved precision}$$

More CPV results for charmed mesons:


$$D^0 \rightarrow \pi^+ \pi^- \pi^0, D_{(s)}^+ \rightarrow K_S^0 K^- \pi^+ \pi^+ \text{ (see backup)}$$

direct CPV in $\Lambda_c^+ \rightarrow \Lambda K^+, \Sigma^0 K^+$

- The sources of raw asymmetry of $\Lambda_c^+ \rightarrow \Lambda h^+$:

$$A_{\text{raw}}(\Lambda_c^+ \rightarrow \Lambda K^+) \approx \mathcal{A}_{CP}^{\Lambda_c^+ \rightarrow \Lambda K^+} + \mathcal{A}_{CP}^{\Lambda \rightarrow p\pi^-} + A_\epsilon^\Lambda + A_\epsilon^{K^+} + A_{FB}^{\Lambda_c^+}$$

- Using (CF) $\Lambda_c^+ \rightarrow \Lambda\pi^+, \Sigma^0\pi^+$ to remove **common sources**.




- $\mathcal{A}_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Lambda K^+) = (+2.1 \pm 2.6 \pm 0.1)\%$
- $\mathcal{A}_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Sigma^0 K^+) = (+2.5 \pm 5.4 \pm 0.4)\%$

First $\mathcal{A}_{CP}^{\text{dir}}$ for SCS two-body decays of charmed baryons.

α -induced CPV in $\Lambda_c^+ \rightarrow \Lambda K^+, \Sigma^0 K^+$

- Measure $\alpha/\bar{\alpha}$ for the separate $\Lambda_c^+/\bar{\Lambda}_c^-$ samples.
- Calculate $\mathcal{A}_{CP}^\alpha \equiv (\alpha_{\Lambda_c^+} + \alpha_{\bar{\Lambda}_c^-})/(\alpha_{\Lambda_c^+} - \alpha_{\bar{\Lambda}_c^-})$.



- $\mathcal{A}_{CP}^\alpha(\Lambda_c^+ \rightarrow \Lambda K^+) = -0.023 \pm 0.086 \pm 0.071$
- $\mathcal{A}_{CP}^\alpha(\Lambda_c^+ \rightarrow \Sigma^0 K^+) = +0.08 \pm 0.35 \pm 0.14$
- First $\mathcal{A}_{CP}^\alpha$ results for SCS decays of charmed baryons.
- No evidence of CPV is found.

粲重子CPV：Belle的结果之后，翘首以盼的Belle II的第一个结果呢？

$A_{CP}(\Lambda_c^+ \rightarrow ph^+h^-, \Xi_c^+ \rightarrow \Sigma^+h^+h^-)$

Preliminary result



- LHCb: $\Delta A_{CP}^{wgt} = A_{CP}(\Lambda_c^+ \rightarrow pK^+K^-) - A_{CP}^{wgt}(\Lambda_c^+ \rightarrow p\pi^+\pi^-) = (0.30 \pm 0.91 \pm 0.61)\%$ [JHEP 03(2018)182]

- Assuming U-spin symmetry

$$A_{CP}(\Lambda_c^+ \rightarrow pK^+K^-) + A_{CP}(\Xi_c^+ \rightarrow \Sigma^+\pi^+\pi^-) = 0$$

$$A_{CP}(\Lambda_c^+ \rightarrow p\pi^+\pi^-) + A_{CP}(\Xi_c^+ \rightarrow \Sigma^+K^+K^-) = 0$$

- Control sample:

- $\Xi_c^+ \rightarrow \Sigma^+h^+h^- : \Lambda_c^+ \rightarrow \Sigma^+h^+h^-$
- $\Lambda_c^+ \rightarrow ph^+h^- : \Lambda_c^+ \rightarrow p\pi^+K^-, D^0 \rightarrow K^-\pi^+\pi^+\pi^-$

- Belle II (428/fb) result:

$$A_{CP}(\Xi_c^+ \rightarrow \Sigma^+K^+K^-) = (3.7 \pm 6.6 \pm 0.6)\%,$$

$$A_{CP}(\Xi_c^+ \rightarrow \Sigma^+\pi^+\pi^-) = (9.5 \pm 6.8 \pm 0.5)\%,$$

$$A_{CP}(\Lambda_c^+ \rightarrow pK^+K^-) = (3.9 \pm 1.7 \pm 0.7)\%,$$


$$A_{CP}(\Lambda_c^+ \rightarrow p\pi^+\pi^-) = (0.3 \pm 1.0 \pm 0.2)\%,$$

first A_{CP} for
3-body SCS

- testing U-spin sum rules:

$$A_{CP}(\Xi_c^+ \rightarrow \Sigma^+\pi^+\pi^-) + A_{CP}(\Lambda_c^+ \rightarrow pK^+K^-) = (13.4 \pm 7.0 \pm 0.9)\%,$$

$$A_{CP}(\Xi_c^+ \rightarrow \Sigma^+K^+K^-) + A_{CP}(\Lambda_c^+ \rightarrow p\pi^+\pi^-) = (4.0 \pm 6.6 \pm 0.7)\%,$$



More data desired.

Outline



- Luminosity
- Recent B results
- Recent charm results
- Recent $q\bar{q}$ and exotic
- Recent tau and dark sector
- Summary


charm CPV

charm spectrum

charmed baryon

近期成果多数由“Belle II劳模”
沈成平（复旦）课题组主导

Belle (II): charm super-factory





PS: no Λ_b^0 sample

$D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma$


Preliminary result



- Precise nature of $D_{s0}(2317)^+$ remains unresolved, underscoring the need for new and improved data.
- The width of $D_s(2317)^+$ is unknown: $\Gamma(D_s(2317)^+) < 3.8 \text{ MeV} @95\% \text{ CL}$.



$$R = \frac{B(D_s(2317)^+ \rightarrow D_s^{*+} \gamma)}{B(D_s(2317)^+ \rightarrow D_s^+ \pi^0)} = [7.14 \pm 0.70 \pm 0.23]\%$$



李郁博(西交)+沈成平(复旦)为主导

Charmed baryon decay

JHEP 10(2024)045, JHEP 03(2025)061,
JHEP 08(2025)195, Preliminary result



BR results of 5 CF and 7 SCS decays of $\Xi_c^{0,+}$:

$$B(\Xi_c^0 \rightarrow \Xi^0 \pi^0)/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = 0.48 \pm 0.02 \pm 0.07$$

$$B(\Xi_c^0 \rightarrow \Xi^0 \eta)/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = 0.11 \pm 0.01 \pm 0.01$$

$$B(\Xi_c^0 \rightarrow \Xi^0 \eta')/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = 0.08 \pm 0.02 \pm 0.01$$

$$B(\Xi_c^+ \rightarrow p K_S^0)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = (2.47 \pm 0.16 \pm 0.07)\%$$

$$B(\Xi_c^+ \rightarrow \Lambda \pi^+)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = (1.56 \pm 0.14 \pm 0.09)\%$$

$$B(\Xi_c^+ \rightarrow \Sigma^0 \pi^+)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = (4.13 \pm 0.26 \pm 0.22)\%$$

$$B(\Xi_c^+ \rightarrow \Sigma^+ K_S^0)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = 0.067 \pm 0.007 \pm 0.003$$



$$B(\Xi_c^+ \rightarrow \Xi^0 \pi^+)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = 0.248 \pm 0.005 \pm 0.009$$

$$B(\Xi_c^+ \rightarrow \Xi^0 K^+)/B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+) = 0.017 \pm 0.003 \pm 0.001$$



$$B(\Xi_c^0 \rightarrow \Lambda \pi^0)/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) < xx\% \text{ @90\% C.L.}$$

$$B(\Xi_c^0 \rightarrow \Lambda \eta)/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = (xxx \pm 0.91 \pm 0.16)\%$$

$$B(\Xi_c^0 \rightarrow \Lambda \eta')/B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = (xxx \pm 0.82 \pm 0.16)\%$$



$$\frac{dN}{d \cos \theta_{\Xi^0}} \propto 1 + \alpha(\Xi_c^0 \rightarrow \Xi^0 h^0) \alpha(\Xi^0 \rightarrow \Lambda \pi^0) \cos \theta_{\Xi^0},$$




$$\alpha(\Xi_c^0 \rightarrow \Xi^0 \pi^0) = -0.90 \pm 0.15(\text{stat}) \pm 0.23(\text{syst})$$

$$B^{+,0} \rightarrow \Sigma_c(2455)^{+,0} \bar{\Xi}_c^{-,0}$$

PRD 112 (2025) L051101



- These two decays proceed through a purely internal W-emission:



- First searches for these decays:

$$B(B^+ \rightarrow \Sigma_c^{++} \bar{\Xi}_c^-) = (5.74 \pm 1.11 \pm 0.42^{+2.47}_{-1.53}) \times 10^{-4}$$





$$B(B^0 \rightarrow \Sigma_c^0 \bar{\Xi}_c^0) = (4.83 \pm 1.12 \pm 0.37^{+0.72}_{-0.60}) \times 10^{-4}$$

Vs. theoretical prediction by


- the QCD sum rule ^[1]: 4×10^{-3} ;
- diquark model ^[2]: $O(10^{-4})$

^[1]Nucl. Phys. B345, 137 (1990).

^[2]Z. Phys. C 51, 445 (1991)




Outline



- Luminosity
- Recent B results
- Recent charm results
- Recent $q\bar{q}$ and exotic
- Recent tau and dark sector
- Summary

Goal: understand the **nature of $\Upsilon(10753)$**



$\Upsilon(10753) \rightarrow \omega\chi_{bJ}$:

PRL 130, 091902 (2023)

$\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(nS)$:

JHEP 07 (2024) 116





$\Upsilon(10753) \rightarrow \omega\eta_b$:

PRD 109, 072013 (2024)

$\Upsilon(10753) \rightarrow B^{(*)}\bar{B}^{(*)}$:

JHEP 10 (2024) 114

$e^+e^- \rightarrow \omega\chi_{bJ}$ and $e^+e^- \rightarrow (\pi^+\pi^-\pi^0)_{\text{non-}\omega}\chi_{bJ}$ Preliminary result



$\Upsilon(10753)$ mass	$(10756.1 \pm 4.3) \text{ MeV}/c^2$
$\Upsilon(10753)$ width	$(32.2 \pm 18.7) \text{ MeV}$

Both are consistent with results from $e^+e^- \rightarrow \pi^+\pi^-\Upsilon(nS)$ [JHEP 07, 116 (2024)]

$$\frac{\sigma(e^+e^- \rightarrow \chi_{bJ}(1P)\omega)}{\sigma(e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-)}$$

- 1.5 at $\sqrt{s} \sim 10.75 \text{ GeV}$
- 0.15 at $\sqrt{s} \sim 10.867 \text{ GeV}$



This may indicate the difference in internal structures of $\Upsilon(10753)$ and $\Upsilon(10860)$.

$e^+e^- \rightarrow \eta Y(1,2S), \gamma \chi_{bJ}$ near 10.75 GeV

arXiv:[2509.01917](#) [hep-ex]
arXiv:[2508.16036](#) [hep-ex]





➤ $e^+e^- \rightarrow \eta Y(1,2S)$



Also search for $e^+e^- \rightarrow \gamma X_b$, where X_b is the bottomonium-sector partner of $X(3872)$. No evidence is found.



➤ $e^+e^- \rightarrow \gamma \chi_{bJ}$




$$B(Y(10753) \rightarrow \gamma \chi_{b1}) < 10^{-3}$$

$e^+e^- \rightarrow h^+h^- J/\psi$ ($h = \pi, K, p$) via ISR

Preliminary result



- indicative of the $Y(4230/4320)$ state
- an excess near 4.1 GeV



- Evidence for $Z_c(3900)^\pm$ (3.3σ)
- consistent with Belle/BESIII result

The precisions of cross section ($h=K, \pi$) based on Belle II (428/fb):


- comparable to those from BaBar/Belle using ISR.
- less precise (5% and 10%) than those from BESIII via energy scans

Belle II: Looking forward to more luminosity; results from ISR or $\gamma\gamma$ process.

- Luminosity
- Recent B results
- Recent charm results
- Recent $q\bar{q}$ and exotic
- Recent tau results
- Summary

Belle II: a tau super-factory

$$\sigma(e^+e^- \rightarrow \tau^+\tau^-(\gamma)) = 0.919 \text{ nb}$$



Advantages at Belle II:



- ✓ High luminosity
- ✓ Good vertexing and tracking capabilities
- ✓ Sophisticated trigger system and particle ID


tau LFV decays

JHEP 09 (2024) 062, PRD 110 (2024) 112003, JHEP 08 (2025) 092




Lepton flavour violation is only allowed by:

- Neutrino oscillations $\mathcal{O}(10^{-55})$
far beyond current experimental sensitivities
- New Physics models $\mathcal{O}(10^{-8})$
e.g. Leptoquarks for $\tau^- \rightarrow \ell^- V^0$ deals with $R(K^{*0})$ anomalies




$\mathcal{B}^{\text{UL}}(\tau \rightarrow \mu\mu\mu) < 1.9 \times 10^{-8}$
more restrictive than Belle

$\mathcal{B}^{\text{UL}}(\tau \rightarrow e(\mu)K_s^0) < 0.8(1.2) \times 10^{-8}$: most stringent




$\mathcal{B}^{\text{UL}}(\tau^- \rightarrow \Lambda \pi^- (\bar{\Lambda} \pi^-)) < 4.7(4.3) \times 10^{-8}$: most stringent



Summary



- Belle II: analyses more data (428/fb before 2023), develops new tools, recently achieved fruitful results, including some world-leading results.
 - B CPV/rare/hadronic, charm CPV/BF/ α /spectrum <- Recoiling/ γ / π^0 / K_S / ν etc.
 $q\bar{q}$ and exotic, τ physics, ($\gamma\gamma$ physics) etc. <- special data at e^+e^- collider
- Now Belle II: 575/fb, only 1% of targeted 50/ab.
- More data, more results + more ideas (from u)
- improvements as expected (未来“可期”)
+ unexpected excitements (未来可期)



❖ 固定教职工10人：

❖ 粒子物理与核物理**理论**：8人

研究强子谱/CP破坏/格点QCD/核物理理论等

❖ 粒子物理**实验**：2人

参加BESIII和Belle II实验

BESIII




研究粲强子/CP破坏/(类)粲偶素等

❖ 博士后1名； 在读研究生~22人


(不完全统计有博士生3人+硕士生19人)

2025.9@长沙




诚邀专家学者莅临指导！

Back up




Belle II big family



► Belle II 合作组（中国大陆）：1207 个成员（134：11%，第三位），124（15）个单位，28个国家/地区。
成员人数排序（>80）：德国，日本，中国，美国，意大利

First charm wave: lifetimes



most precise charm lifetimes:




$$\begin{aligned}\tau(D^0) &= 410.5 \pm 1.1 \pm 0.8 \text{ fs}, \\ \tau(D^+) &= 1030.4 \pm 4.7 \pm 3.1 \text{ fs}, \\ \tau(D_s^+) &= 499.5 \pm 1.7 \pm 0.9 \text{ fs}, \\ \tau(\Lambda_c^+) &= 203.20 \pm 0.89 \pm 0.77 \text{ fs}\end{aligned}$$

第一波粲物理结果: 基于早期数据的粲强子寿命的精确测量


PRL 127, 211801 (2021); PRL 131, 171803 (2023);

PRD 107, L031103 (2023); PRL 130, 071802 (2023).



Charm CPV: $A_{CP}(D^+ \rightarrow \pi^+ \pi^0)$

PRD 112, L031101 (2025)



$$N_{\text{sig}} = 5130 \pm 110$$

$$A_{CP} = (-3.9 \pm 1.8 \pm 0.2)\%$$

$$N_{\text{sig}} = 18510 \pm 240$$

$$A_{CP} = (-1.1 \pm 1.0 \pm 0.1)\%$$

Using $D^+ \rightarrow \pi^+ K_S^0$ (obtain $\sim 1.6\text{M}$ signals) to eliminate two common asymmetry sources: $A_{\text{prod}}^D + A_{\varepsilon}^{\pi^+}$. Thus, the CP asymmetry of interest is

$$A_{CP} = A_{\text{raw}}^{\pi^+ \pi^0} - A_{\text{raw}}^{\pi^+ K_S^0} + A_{\bar{K}^0}$$

Belle II (428/fb) result:

$$A_{CP} = (-1.8 \pm 0.9 \pm 0.1)\%$$

→ most precise; and 30% improved precision compared to Belle's result $\sigma=1.26\%$ (980/fb)

[PRD 97, 011101 (2018)]

Charm CPV: $A_{CP}(D^0 \rightarrow \pi^+\pi^-\pi^0)$

Preliminary result



$$A_{\text{raw}}^{\pi\pi\pi^0} = \frac{N(D^0 \rightarrow \pi^+\pi^-\pi^0) - N(\bar{D}^0 \rightarrow \pi^+\pi^-\pi^0)}{N(D^0 \rightarrow \pi^+\pi^-\pi^0) + N(\bar{D}^0 \rightarrow \pi^+\pi^-\pi^0)}$$

$$A_{\text{raw}}^{\pi\pi\pi^0} \simeq A_{CP} + A_{\text{prod}} + A_{\epsilon}^{\pi\pi\pi^0} + A_{\epsilon}^{\pi_{\text{tag}}}$$




$$A_{\epsilon}^{\pi_{\text{tag}}} = A_{\text{raw}}^{\text{tagged}} - A_{\text{raw}}^{\text{untagged}}$$

$$A_{\text{raw}}^{\text{tagged}} \simeq A_{\text{prod}} + A_{\epsilon}^{K\pi} + A_{\epsilon}^{\pi_{\text{tag}}}$$

$$A_{\text{raw}}^{\text{untagged}} \simeq A_{\text{prod}} + A_{\epsilon}^{K\pi},$$

$$A_{\pm i} = A_{\text{raw},\pm i}^{\pi\pi\pi^0} - A_{\text{raw},\pm i}^{\text{tagged}} + A_{\text{raw},\pm i}^{\text{untagged}}$$

$$A_{CP}^i = \frac{A_{+i} + A_{-i}}{2}$$



Time-integrated CP asymmetry using 428/fb@Belle II:


$$A_{CP}(D^0 \rightarrow \pi^+\pi^-\pi^0) = (0.29 \pm 0.27 \pm 0.13)\%$$

- consistent with CP symmetry
- 35% more precise than the world's best result from BaBar, despite an increase of only ~10%*Lumin.

Next step: Dalitz-plot- or time-dependent searches for CPV

Charm CPV: $D_{(s)}^+ \rightarrow K_S^0 K^- \pi^+ \pi^+$


JHEP 04 (2025) 036



三重积变量:

$$C_{TP} = \vec{p}_i \cdot (\vec{p}_j \times \vec{p}_k)$$




$M \rightarrow P_i P_j P_k P_l$



四重积变量:

$$C_{QP} = (\vec{p}_{K^-} \times \vec{p}_{\pi^+}) \cdot (\vec{p}_{K_S^0} \times \vec{p}_{\pi^+})$$

$C_{QP} > 0$: $\vec{p}_{K^-}$ at left-side of $\vec{p}_{K_S^0} \pi^+$ ($\vec{p}_{K_S^0} \times \vec{p}_{\pi^+}$) plane



$$A_X(D_{(s)}^+) = \frac{N_+(X > 0) - N_+(X < 0)}{N_+(X > 0) + N_+(X < 0)}$$
$$A_{\bar{X}}(D_{(s)}^-) = \frac{N_-(\bar{X} > 0) - N_-(\bar{X} < 0)}{N_-(\bar{X} > 0) + N_-(\bar{X} < 0)}$$
$$\mathcal{A}_{CP}^X = \frac{1}{2}(A_X(D_{(s)}^+) - A_{\bar{X}}(D_{(s)}^-))$$

X	Combined $\mathcal{A}_{CP}^X (10^{-3})$
C_{TP}	$-2.3 \pm 4.5 \pm 1.5$
C_{QP}	$-0.7 \pm 4.5 \pm 1.7$
$C_{TP} C_{QP}$	$+3.9 \pm 4.5 \pm 1.1$
$\cos \theta_{K_S^0} \cos \theta_{K^-}$	$-2.9 \pm 4.5 \pm 2.1$
$C_{TP} \cos \theta_{K_S^0} \cos \theta_{K^-}$	$+1.0 \pm 4.5 \pm 1.4$
$C_{QP} \cos \theta_{K_S^0} \cos \theta_{K^-}$	$+11.6 \pm 4.5 \pm 1.1$

Belle 980/fb: 0.59%; Belle II 424/fb: 0.70%

Belle II首批粲CPV结果

精度4.5‰接近 $A_{CP}^{dir}(D^0 \rightarrow \pi^+ \pi^-) = 2.3\text{‰}$

李龙科 (湖南师大) 为主导