


Toponium: Experimental Results



李海峰




山东大学（青岛）

TeV物理前沿专题研讨会，2025年10月11-12日，青岛

Top Quark Pair Production at LHC


- LHC is a $t\bar{t}$ factory
 - $\sigma_{t\bar{t}} = 834 \text{ pb}$ at LHC Run 2
 - 0.83M $t\bar{t}$ events per fb^{-1}
 - Due to the short life time, can measurement $t\bar{t}$ spin correlations
- With those **huge amount** of $t\bar{t}$ data, ATLAS has carried out precision measurements in top quark physics
- Thanks to the advanced MC generators and high-order QCD/EW calculations
- In this talk, **focus on $t\bar{t}$ production threshold region** with 140 fb^{-1} LHC Run 2 pp data




Threshold Region Measurement is Challenging

Experimentally very challenging: modelling of $t\bar{t}$ close to threshold region; tiny effect of quasi-bound state

Previous hints



First Quantum Entanglement (QE) measurement using $t\bar{t}$ at LHC



Threshold region has received a lot of attention recently in the context of quantum entanglement

Stronger QE in data than MC.
Missing toponium contributions?

Top quark and $t\bar{t}$ Threshold Region


- Top quark is very special. Heaviest quark in the SM. Has largest Yukawa coupling to the Higgs field
- Very short life time $\rightarrow$ decays before forming any real hadron
- QCD predicts a quasi-bound state close to the threshold for low momentum top quarks (the prediction was made even before the top quark discovery)

部分理论文章:

V.S. Fadin and V.A. Khoze, JETP Lett. 46 (1987) 525

Y. Sumino *et al.*, Phys. Rev. D 47 (1993) 56


W.-L. Ju *et al.*, JHEP 06 (2020) 158



ATLAS Results

Top-antitop production near threshold

From B. Fuks



The toponium Green's function

From B. Fuks

$$K_{abcd}(x, y, z) = \langle 0 | T \{ t_c(y) \bar{t}_d(z) : \bar{t}_a(x) t_b(x) : \} | 0 \rangle$$

$$= \frac{(1 + \gamma^0)_{ca}}{2} \frac{(1 - \gamma^0)_{bd}}{2} \int d^3r \left[K_1(y; (z^0, \vec{r})) K_2(z^0, \vec{r}, \vec{z}; x^0, \vec{x}, \vec{x}) + K_1(z; (y^0, \vec{r})) K_2(y^0, \vec{y}, \vec{r}; x^0, \vec{x}, \vec{x}) \right]$$

Non-relativistic spin projection operators

Antitop-decay first

Top-decay first

1-particle-state and 2-particle-state propagators

The toponium Green's function

- Solution to the Lippmann-Schwinger equation
 - Fourier transform of the QCD potential
 - S-wave contributions
- To be solved numerically

$$\widetilde{G}(E; p) = \widetilde{G}_0(E; p) + \int \frac{d^3q}{(2\pi)^3} \widetilde{V}_{\text{QCD}}(\vec{p} - \vec{q}) \widetilde{G}(E; q)$$

Free Green's function


Quasi-bound State from NRQCD

- S-wave, color-singlet state with Green's function of non-relativistic (NR) QCD by [B. Fuks *et al.*, Eur. Phys. J. C 85 \(2025\) 157](#)
- Generate $gg \rightarrow tt \rightarrow b\ell\nu b\ell\nu$ with MG5_aMC. Spin correlations included
- Reweight matrix element with QCD Green's functions

$$|\mathcal{M}|^2 \rightarrow |\mathcal{M}|^2 \left| \frac{\tilde{G}(E; p^*)}{\tilde{G}_0(E; p^*)} \right|^2$$

$\tilde{G}$: Green's function considering QCD potential

$\tilde{G}_0$: Free Green's function



This model includes NRQCD calculations. More complete w.r.t. previous simplified models (using scalar/pseudoscalar as an effective model)

Background Modelling

Extremely challenging measurement: need precise modelling of the $t\bar{t}$ threshold region

- $t\bar{t}$: main background. Powheg v2 hvq + Pythia8, using narrow-width approximation (NWA). $m_t = 172.5$ GeV
 - 2D reweighting in $(\cos\theta^*, M(t\bar{t}))$ to NNLO QCD (from MATRIX) and NLO EW (HATHOR)
 - θ^* : angle between the momentum of the top quark in the $t\bar{t}$ center-of-mass frame and the momentum of the $t\bar{t}$ system in the lab. frame
- $t\bar{t}$: alternative MC sample (for syst.), Powheg v2 bb4l + Pythia8
 - Simulate $pp \rightarrow b\ell\nu b\ell\nu$ including off-shell, non-resonant contributions, and exact spin correlations at NLO

Event Selections

Target for dilepton channel $tt \rightarrow b\ell\nu b\ell\nu$

SR: Signal Region;

CR: Control Region

SRs	CR-Z	CR-Fakes
$= 2\ell$ with $p_T(\ell) \geq 10$ GeV ≥ 1 trigger-matched lepton with $p_T \geq 25/27/28$ GeV ≥ 2 jets with $p_T \geq 25$ GeV ≥ 1 b -tagged jet (70% efficiency WP) $m_{\ell\ell} \geq 15$ GeV $m_{t\bar{t}} \leq 500$ GeV		
$E_T^{\text{miss}} \geq 60$ GeV for OSSF events		—
$\ell^\pm \ell'^\mp$	$e^\pm e^\mp / \mu^\pm \mu^\mp$	$\ell^\pm \ell'^\pm$
$ m_{\ell\ell} - m_Z \geq 10$ GeV	$ m_{\ell\ell} - m_Z \leq 10$ GeV	$ m_{\ell\ell} - m_Z \geq 10$ GeV

OSSF: opposite-sign, same-flavor

CRs are for correcting Z+jets and Fakes normalization in fit


Event Categorization

SR events are categorized into 9 regions based on two observables: c_{hel} and c_{han}

$$c_{hel} = \vec{\ell}_+ \cdot \vec{\ell}_-,$$

where the $\vec{\ell}_\pm$ are the lepton directions in $t\bar{t}$ center-of-mass frame, and then in turn boosted into t and $\bar{t}$ frames. This distribution has a maximum slope for a spin-singlet state

c_{han} : flip the $\vec{\ell}$ in t direction. This distribution has a maximum slope for a spin-triplet state



c_{hel} is useful to separate pseudoscalar from other contributions

Event Categorization and Fitting

	$-1 < c_{hel} < -\frac{1}{3}$	$-\frac{1}{3} < c_{hel} < \frac{1}{3}$	$\frac{1}{3} < c_{hel} < 1$
$-1 < c_{han} < -\frac{1}{3}$	SR1	SR2	SR3
$-\frac{1}{3} < c_{han} < \frac{1}{3}$	SR4	SR5	SR6
$\frac{1}{3} < c_{han} < 1$	SR7	SR8	SR9


CR-Fakes ee	CR-Fakes $e\mu$	CR-Fakes $\mu\mu$	CR-Z
---------------	-----------------	-------------------	------

Simultaneous fitting to $m_{t\bar{t}}$ with 13 categories with profile likelihood method

Background Estimations

- $t\bar{t}$: with a free-floating scale factor (SF); tW : estimation from MC
- **Z+jets**: get some contributions from $Z \rightarrow \tau\tau$. Use the CR-Z to normalize the Z+b process


- **fake / non-prompt leptons**: Fakes represent 1.5% of SR yields. Data-driven estimation with 3 CR-Fakes



Results: baseline $t\bar{t}$ + quasi-bound state (NRQCD)

Observed (expected)
significance: 7.7σ (5.7σ)

Goodness-of-Fit: 0.93



$$\sigma(t\bar{t}_{\text{NRQCD}}) = 9.0 \pm 1.3 \text{ pb} = 9.0 \pm 1.2 \text{ (stat.)} \pm 0.6 \text{ (syst.) pb}$$



Run: 338183
Event: 3295623881
2017-10-14 09:08:09 CEST

muon

electron

b-jet 1


b-jet 2

Missing E_T

$$m_{t\bar{t}} = 342 \text{ GeV}$$


$$c_{hel} = 0.97$$

$$c_{chan} = 0.94$$



Impacts of Systematics

- Quasi-bound state modelling: Parton shower [Herwig7]
- $t\bar{t}$ decay and off-shell [comparison to bb4l]
- NNLO QCD rew.: NNLO QCD scale variations
- No strong pulls or constraints
- Largest effects from toponium modelling and off-shell effect modelling



Impacts of Systematics


ATLAS-CONF-2025-008

- Quasi-bound state modelling: Parton shower [Herwig7]
- $t\bar{t}$ decay and off-shell [comparison to bb4l]
- NNLO QCD rew.: NNLO QCD scale variations
- No strong pulls or constraints
- Largest effects from toponium modelling and off-shell effect modelling

Category	Impact
$t\bar{t}_{\text{NRQCD}}$ modelling	5.3%
$t\bar{t}$ modelling	3.5%
Jet energy scale (pileup)	1.3%
b -tagging	1.2%
Instrumental (other)	0.9%
Limited MC statistics	0.7%
Jet energy scale (flavour)	0.5%
Background normalisations	0.4%
tW modelling	0.4%
Jet energy scale (η inter-calibration)	0.4%
Jet energy scale (other)	0.3%
Jet energy resolution	0.3%
Leptons	0.2%
Total syst. uncertainties	6.8%
Total stat. uncertainties	13%

Ratios of the pre-fit distributions for $t\bar{t}$ MC models vs. baseline Powheg hvq

- Low $m_{t\bar{t}}$ region: bb4l is more similar to hvq+toponium than hvq only
- High $m_{t\bar{t}}$ region: bb4l differs from hvq+toponium model




CMS Results

CMS Results

- [arXiv:2503.22382](https://arxiv.org/abs/2503.22382), [Rep. Prog. Phys. 88 \(2025\) 087801](#)
- Use very similar analysis method compared with ATLAS
- Use toy model for signal

显著度大于 5σ




Systematics



CMS

- Fit constraint (FO pQCD + BG + η_t)
- Fit constraint (FO pQCD + BG only)
- +1 σ impact (FO pQCD + BG + η_t)
- 1 σ impact (FO pQCD + BG + η_t)




$$\hat{\sigma}(\eta_t) = 8.8^{+1.2}_{-1.4} \text{ pb}$$



S+B fitting




Comparison of different models



比较ATLAS和CMS的signal model


LHC Top WG meeting 4-6 June 2025, CERN



Elusive romance of top-quark pairs observed at the LHC

The CMS and ATLAS experiments at CERN's Large Hadron Collider have observed an unforeseen feature in the behaviour of top quarks that suggests that these heaviest of all elementary particles form a fleeting union

8 JULY, 2025



Artist's impression of the short-lived union of a top quark and a top antiquark formed by the exchange of gluons. (Image: D. Dominguez/CERN)

An unforeseen feature in proton-proton collisions previously observed by the CMS experiment at CERN's Large Hadron Collider (LHC) has now been confirmed by its sister experiment ATLAS. The result, reported yesterday at the European Physical Society's High-Energy Physics conference in Marseille, suggests that top quarks – the heaviest and shortest-lived of all the elementary particles – can momentarily pair up with their [antimatter](#) counterparts to produce a “quasi-bound-state” called toponium. Further input based on complex theoretical calculations of the strong nuclear force -- called quantum chromodynamics (QCD) -- will enable physicists to understand the true nature of this elusive dance.

CERN COURIER

Reporting on international high-energy physics

Physics ▾

Technology ▾


Community ▾

Magazine

STRONG INTERACTIONS | NEWS

ATLAS confirms top–antitop excess

9 September 2025




Quasi-bound candidate An event display of an interaction consistent with the formation of toponium in the ATLAS detector. The final state

At the LHC, almost all top–antitop pairs are produced in a smooth invariant-mass spectrum described by perturbative QCD. In March, the CMS collaboration announced the discovery of an additional 1% localised near the energy threshold to produce a top quark and its antiquark ([CERN Courier May/June 2025 p7](#)). The ATLAS collaboration has now confirmed this observation.

“The measurement was challenging due to the small cross section and the limited

Celebration after toponium talks from ATLAS and CMS during EPS-HEP2025
First report about ATLAS results



Conclusions

- An excess of events is observed over the NNLO perturbative QCD prediction, with **7.7 σ observed (5.7 σ expected)** near the $t\bar{t}$ production threshold by ATLAS with LHC Run 2 data. [ATLAS-CONF-2025-008], [ATLAS Physics Briefing]
- This excess is consistent with **color-singlet, S-wave, quasi-bound $t\bar{t}$ states** predicted by NRQCD with **cross-section of 9.0 ± 1.3 pb**
- A more complete model from NRQCD calculation is used in this analysis. Important advantage compared with recent CMS results ([Rep. Prog. Phys. 88 \(2025\) 087801](#))
- This results show the strength of the LHC as a precision machine
- **Observation of toponium** opens a new field to study NRQCD with top quarks
- More work to characterize this excess and to better quantify the impact of off-shell top-quark decays



Top quark and anti top quark “shake their hands” before they decay

Backup

Simplified model for $t\bar{t}$ quasi-bound states

- A pseudo-scalar spin-singlet resonance or as a combination of scalar and pseudo-scalar resonances. Mostly pseudo-scalar $^1S^{[1]}$ configuration
- The contributions from states with higher total angular momentum J and color-octet states are expected to be sub-dominant and are neglected in this model
- In contrast, the main $t\bar{t}$ quasi-bound-state model used in this study includes the full set of S -wave color-singlet contributions, incorporating both bound state and scattering-state effects


More Information for Fitting

- Profile likelihood fitting to $m_{t\bar{t}}$
- **Control regions**: use the 3 CR-Fakes to extract scale factors (SF) for heavy-flavor and photon-conversion electron fakes, and heavy-flavor muon fakes; use the CR-Z to normalize the Z+b process
- **Signal regions**: use the 9 SRs to extract a SF for regular $t\bar{t}$ and signal strength for quasi-bound state $t\bar{t}$
- **Nuisance parameters (NPs) correlation scheme for constraint ones**: In the case of NPs that are strongly constrained to less than 50% of their prior uncertainty, the original systematic variation is treated as partially (50%) correlated between SRs and CRs


Alternative fit: Powheg hvq + simplified model of toponium

Observed
(expected)
significance
of 7.8σ (4.0σ)

Goodness-of-
Fit: 0.95




ATLAS-CONF-2025-008




Alternative fit: Powheg bb4l + quasi-bound state (NRQCD)

Observed
(expected)
significance of
 4.3σ (6.3σ)

Goodness-of-
Fit: 0.54



ATLAS-CONF-2025-008



Reconstruction of the $t\bar{t}$ System

- Ellipse Method [NIM A 736 (2014) 169]: geometric/analytic approach that imposes W and top mass constraints
- The two b -tagged jets associated with the decays of the top and antitop quark are chosen from all selected hadronic jets. If more than two of them are b -tagged, the two leading b -tagged jets are selected. If there is only one b -tagged jet, the highest- p_T jet among the remaining untagged ones is selected.
- The Ellipse Method provides a solution for about 95% of $t\bar{t}$ dilepton events

Systematics

- **Experimental systematics**: Electron, muon, jet, b-jet tagging, Missing E_T , pileup, and luminosity
- **$t\bar{t}$ modelling**: Parton shower [Herwig7], decay and off-shell [comparison to bb4l], matching [$p_T^{\text{hard}}, h_{\text{damp}}$], recoil-to-top, underlying event [A14 Var1], color reconnection [CR0 vs CR1/CR2 models], top mass [± 0.5 GeV], ISR/FSR, PDF4LHC15, NNLO QCD scale variations, NLO EW scheme
- **tW modelling**: Parton shower [Herwig7], matching [$p_T^{\text{hard}}, h_{\text{damp}}$], interference scheme [DR/DS], top mass [± 0.5 GeV]
- **Background normalizations**: 4% tW , 20% $t\bar{t}+b$, 40% $t\bar{t}+c$, 50% $\text{top}+X$, 50% $Z+c/\text{light}$, 50% diboson
- **Quasi-bound state modelling**: μ_r/μ_f variations, PDF4LHC21, parton shower [Herwig7], ISR/FSR

