

# ***Threshold Production of Unstable Top***

**Alexander Penin**

*University of Alberta & TTP Karlsruhe*

**Quarkonium Workshop**

*IHEP, Beijing, 22-26 April 2013*

# Topics discussed

- Top-antitop threshold production
  - *brief introduction and review*

# Topics discussed

- Top-antitop threshold production
  - *brief introduction and review*
- Top width effect:
  - *beyond the complex energy shift*
  - *effective theory of unstable particles “ $\rho$ NRQCD”*
  - *unstable top production in NLO and NNLO*

# Topics discussed

- Top-antitop threshold production
  - *brief introduction and review*
- Top width effect:
  - *beyond the complex energy shift*
  - *effective theory of unstable particles “ $\rho$ NRQCD”*
  - *unstable top production in NLO and NNLO*
- Based on: *A. Penin, J. Piclum, JHEP 1201 (2012) 034*

# Why top-antitop threshold at a LC?

## ● Theory

- *top quark width is a natural infrared cutoff*
- ➔ *first principle QCD predictions*

## ● Experiment

- *as clean as possible for a strongly interacting particle*


## ● Phenomenology

- *most precise determination of top quark*

*mass, width, vector couplings*


# Born cross section

$$R(e^+e^- \rightarrow t\bar{t})$$



# Coulomb and finite width effects

$R(e^+e^- \rightarrow t\bar{t})$



$$R_{\text{res}} \sim \frac{\alpha_s^3}{m_t \Gamma_t}, \quad E_{\text{res}} \sim \alpha_s^2 m_t$$

# Perturbative expansion

- Apparent slow convergence

→ *Full  $N^3LO$  analysis is mandatory*



# Perturbative expansion

- Apparent slow convergence

→ *Full  $N^3LO$  analysis is mandatory*

- *Resonance energy*

A. Penin, M. Steinhauser,  
Phys.Lett. B538 (2002) 335-345

- *Resonance cross section*

Alberta (A. Penin *et al.*)  
Karlsruhe (M. Steinhauser *et al.*)  
Munich (M. Beneke *et al.*)

# Finite top lifetime

- Resonant approximation

- *complex energy shift*  $E \rightarrow E + i\Gamma_t$

(V.Fadin, V.Khoze, JETP Lett. 46 (1987) 525)

- *not consistent in pNRQCD beyond LO!*

# Finite top lifetime

## ● Resonant approximation

- *complex energy shift*  $E \rightarrow E + i\Gamma_t$

(V.Fadin, V.Khoze, JETP Lett. 46 (1987) 525)

- *not consistent in pNRQCD beyond LO!*

## ● Nonresonant contribution

- *Phase space matching (tight cuts on top invariant mass)*

(A. Hoang, C. Reißer, P. Ruiz-Femenía, Phys. Rev. D82 (2010) 014005)


- *QCD effective theory of unstable particles to NLO*

(M. Beneke, B. Jantzen, P. Ruiz-Femenía, Nucl. Phys. B840 (2010) 186)

- *NRQCD effective theory of unstable particles to NNLO*

(A. Penin, J. Piclum, JHEP 1201 (2012) 034)

# Stable top




Optical theorem:

$$R_{res}^{Born} \sim \Im \int \frac{d^{d-1} \mathbf{p}}{(2\pi)^{d-1}} \frac{1}{\mathbf{p}^2 - m_t E - i\epsilon} \sim \Im \sqrt{-E - i\epsilon},$$

On-shell top:

$$\Im \left[ \frac{1}{\mathbf{p}^2 - m_t E - i\epsilon} \right] \sim \delta(\mathbf{p}^2 - m_t E),$$

# Unstable top




Imaginary part of mass operator:

here  $\rho = 1 - M_W/m_t$ ,  $z = (\mathbf{p}^2 - m_t E)/m_t^2 \ll 1$

$$\Im[\Sigma(z)] = \frac{\Gamma_t}{2} - \frac{\Gamma_t}{2} \left[ \theta(z - \rho) + \left( \frac{2z}{\rho} - \frac{z^2}{\rho^2} \right) \theta(\rho - z) + \mathcal{O}(\rho, z) \right]$$

# Unstable top



Imaginary part of mass operator:

here  $\rho = 1 - M_W/m_t$ ,  $z = (\mathbf{p}^2 - m_t E)/m_t^2 \ll 1$

$$\Im[\Sigma(z)] = \frac{\Gamma_t}{2} - \frac{\Gamma_t}{2} \left[ \theta(z - \rho) + \left( \frac{2z}{\rho} - \frac{z^2}{\rho^2} \right) \theta(\rho - z) + \mathcal{O}(\rho, z) \right]$$

*Resonant contribution*

*Nonresonant contribution*

# Resonant contribution

- Complex energy shift:

- *Dyson resummation*

$$\frac{1}{\mathbf{p}^2 - m_t E - i\epsilon} \rightarrow \frac{1}{\mathbf{p}^2 - m_t E - im_t \Gamma_t}$$

- *Breit-Wigner resonance*

$$\delta(\mathbf{p}^2 - m_t E) \rightarrow \frac{1}{\pi} \frac{\Gamma_t}{(\mathbf{p}^2 - m_t E)^2 + m_t^2 \Gamma_t^2},$$

- *Born cross section*

$$R_{res}^{Born} \sim \Im \left[ \sqrt{-E - i\Gamma_t} \right]$$

- Invariant mass distribution:

- $2\mathbf{p}^2 \approx m_t^2 - (p_W + p_b)^2$

# Nonresonant contribution

- On-shell  $t \Leftrightarrow$  on-shell  $W$  and  $b$ 
  - *kinematical constraint*  $M_W^2 < (p_W + p_b)^2 < m_t^2$
  - *natural cutoff on spatial momentum*  $0 < \mathbf{p}^2 < \rho m_t^2$
  - $\Im[\Sigma] - \Gamma_t/2 \neq 0$  for  $\mathbf{p}^2 \neq 0 \Leftrightarrow$  "nonresonant"



# Nonresonant contribution

- On-shell  $t \Leftrightarrow$  on-shell  $W$  and  $b$

- *kinematical constraint*  $M_W^2 < (p_W + p_b)^2 < m_t^2$

- *natural cutoff on spatial momentum*  $0 < \mathbf{p}^2 < \rho m_t^2$

- $\Im[\Sigma] - \Gamma_t/2 \neq 0$  for  $\mathbf{p}^2 \neq 0 \Leftrightarrow$  "nonresonant"

- Approximation  $\rho \ll 1$

- *nonrelativistic  $t$  and  $W$ , ultrarelativistic  $b$*

- *expansion in  $\rho$  similar to pNRQCD expansion in  $v^2 \sim E/m_t$*

- *actual value  $\rho = 0.53 \dots$*

# Nonrelativistic effective theory of unstable top

## • Scales

### • $pNRQCD$ :

*hard*  $m_t$       *soft*  $v m_t$       *ultrasoft*  $v^2 m_t$

### • $\rho NRQCD$ :

*hard*  $m_t$        $\rho$ -*soft*  $\rho^{1/2} m_t$        $\rho$ -*ultrasoft*  $\rho m_t$

# Nonrelativistic effective theory of unstable top

## • Scales

### • $pNRQCD$ :

*hard*  $m_t$       *soft*  $v m_t$       *ultrasoft*  $v^2 m_t$

### • $\rho NRQCD$ :

*hard*  $m_t$        $\rho$ -*soft*  $\rho^{1/2} m_t$        $\rho$ -*ultrasoft*  $\rho m_t$

## • Scale hierarchy and power counting

•  $pNRQCD$  scaling:  $\alpha_{ew}^{1/2} \sim \alpha_s \sim v \ll 1$ ,  $\Gamma_t/m_t \sim \alpha_{ew}$

• complimentary expansion in  $\rho$  with  $v \ll \rho^{1/2} \ll 1$

•  $\rho$ -Coulomb terms  $\alpha_s/\rho^{1/2} \ll 1$

# Nonrelativistic effective theory of unstable top

- How to expand?
  - $\rho$ - $pNRQCD$  Feynman rules
  - expansion by regions

# Nonrelativistic effective theory of unstable top

● How to expand?

✗  $\rho$ - $p$ NRQCD Feynman rules

✓ expansion by regions

# NLO nonresonant contribution

- Power counting

- *resonant contribution*
- *nonresonant contribution*

$$\Im\sqrt{-E - i\Gamma_t} \sim v$$
$$\Gamma_t \sim v^2$$

# NLO nonresonant contribution

## ● Power counting

- *resonant contribution*
- *nonresonant contribution*

$$\Im\sqrt{-E - i\Gamma_t} \sim v$$
$$\Gamma_t \sim v^2$$

## ● Calculation steps

- *treat  $\Im[\Sigma] - \Gamma_t/2$  as a perturbation*
  - *add all the two-loop diagrams with  $t$ - $W$ - $b$  cut*
  - *expand in  $E/\rho m_t$ , expand in  $\rho \Leftrightarrow$  single region left:  
 $t$  and  $W$  are  $\rho$ -potential,  $b$  is  $\rho$ -ultrasoft*
- ➔ *recover nonrelativistic propagators and vertices*

# NLO nonresonant contribution

## ● Power counting

- *resonant contribution*
- *nonresonant contribution*

$$\Im\sqrt{-E - i\Gamma_t} \sim v$$
$$\Gamma_t \sim v^2$$

## ● Calculation steps











- *treat  $\Im[\Sigma] - \Gamma_t/2$  as a perturbation*
- *add all the two-loop diagrams with  $t$ - $W$ - $b$  cut*
- *expand in  $E/\rho m_t$ , expand in  $\rho \Leftrightarrow$  single region left:  
 $t$  and  $W$  are  $\rho$ -potential,  $b$  is  $\rho$ -ultrasoft*

➔ *recover nonrelativistic propagators and vertices*











- *no expansion in  $\rho \Leftrightarrow$  fully relativistic calculation (M. Beneke et al.)*



# NLO diagrams



# NLO diagrams



# NLO result

- leading term of  $\rho$ -expansion

$$R_{nr}^{NLO} = -\frac{24}{\pi\rho^{1/2}} \frac{\Gamma_t}{m_t} \left[ \frac{4}{9} + \text{“Z”} - \frac{1}{\sin^4 \theta_W} \left( \frac{17}{48} - \frac{9\sqrt{2}}{32} \ln(1 + \sqrt{2}) \right) \right]$$

# NLO result

- leading term of  $\rho$ -expansion


$$R_{nr}^{NLO} = -\frac{24}{\pi\rho^{1/2}} \frac{\Gamma_t}{m_t} \left[ \frac{4}{9} + \text{“Z”} - \frac{1}{\sin^4 \theta_W} \left( \frac{17}{48} - \frac{9\sqrt{2}}{32} \ln(1 + \sqrt{2}) \right) \right]$$

- Convergence?

- *generally not bad*
- *for some diagrams Padé is necessary*


# Convergence

*diagram "j"*



# Convergence

*all diagrams*




*dash line - leading  $\rho$ -dependence*

*solid line - exact  $\rho$ -dependence*

# NNLO nonresonant contribution

$\rho$ -leading diagrams



## ● Regions of gluon momentum

- (a) and (c) - hard, potential,  $\rho$ -potential
- (b) and (d) - hard,  $\rho$ -soft

# NNLO result

## ● leading term of $\rho$ -expansion

$$\begin{aligned} R_{nr}^{N^2LO} = & \frac{3C_F\alpha_s}{\pi^2\rho^{1/2}} \frac{\Gamma_t}{m_t} \left\{ \left[ \frac{4}{9} + \text{“Z”} \right] \left[ \frac{\pi^2}{\rho^{1/2}} \left( 3 \ln \left( \sqrt{E^2 + \Gamma_t^2/\rho m_t} \right) + \frac{3}{2} + 6 \ln 2 \right) + (18 + 24 \ln 2) \right] \right. \\ & + \frac{1}{\sin^4 \theta_W} \left[ \frac{22}{3} + \frac{17\pi^2}{6} - \frac{17}{2} \ln 2 + (2 - 3\pi^2 + 9 \ln 2) \frac{3\sqrt{2}}{4} \ln (1 + \sqrt{2}) \right. \\ & \left. \left. - \frac{27\sqrt{2}}{8} \left( \ln^2 (1 + \sqrt{2}) + \text{Li}_2 (2\sqrt{2} - 2) \right) \right] \right\}. \end{aligned}$$



# NNLO result


## ● leading term of $\rho$ -expansion

$$R_{nr}^{N^2LO} = \frac{3C_F\alpha_s}{\pi^2\rho^{1/2}} \frac{\Gamma_t}{m_t} \left\{ \left[ \frac{4}{9} + \text{“Z”} \right] \left[ \frac{\pi^2}{\rho^{1/2}} \left( 3 \ln \left( \sqrt{E^2 + \Gamma_t^2/\rho m_t} \right) + \frac{3}{2} + 6 \ln 2 \right) + (18 + 24 \ln 2) \right] \right. \\ \left. + \frac{1}{\sin^4 \theta_W} \left[ \frac{22}{3} + \frac{17\pi^2}{6} - \frac{17}{2} \ln 2 + (2 - 3\pi^2 + 9 \ln 2) \frac{3\sqrt{2}}{4} \ln(1 + \sqrt{2}) \right. \right. \\ \left. \left. - \frac{27\sqrt{2}}{8} \left( \ln^2(1 + \sqrt{2}) + \text{Li}_2(2\sqrt{2} - 2) \right) \right] \right\}.$$

●  $\rho$ -Coulomb term  $\alpha_s/\rho^{1/2}$

● new type of logs  $\ln(E/\rho m_t) \sim \ln(v^2/\rho)$

# Numerics



*dot line - LO*

*dash line - LO+NLO nonresonant*

*solid line - LO+NNLO nonresonant*

# Summary

- Effective theory of  $\rho$ NRQCD

# Summary

- Effective theory of  $\rho$ NRQCD
  - *based on nonrelativistic expansion in  $\rho = 1 - m_t/M_W$*
  - *systematically accounts for finite width effects in threshold top-antitop production*
  - *optimized for high-order calculations*
  - *solve the problem of the spurious divergences*

# Summary

- Effective theory of  $\rho$ NRQCD
  - *based on nonrelativistic expansion in  $\rho = 1 - m_t/M_W$*
  - *systematically accounts for finite width effects in threshold top-antitop production*
  - *optimized for high-order calculations*
  - *solve the problem of the spurious divergences*
  - *conceptually clear and aesthetically appealing*