

正负电子对撞机上的新物理和暗物质研究

searching for new light dark particles

刘佐伟

南京大学

味物理前沿研讨会暨味物理讲座100期特别活动

2026年1月30日 - 2月4日

dark sector


Standard
Model

dark
sector


dark sector



parameter space of interest for electron colliders




parameter space of interest for electron colliders



dark matter detection

[Cirelli, Strumia, Zupan, 2406.01705v3]

Spin-independent DM detection



see also

2507.11930 (PandaX-4T)

2510.21458 (CDEX)

Outline

- 1 dark photon & millicharged particles
- 2 dark matter
- 3 leptophilic gauge boson
- 4 axion & axion-like particles (ALPs)

1

dark photon & millicharged particles

Hypercharge portal models

Hypercharge portal model for dark photon & millicharged particles

$$SU(3)_c \times SU(2)_L \times U(1)_Y$$

SM
sector


$$U(1)_X$$

dark
sector

[Holdom 1986; Foot & He 1991; Cheung & Yuan 2007; Feldman, ZL, Nath 2007]

Hypercharge portal models

Hypercharge portal model for dark photon & millicharged particles



[Holdom 1986; Foot & He 1991; Cheung & Yuan 2007; Feldman, ZL, Nath 2007]

Kinetic mixing & mass mixing

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$$

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi -\frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} -\frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

Kinetic mixing & mass mixing

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$$

$$\mathcal{L} = -\frac{1}{4} A_{\mu\nu} A^{\mu\nu} - \frac{1}{4} X_{\mu\nu} X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

Kinetic mixing & mass mixing

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$$

$$\mathcal{L} = -\frac{1}{4} A_{\mu\nu} A^{\mu\nu} - \frac{1}{4} X_{\mu\nu} X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

Kinetic mixing & mass mixing

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$$

$$\mathcal{L} = -\frac{1}{4} A_{\mu\nu} A^{\mu\nu} - \frac{1}{4} X_{\mu\nu} X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

kinetic mixing

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

Kinetic mixing & mass mixing

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_X$$

$$\mathcal{L} = -\frac{1}{4} A_{\mu\nu} A^{\mu\nu} - \frac{1}{4} X_{\mu\nu} X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

↑ kinetic mixing ↑ mass mixing

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical
- X_μ becomes **dark photon** (often denoted as A'_μ) if $M_1 \ll M_Z$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical
- X_μ becomes **dark photon** (often denoted as A'_μ) if $M_1 \ll M_Z$
- Dark photon A'_μ interactions: $\epsilon e Q_f A'_\mu \bar{f} \gamma^\mu f$ (visible) and $g_D A'_\mu \bar{\chi} \gamma^\mu \chi$ (invisible)

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi -\frac{\tilde{\epsilon}_1}{2}A_{\mu\nu}X^{\mu\nu} -\frac{M_1^2}{2}(\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical
- X_μ becomes **dark photon** (often denoted as A'_μ) if $M_1 \ll M_Z$
- Dark photon A'_μ interactions: $\epsilon e Q_f A'_\mu \bar{f} \gamma^\mu f$ (visible) and $g_D A'_\mu \bar{\chi} \gamma^\mu \chi$ (invisible)
- X_μ is hypercharge-like if $M_1 \gg M_Z$ (better probed at LHC)

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi -\frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} -\frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical
- X_μ becomes **dark photon** (often denoted as A'_μ) if $M_1 \ll M_Z$
- Dark photon A'_μ interactions: $\epsilon e Q_f A'_\mu \bar{f} \gamma^\mu f$ (visible) and $g_D A'_\mu \bar{\chi} \gamma^\mu \chi$ (invisible)
- X_μ is hypercharge-like if $M_1 \gg M_Z$ (better probed at LHC)
- If dark photon is massive, hidden fermion χ is **millicharged** only when $\tilde{\epsilon}_2 \neq 0$

[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon & millicharged particles

$$\mathcal{L} = -\frac{1}{4}A_{\mu\nu}A^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + g_D X_\mu \bar{\chi} \gamma^\mu \chi - \frac{\tilde{\epsilon}_1}{2} A_{\mu\nu} X^{\mu\nu} - \frac{M_1^2}{2} (\partial_\mu \sigma + X_\mu + \tilde{\epsilon}_2 A_\mu)^2$$

- kinetic mixing $\tilde{\epsilon}_1$ & mass mixing $\tilde{\epsilon}_2$ are degenerate; only $\epsilon \sim (\tilde{\epsilon}_2 - \tilde{\epsilon}_1)$ is physical
- X_μ becomes **dark photon** (often denoted as A'_μ) if $M_1 \ll M_Z$
- Dark photon A'_μ interactions: $\epsilon e Q_f A'_\mu \bar{f} \gamma^\mu f$ (visible) and $g_D A'_\mu \bar{\chi} \gamma^\mu \chi$ (invisible)
- X_μ is hypercharge-like if $M_1 \gg M_Z$ (better probed at LHC)
- If dark photon is massive, hidden fermion χ is **millicharged** only when $\tilde{\epsilon}_2 \neq 0$
- Millicharged particle χ interacts with photon via $\epsilon e A_\mu \bar{\chi} \gamma^\mu \chi$


[Feldman, ZL, Nath, hep-ph/0702123, 437 cites]

[see also Fabbrichesi+, 2005.01515, Dark Photon Review]

dark photon


Limits on dark photon with mass below 1 MeV

[<https://cajohare.github.io/AxionLimits/>]




ultralight dark photons
are DM candidates

Limits on dark photon with mass above 1 MeV



[Fabbrichesi+, 2005.01515, Dark Photon Review]

Limits on dark photon with mass above 1 MeV



probed by e^+e^- colliders


[Fabbrichesi+, 2005.01515, Dark Photon Review]

Dark photon production @ electron colliders


Dark photon A'_μ interactions: $\epsilon e Q_f A'_\mu \bar{f} \gamma^\mu f$ (visible) and $g_D A'_\mu \bar{\chi} \gamma^\mu \chi$ (invisible)

Production channel


(1) radiative return



(2) hadron decays



Dark photon visible decay branching ratio



only visible channels here

invisible ones can dominate
over the visible ones

[Mingxuan Du, ZL, Van Que Tran, 1912.00422]


Dark photon signatures @ electron colliders

Two different collider signatures for dark photon


visible decay ($ee/\mu\mu$)

(1) prompt



(2) long-lived



invisible decay: mono-photon




Belle II sensitivity on prompt dark photon **visible** decay




Belle-II, 2207.06307, Snowmass

Belle II sensitivity on long-lived dark photon




[Ferber, Garcia-Cely, Schmidt-Hoberg, 2202.03452]




photon + displaced vertex
e & non-e final states

Belle II sensitivity on dark photon invisible decay




Belle-II, 2207.06307, Snowmass



monochromatic photon
for on-shell dark photon

challenge: reducible BG
(see later slides)

BESIII limits on dark photon invisible decay




BESIII, 2209.13893

14.9/fb at 4.13-4.60 GeV


2

dark matter & millicharged particles


Mono-photon signature at electron colliders



Mono-photon signature at electron colliders



Mono-photon signature at electron colliders



probe a variety of models:

- EFT DM
- light mediator DM
- millicharged particles
(undetectable ionization)
- invisible dark photon

Different sub-GeV DM models (diff mediator mass)

EFT DM	$\frac{1}{\Lambda_V^2} \bar{\chi} \gamma_\mu \chi \bar{\ell} \gamma^\mu \ell$	$\frac{1}{\Lambda_A^2} \bar{\chi} \gamma_\mu \gamma_5 \chi \bar{\ell} \gamma^\mu \gamma_5 \ell$	$\frac{1}{\Lambda_S^2} \bar{\chi} \chi \bar{\ell} \ell$	$\frac{1}{\Lambda_t^2} \bar{\chi} \ell \bar{\ell} \chi$
--------	---	---	---	---

[ZL, Zhang, 1808.00983]

[Liang, ZL, Ma, Zhang, 1909.06847]

[Liang, ZL, Yang, 2111.15533]

Different sub-GeV DM models (diff mediator mass)

$$\text{EFT DM} \quad \frac{1}{\Lambda_V^2} \bar{\chi} \gamma_\mu \chi \bar{\ell} \gamma^\mu \ell \quad \frac{1}{\Lambda_A^2} \bar{\chi} \gamma_\mu \gamma_5 \chi \bar{\ell} \gamma^\mu \gamma_5 \ell \quad \frac{1}{\Lambda_S^2} \bar{\chi} \chi \bar{\ell} \ell \quad \frac{1}{\Lambda_t^2} \bar{\chi} \ell \bar{\ell} \chi$$

$$\text{Light mediator DM} \quad Z'_\mu \bar{\chi} \gamma^\mu (g_v^\chi - g_a^\chi \gamma_5) \chi + Z'_\mu \bar{\ell} \gamma^\mu (g_v^\ell - g_a^\ell \gamma_5) \ell$$

[ZL, Zhang, 1808.00983]

[Liang, ZL, Ma, Zhang, 1909.06847]

[Liang, ZL, Yang, 2111.15533]

Different sub-GeV DM models (diff mediator mass)

EFT DM

$$\frac{1}{\Lambda_V^2} \bar{\chi} \gamma_\mu \chi \bar{\ell} \gamma^\mu \ell \quad \frac{1}{\Lambda_A^2} \bar{\chi} \gamma_\mu \gamma_5 \chi \bar{\ell} \gamma^\mu \gamma_5 \ell \quad \frac{1}{\Lambda_S^2} \bar{\chi} \chi \bar{\ell} \ell \quad \frac{1}{\Lambda_t^2} \bar{\chi} \ell \bar{\ell} \chi$$

Light mediator DM

$$Z'_\mu \bar{\chi} \gamma^\mu (g_v^\chi - g_a^\chi \gamma_5) \chi + Z'_\mu \bar{\ell} \gamma^\mu (g_v^\ell - g_a^\ell \gamma_5) \ell$$

Millicharged particles

$$e e A_\mu \bar{\chi} \gamma^\mu \chi$$


[ZL, Zhang, 1808.00983]

[Liang, ZL, Ma, Zhang, 1909.06847]

[Liang, ZL, Yang, 2111.15533]


Irreducible background for mono-photon

Irreducible BG: $e^+e^- \rightarrow \gamma\nu\nu$



more challenging: reducible BG (due to limitations of the detectors)

Reducible background @ Belle II



Reducible background @ Belle II


ECL angles
(lab frame)

$$12.4^\circ < \theta < 31.4^\circ$$


$$32.2^\circ < \theta < 128.7^\circ$$

$$130.7^\circ < \theta < 155.1^\circ$$

[Belle II 1808.10567]



Reducible background @ Belle II



ECL angles
(lab frame)


$$12.4^\circ < \theta < 31.4^\circ$$

$$32.2^\circ < \theta < 128.7^\circ$$

$$130.7^\circ < \theta < 155.1^\circ$$

[Belle II 1808.10567]

Reducible background @ Belle II



ECL angles
(lab frame)

$$12.4^\circ < \theta < 31.4^\circ$$

$$32.2^\circ < \theta < 128.7^\circ$$

$$130.7^\circ < \theta < 155.1^\circ$$

[Belle II 1808.10567]


reducible BG

BG due to ECL gaps (gBG)

BG due to beam (bBG)

B: Invisible Dark Photon searches



[from Ferber's talk]




$ee \rightarrow 2\gamma$ and 3γ
 1 γ in ECL 90° gap
 1 γ out of ECL acceptance

$$E_y = \frac{s - M_{A'}^2}{2\sqrt{s}}$$

$ee \rightarrow eey$
 both electrons
 out of tracking acceptance




$ee \rightarrow 2\gamma$
 1 γ in ECL BWD or FWD gap



$ee \rightarrow 3\gamma$
 1 γ in ECL BWD gap
 1 γ out of ECL acceptance

B: Invisible Dark Photon searches



[from Ferber's talk]




$ee \rightarrow 2\gamma$ and 3γ
 1 γ in ECL 90° gap
 1 γ out of ECL acceptance

$$E_y = \frac{s - M_{A'}^2}{2\sqrt{s}}$$

$ee \rightarrow eey$
 both electrons
 out of tracking acceptance



$ee \rightarrow 2\gamma$
 1 γ in ECL BWD or FWD gap




$ee \rightarrow 3\gamma$
 1 γ in ECL BWD gap
 1 γ out of ECL acceptance


EFT DM probed by Belle II (50/ab)

[Liang, ZL, Yang, 2111.15533]

LEP is better than Belle II for EFT-DM





Electron colliders are better than DMDD



$$\frac{1}{\Lambda_V^2} \bar{\chi} \gamma_\mu \chi \bar{\ell} \gamma^\mu \ell$$

Light mediator DM probed by Belle II (50/ab) [2111.15533]

Belle II is better than LEP for light mediator DM Electron colliders can be better than DMDD




$$Z'_\mu \bar{\chi} \gamma^\mu (g_v^\chi - g_a^\chi \gamma_5) \chi + Z'_\mu \bar{\ell} \gamma^\mu (g_v^\ell - g_a^\ell \gamma_5) \ell$$

$$M_{Z'} = 5 \text{ GeV}, 0.6 \text{ GeV}, 10 \text{ MeV}$$

PandaX-4T limits on light DM


Search for Light Dark Matter with 259 Days of Data in PandaX-4T

[PandaX-4T, 2507.11930]



Belle II vs dark matter indirect detection experiments

Electron colliders can be better than DMID




Photons from DM annihilation

INTEGRAL

Fermi-LAT


millicharged particles

Millicharge in BSM can be quite “large”



[Jaeckel, Ringwald, 1002.0329]


Millicharge in BSM can be quite “large”




← probed by e^+e^- colliders

[Jaeckel, Ringwald, 1002.0329]

Millicharged particles probed by electron colliders



mono-photon




electron collider can probe
new parameter space

[ZL, Zhang, 1808.00983]

[Liang, ZL, Ma, Zhang, 1909.06847]

Run 3 milliQan bar detector




[2506.02251]

Low colliding energy is better to probe MCPs

if irreducible BG is the only important BG

BESIII/STCF vs Belle II




$$e^+e^- \rightarrow \gamma \bar{\nu} \nu$$

$$e^+e^- \rightarrow \gamma \bar{\chi} \chi$$

**new dark matter detection channel
at electron colliders**


Previous dark matter detection channels at colliders

Most studies focus on mono-X channel with SM X produced at the primary vertex




Previous dark matter detection channels at colliders

Most studies focus on mono-X channel with SM X produced at the primary vertex




New dark matter channel at colliders



A pair of SM particles
produced at the primary vertex


New dark matter channel at colliders



One SM particle interacts with the detector to produce a pair of DM particles

A pair of SM particles produced at the primary vertex

New dark matter channel at colliders




One SM particle interacts with the detector to produce a pair of DM particles

A pair of SM particles produced at the primary vertex

fixed target in collider


New dark matter channel @ electron colliders



[Liang, ZL, Yang, 2212.04252]

- $e^+e^- \rightarrow e^+e^-$
- e^- deposit energy in ECL
- e^+ interact with ECL to produce DM
- signature: e^- , e^+ , and missing energy (<5% e^+ energy deposited in ECL)

New dark matter channel @ electron colliders




[Liang, ZL, Yang, 2212.04252]


- $e^+e^- \rightarrow e^+e^-$
- e^- deposit energy in ECL
- e^+ interact with ECL to produce DM
- signature: e^- , e^+ , and missing energy (<5% e^+ energy deposited in ECL)

disappearing positron track

Positron interaction with ECL



annihilation w/
atomic electron



bremsstrahlung w/
target nucleus

[Liang, ZL, Yang, 2212.04252]


Background estimation

BG: $e^+ + \text{ECL} \rightarrow \gamma/n$ which escape detection

Use KLM to veto such BG


- photon BG events: ~ 13
- neutron BG events: ~ 81

[Liang, ZL, Yang, 2212.04252]





Belle II sensitivity on invisible dark photon

[Liang, ZL, Yang, 2212.04252]



Belle II sensitivity with ISR included

[Shao-Feng Ge, Jinhua Liang, ZL, Ui Min, 2505.10302]



3


Leptophilic gauge boson

Anomaly free extension to the standard model

Gauged lepton flavor $L_i - L_j$ are anomaly free extensions of the standard model


[Xiao-Gang He, Girish C. Joshi, H. Lew, R.R. Volkas, Phys.Rev.D 44 (1991) 2118-2132]

signature @ electron colliders




Belle II constraints on fully-invisible $L_\mu - L_\tau$ boson Z'

$\mu\mu + \text{miss}$



Belle-II, 2212.03066

Belle II constraints on vanilla $L_\mu - L_\tau$ boson Z'



Belle-II, 2212.03066

$\mu\mu + \text{miss}$

vanilla: only couple to SM fermions

invisible decay:

- Belle II
- NA64-e

decay to muons:

- Belle
- BaBar
- CMS

4

axion & axion-like particles

Axion & axion-like particles

Axion is proposed as a solution to the strong CP problem (QCD axion)

[Peccei & Quinn 1977]


Axion-like particles (ALPs) appears in many beyond-the-standard-model theories.

[see e.g. Ringwald 1210.5081]

$$\mathcal{L}_{\text{int}} = -\frac{1}{4} g_{a\gamma\gamma} F^{\mu\nu} \tilde{F}_{\mu\nu} a = g_{a\gamma\gamma} \mathbf{E} \cdot \mathbf{B} a$$

axion-photon couplings



Constraints on axion-photon coupling



[2305.01715, FIPs 2022 Report]


Belle II limits on ALPs

$$\mathcal{L}_{\text{int}} = -\frac{1}{4} g_{a\gamma\gamma} F^{\mu\nu} \tilde{F}_{\mu\nu} a$$



Belle-II, 2208.01101


Belle II sensitivity on ALP



Belle-II, 2207.06307

BESIII limits on ALPs

$$(2.71 \pm 0.01) \times 10^9 \psi(3683) \rightarrow \pi^+ \pi^- J/\psi, J/\psi \rightarrow \gamma a, a \rightarrow \gamma\gamma$$



BESIII, 2211.12699

**lepton flavor violating axions/
axion-like particles**

Lepton flavor violating axion-like particles

Lepton flavor violating axion-like particles

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{\partial_\mu a}{2f_a} \bar{\ell}_i C_{\ell_i \ell_i}^A \gamma^\mu \gamma_5 \ell_i + \sum_{i \neq j} \frac{\partial_\mu a}{2f_a} \bar{\ell}_i \gamma^\mu (C_{\ell_i \ell_j}^V + C_{\ell_i \ell_j}^A \gamma_5) \ell_j,$$

[Motoi Endo, Syuhei Iguro, Teppei Kitahara, 2002.05948]




[Lorenzo Calibbi, Diego Redigolo, Robert Ziegler, Jure Zupan, 2006.04795]

[Hooman Davoudiasl, Roman Marcarelli, Ethan T. Neil, 2112.04513]

[Yonglin Li, Zuowei Liu, 2501.12075]

$e\mu$ flavor violating axion-like particles @ Belle II

$$(y_A)_{e\mu} a \bar{e} \gamma_5 \mu$$




same sign & same-flavor lepton pairs via on-shell ALP: $e^+e^- \rightarrow \mu^\pm e^\mp a \rightarrow \mu^\pm \mu^\pm e^\mp e^\mp$

[Motoi Endo, Syuhei Iguro, Teppei Kitahara, 2002.05948]

$e\mu$ flavor violating axion-like particles @ Belle II


[Motoi Endo, Syuhei Iguro, Teppei Kitahara, 2002.05948]



$$(y_A)_{e\mu} a \bar{e} \gamma_5 \mu$$

Supernova constraints on $e\mu$ flavor violating ALPs

constraints from supernova cooling & low-energy supernova explosion energy



$$\mathcal{L}_{\text{int}} = -ig_{ae\mu} a \bar{e} \gamma_5 \mu + \text{h.c.}$$

$$e + \mu \rightarrow a$$

probes new para space for SN

[Li, ZL, 2501.12075]

[Huang, ZL, 2506.16922]

[Huang, Li, ZL, 2510.22523]


Summary

There are a variety of well-motivated new light particles that can be searched for at electron-positron colliders:

- dark photon
- millicharged particles
- dark matter
- axion and ALPS
- leptophilic boson

backup slides


BESIII limits on prompt dark photon **visible** decay



BESIII, 1705.04265

2.93/fb at 3.773 GeV

SENSEI constraints on millicharged particles



SENSEI

[2305.04964]