

Observation for TeV candidate of HMXB XTE J1858+034

Presenter: Xinpeng Cui

School: Guizhou Normal University

Collaborator: Shaoqiang Xi Xiaoli Huang

XTE J1858+034 is located at the center of the TeV source LHAASO J1858+0330. The source LHAASO J1858+0330 exhibits an extended structure of 0.52° .




Figure 1: The left panel shows the TS map within an 8.8° multiplied by 8° region centered on LHAASO J1858+0330 (R.A. = 284.71°, Decl. = 3.55°), while the right panel displays the TS map after excluding all other sources except this one.

Table 1: Morphological Fit Results for J1858+0330

Energy	Spatial Model	TS	R.A. deg	Decl. deg	Major Axis deg	Minor Axis deg	Position Angle	AIC
> 2TeV	Point	264.4	284.5815	3.6055				0
	Gaussian	574.0	284.6902	3.5778	0.5199±0.024			-299.084
	Elliptical gaussian	577.3	284.6905	3.5794	0.5336±0.033	0.4972±0.031	0	-295.722
	Disk	531.9	284.7250	3.5682	0.6551±0.016			-260.106

Note that TS differences cannot be used to quantitatively determine the best-fit model when models are non-nested. The Akaike information criterion (AIC; Akaike, 1974) was employed for model comparison. The AIC values are subtracted from that of the point model for a clear comparison. A comparison shows that the Gaussian model yields the smallest AIC value and is therefore adopted.




Figure 2: The left panel displays the TS map for WCDA ($2 \text{ TeV} < E < 25 \text{ TeV}$), while the right panel presents the TS map for KM2A ($25 \text{ TeV} < E < 100 \text{ TeV}$).




Figure 3: Energy spectrum of J1858+0330 observed by LHAASO.

Thank You!