

Studies of 1^{-+} states at BESIII


刘北江

中国科学院高能物理研究所

第十一届XYZ研讨会 2026.07.22-26, 贵阳

Mysterious gluons

- How does QCD give rise to hadrons?
 - What is the origin of confinement?
 - How is the mass generated in QCD?
- Role of gluons in hadrons
 - How to access soft gluon dynamics
- Key thing to search for: hadrons explicitly manifest the **gluonic degrees of freedom**
 - Gluonic excitations: **Blueballs and hybrids**, which provide critical information on the strong force and QCD vacuum



Eur.Phys.J.A 60 (2024) , 173

Light hadrons with exotic quantum numbers

- Finding unambiguous signature for exotics



- Efforts concentrate on Spin-exotic

- Forbidden for $q\bar{q}$:

$$J^{PC} = 0^{--}, \text{even}^{+-}, \text{odd}^{-+}$$

Experiments:

- Hadroproduction: GAMS, VES, E852, COMPASS, [AMBER](#)
- $p\bar{p}$ annihilation: Crystal Barrel, OBELIX, [PANDA](#)
- Photoproduction: [GlueX\(2017-\)](#), CLAS





$$\vec{J} = \vec{L} + \vec{S} \quad P = (-1)^{L+1} \quad C = (-1)^{L+S}$$

Allowed J^{PC} : $0^{-+}, 0^{++}, 1^{--}, 1^{+-}, 2^{++}, \dots$


Predictions

PRD 88 094505(2013)




“Constituent gluon”

- $(J^{PC})_g = 1^{+-}$
- $M_g \sim 1.3$ GeV



PRD 103, 054502(2021)




Decay width of 1^{-+} hybrid π_1

On lattice,


- Meson excitations similar to quark model
- Hybrid supermultiplet: 0^{-+} , 1^{--} , 2^{-+} , 1^{-+}
- Lightest spin-exotic state in LQCD: 1^{-+} hybrid

Spin-exotic mesons

- Candidates over 3 decades
 - $\pi_1(1400)$, $\pi_1(1600)$, $\pi_1(2015)$ (needs confirmation), all isovectors
 - Recent coupled-channel analyses suggest $\pi_1(1400)$ and $\pi_1(1600)$ as one pole



[PRL 122, 042002 (2019), EPJ C 81, 1056 (2021)]




Review: PRC 82, 025208 (2010), PPNP 82, 21 (2015), EPJC 83 (2023) 1125


	Decay mode	Reaction	Experiment
$\pi_1(1400)$	$\eta\pi$	$\pi^-p \rightarrow \pi^- \eta p$ $\pi^-p \rightarrow \pi^0 \eta n$ $\pi^-p \rightarrow \pi^- \eta p$ $\pi^-p \rightarrow \pi^0 \eta n$ $\bar{p}n \rightarrow \pi^- \pi^0 \eta$ $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$	GAMS KEK E852 E852 CBAR CBAR
	$\rho\pi$	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$	Obelix
$\pi_1(1600)$	$\eta'\pi$	$\pi^-Be \rightarrow \eta' \pi^- \pi^0 Be$ $\pi^-p \rightarrow \pi^- \eta' p$	VES E852
	$b_1\pi$	$\pi^-Be \rightarrow \omega \pi^- \pi^0 Be$ $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$ $\pi^-p \rightarrow \omega \pi^- \pi^0 p$	VES CBAR E852
	$\rho\pi$	$\pi^-Pb \rightarrow \pi^+ \pi^- \pi^- X$ $\pi^-p \rightarrow \pi^+ \pi^- \pi^- p$	COMPASS E852
	$f_1\pi$	$\pi^-p \rightarrow p \eta \pi^+ \pi^- \pi^-$ $\pi^-A \rightarrow \eta \pi^+ \pi^- \pi^- A$	E852 VES
$\pi_1(2015)$	$f_1\pi$	$\pi^-p \rightarrow \omega \pi^- \pi^0 p$	E852
	$b_1\pi$	$\pi^-p \rightarrow p \eta \pi^+ \pi^- \pi^-$	

Recent progress on $\pi_1(1600)$ from COMPASS

Spin-exotic $J^{PC}=1^{-+}$ waves at COMPASS
preliminary



Double Regge processes in $\pi\eta^{(\prime)}$



- Significant contribution of exotic π_1 trajectory: 9.9σ ($\eta\pi^-$) 5.5σ ($\eta'\pi^-$)


[D. Spuelbeck MESON2026]

Explore $\pi_1(1600)$ in photoproduction

- $\pi_1(1600) \rightarrow \eta\pi$ expected to be very small
- $\pi_1(1600) \rightarrow \eta'\pi$ potentially dominating the spectrum




Phys. Rev. Lett. 133, 261903(2024)




Studies of π_1 in $\chi_{c1} \rightarrow \pi^+ \pi^- \eta^{(\prime)}$

PR D84 112009 (2011)

2.6×10^7 $\psi(3686)$ @CLEO – c




No evidence of
 $\pi_1 \rightarrow \eta\pi$




Evidence of $\pi_1 \rightarrow \eta'\pi$
(without significant
BW phase motion)

PR D95 032002(2017)

44.8×10^7 $\psi(3686)$ @BESIII



Observation of π_1 in $\chi_{c1} \rightarrow \pi^+ \pi^- \eta'$



- $\pi_1(1600)$ observed $> 21\sigma$
- with a significant BW phase motion $> 11\sigma$
- $J^{PC} = 1^{-+}$, better than other assignments $> 17\sigma$

BW with mass-dependent width

$$M = (1828 \pm 8^{+11}_{-33}) \text{ MeV}/c^2,$$

$$\Gamma = (638 \pm 26^{+35}_{-86}) \text{ MeV}/c^2$$

$$B[\chi_{c1} \rightarrow \pi_1(1600)^\pm \pi^\mp] \times B[\pi_1(1600)^\pm \rightarrow \pi^\pm \eta'] = (4.30 \pm 0.14^{+1.04}_{-1.03}) \times 10^{-4}$$

pole: $(1690 \pm 16^{+36}_{-44}) - i(217 \pm 5^{+7}_{-19}) \text{ MeV}$

BW with constant width

$$M = (1675 \pm 9^{+32}_{-32}) \text{ MeV}/c^2,$$

$$\Gamma = (443 \pm 15^{+18}_{-56}) \text{ MeV}/c^2$$

$$B[\chi_{c1} \rightarrow \pi_1(1600)^\pm \pi^\mp] \times B[\pi_1(1600)^\pm \rightarrow \pi^\pm \eta'] = (4.27 \pm 0.14^{+0.86}_{-1.32}) \times 10^{-4}$$

pole: $(1689 \pm 10^{+33}_{-35}) - i(220 \pm 7^{+9}_{-27}) \text{ MeV}$



- Pole position is consistent with CLEO-c, COMPASS and JPAC

1^{-+} Hybrids

- **Isoscalar 1^{-+}** is critical to establish the hybrid nonet
 - Can be produced in the gluon-rich charmonium decays
 - Can decay to $\eta\eta'$ in P-wave

PRD 83,014021 (2011), PRD 83,014006 (2011), EPJ P135, 945(2020)

→ Search for $\eta_1 (1^{-+})$ in $J/\psi \rightarrow \gamma\eta\eta'$





$$\Gamma(J/\psi \rightarrow \gamma H) \sim O(\alpha\alpha_s^3)$$

Observation of An Exotic 1^{-+} Isoscalar State $\eta_1(1855)$

PRL 129 192002(2022) , PRD 106 072012(2022)

- An **isoscalar 1^{-+} , $\eta_1(1855)$** , has been **observed in $J/\psi \rightarrow \gamma\eta\eta'$ ($>19\sigma$)**
 $M = (1855 \pm 9_{-1}^{+6}) \text{ MeV}/c^2$, $\Gamma = (188 \pm 18_{-8}^{+3}) \text{ MeV}/c^2$
 $B(J/\psi \rightarrow \gamma\eta_1(1855) \rightarrow \gamma\eta\eta') = (2.70 \pm 0.41_{-0.35}^{+0.16}) \times 10^{-6}$
 - **Mass consistent with hybrid on LQCD**
- $\eta\eta'$ in P-waves uniquely indicates **1^{-+} exotic quantum numbers**




Observation of An Exotic 1^-+ Isoscalar State $\eta_1(1855)$

PRL 129 192002(2022) , PRD 106 072012(2022)

- Inspired many interpretations:
Hybrid/ $K\bar{K}_1$ Molecule/Tetraquark?




NPA 1047 122874(2024); Rept.Prog.Phys. 86 (2023) 026201;
PRD 107 (2023) 7, 074028; SCPMA 65 (2022) 6, 261011;
CPC 46 , 051001(2022); CPL 39, 051201 (2022);
PLB 834, 137478(2022); PRD 106 , 074003(2022); PRD 106,
036005(2022) ;...

- Opens a new direction to completing the picture of spin-exotics
 - As a “recent achievements and highlights” in hadron spectroscopy in the NuPECC LRP
 - 50 years of QCD: Exotic mesons, “observation of an $\eta_1(1855)$ state could be a breakthrough” [EPJ.C 83 (2023) 1125]



PWA results of $J/\psi \rightarrow \omega\eta\eta'$

BESIII preliminary




OZI filter to access quark contents

For comparison,

$$\begin{aligned}
 & B(J/\psi \rightarrow \gamma \eta_1(1855)) \\
 & \cdot B(\eta_1(1855) \rightarrow \eta \eta') \\
 & = (2.70 \pm 0.41^{+0.16}_{-0.35}) \times 10^{-6}
 \end{aligned}$$


$J/\psi \rightarrow \gamma\gamma\phi$, a $s\bar{s}$ flavor filter

PR D97 051101 (2018)



Amplitude analysis with
ML techniques for
background subtraction

← Fit to mass spectrum



From the amplitude analysis,

- $\eta(1405)$ is observed, while $\eta(1475)$ can not be excluded
- $X(1835) \rightarrow \gamma\phi$ suggests its assignment of η' excitation
- $\eta_c \rightarrow \gamma\phi$ are observed. The very first radiative decay mode of η_c
- Observation of $f_2(1950)$ and $f_0(2200) \rightarrow \gamma\phi$

unfavored their glueball interpretations [PRD 108, 014023, Sci.China Phys.Mech.Astron. 67 (2024) 11, 111012]


- **No evidence of $X(2370)/\eta_1(1855)$, well consistent with the predictions for glueball/hybrid**
[PRD 107, 114020, NPA 1037, 122683]

BESIII PhysRevD.111.052011(2025)



$J/\psi \rightarrow \gamma\gamma\omega$, a $n\bar{n}$ flavor filter

BESIII Preliminary




Resonance	[this analysis] $J/\psi \rightarrow \gamma R \rightarrow \gamma\gamma\omega (\times 10^{-6})$	[PRD 111, 052011 (2025)] $J/\psi \rightarrow \gamma R \rightarrow \gamma\gamma\phi (\times 10^{-6})$	$\frac{\Gamma(R \rightarrow \gamma\omega)}{\Gamma(R \rightarrow \gamma\phi)}$
$f_1(1285)$	$1.73 \pm 0.12^{+0.13}_{-0.18}$	$0.29 \pm 0.03^{+0.11}_{-0.09}$	3.5 ± 1.5
$f_1(1420)$	$0.88 \pm 0.10^{+0.20}_{-0.49}$	$0.55 \pm 0.07^{+0.18}_{-0.17}$	1.1 ± 0.7
$\eta(1405)$	$3.54 \pm 0.20^{+0.65}_{-0.24}$	$3.57 \pm 0.18^{+0.59}_{-0.61}$	0.3 ± 0.1
η_c	$0.20 \pm 0.06^{+0.03}_{-0.09}$	$0.21 \pm 0.03^{+0.05}_{-0.07}$	0.8 ± 0.5
$\eta(1295)$	< 0.35	< 0.84	
$\eta(1475)$	< 0.14	< 0.38	
$X(1835)$	< 0.11	$3.37 \pm 0.19^{+0.78}_{-1.10}$	
$X(2370)$	< 0.04	< 0.11	
$\eta_1(1855)$	< 0.33	< 4.74	

* $B(J/\psi \rightarrow \gamma X(2370))$ is estimated to be greater than 1×10^{-3}


The suppression of $X(2370)/\eta_1(1855)$ indicate gluonic content

Search for η_1 in $\chi_{c1} \rightarrow \eta\eta\eta'$


BESIII CPC 49, 10 (2025)




(a)




(b)






(a)



(b)



(c)




(f)

$$\mathcal{B}(\chi_{c1} \rightarrow \eta\eta\eta') = (1.39 \pm 0.13(\text{stat.}) \pm 0.09(\text{sys.})) \times 10^{-4}$$

$$\mathcal{B}(\chi_{c1} \rightarrow \eta_1(1855)\eta) \cdot \mathcal{B}(\eta_1(1855) \rightarrow \eta\eta') < 9.79 \times 10^{-5}.$$

$\bar{c}c g$ hybrids

LQCD predicts similar supermultiplet as light hybrids







JHEP 07 (2012), 126

- Y(4230)?
 - Seem to be contradicting with the exp. observation:
 $\sigma(D^* \bar{D} + c.c.) \gg \sigma(D^* \bar{D}^*)$
- Transitions between 1^{--} and $\{0, 1, 2\}^{-+}$
- Molecule states of 1^{-+} and 1^{--} ?

Further studied at
BESIII and Belle II


Search for 1^{-+} charmonium-like hybrid via $e^+e^- \rightarrow \gamma\eta^{(\prime)}\eta_c$ at center-of-mass energies between 4.258 and 4.681 GeV

BESIII PR D 111, 112007 (2025)






Search for a 1^{-+} molecular state via $e^+e^- \rightarrow \gamma D_s^+ D_{s1}^-(2536) + c.c.$

BESIII PRD 112, 032002(2025)



1^{-+} Hybrids

- How to discriminate hybrid/molecule/tetraquark?
 - $\eta_1(1855) \rightarrow f_1\eta, K_1K$ are important
 - $\Gamma_{\pi_1 \rightarrow b_1\pi} / \Gamma_{\pi_1 \rightarrow \eta'\pi}$ is not available yet
- Where is the $\eta_1^{(\prime)}$?
- Why η' is the observation channel for $\pi_1(1600)$ and $\eta_1(1855)$? $U_1(A)$ anomaly?
- Does K_1 exist and how to identify it?
- Where are the other $J^{PC} = (0^{--},)\text{even}^{+-}, \text{odd}^{-+}$ states? Analog in $\bar{c}c$ exists?
- New results via various probes are eagerly awaited



Thank you

backup

Observation of **An Exotic $1^- +$ Isoscalar State $\eta_1(1855)$**

PRL 129 192002(2022) , PRD 106 072012(2022)

- Angular distribution as a function of $M(\eta\eta')$ expressed **model-independently**

$$\langle Y_l^0 \rangle \equiv \sum_{i=1}^{N_k} W_i Y_l^0(\cos\theta_\eta^i)$$

- Related to the spin-0(S), spin-1(P), spin-2(D) amplitudes in $\eta\eta'$ by:

$$\sqrt{4\pi}\langle Y_0^0 \rangle = S_0^2 + P_0^2 + P_1^2 + D_0^2 + D_1^2 + D_2^2,$$

$$\sqrt{4\pi}\langle Y_1^0 \rangle = 2S_0P_0\cos\phi_{P_0} + \frac{2}{\sqrt{5}}(2P_0D_0\cos(\phi_{P_0} - \phi_{D_0}) + \sqrt{3}P_1D_1\cos(\phi_{P_1} - \phi_{D_1})),$$

$$\sqrt{4\pi}\langle Y_2^0 \rangle = \frac{1}{7\sqrt{5}}(14P_0^2 - 7P_1^2 + 10D_0^2 + 5D_1^2 - 10D_2^2) + 2S_0D_0\cos\phi_{D_0},$$


$$\sqrt{4\pi}\langle Y_3^0 \rangle = \frac{6}{\sqrt{35}}(\sqrt{3}P_0D_0\cos(\phi_{P_0} - \phi_{D_0}) - P_1D_1\cos(\phi_{P_1} - \phi_{D_1})),$$

$$\sqrt{4\pi}\langle Y_4^0 \rangle = \frac{1}{7}(6D_0^2 - 4D_1^2 + D_2^2).$$


- Narrow structure** in $\langle Y_1^0 \rangle$

➤ **Cannot be described by resonances in $\gamma\eta(\eta')$**

- $\eta_1(1855) \rightarrow \eta\eta'$ needed**



What about K_1 ?



Exotic Kaons are difficult to be established,
because they do not have exotic quantum numbers


Decay property is the key

- $K_1(1270)$
- $K_1(1400)$
- $K^*(1410)$
- $K_0^*(1430)$
- $K_2^*(1430)$
- $K(1460)$
- $K_2(1580)$
- $K(1630)$
- $K_1(1650)$
- $K^*(1680)$
- $K_2(1770)$
- $K_3^*(1780)$
- $K_2(1820)$
- $K(1830)$
- $K_0^*(1950)$
- $K_2^*(1980)$

$$\chi_{c1} \rightarrow K^+ K^- \eta'$$

PWA with 106 million $\psi(3686)$

PhysRevD.89.074030



Now, BESIII has 2.7 billion $\psi(3686)$. An updated analysis is underworking