

Introduction to Neutrino Oscillations

Shun Zhou
(IHEP, Beijing)

A. History of Neutrino Oscillations


B. Neutrino Oscillations in Vacuum

C. Neutrino Oscillations in Matter

The First China-Europe Neutrino Astronomy Summer School

September 14-23, 2026

Part A: History of Neutrino Oscillations






mass →	$\approx 2.3 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 173.07 \text{ GeV}/c^2$	0	$\approx 126 \text{ GeV}/c^2$
charge →	$2/3$	$2/3$	$2/3$	0	0
spin →	$1/2$	$1/2$	$1/2$	1	0
	u up	c charm	t top	g gluon	H Higgs boson
	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-1/3$	$-1/3$	$-1/3$	0	
	$1/2$	$1/2$	$1/2$	1	
	d down	s strange	b bottom	γ photon	
	$0.511 \text{ MeV}/c^2$	$105.7 \text{ MeV}/c^2$	$1.777 \text{ GeV}/c^2$	$91.2 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$1/2$	$1/2$	$1/2$	1	
	e electron	μ muon	τ tau	Z Z boson	
	$< 2.2 \text{ eV}/c^2$	$< 0.17 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$	$80.4 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$1/2$	$1/2$	$1/2$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	

Standard Model of Elementary Particles

- Three generations of quarks and leptons-building blocks of matter
- Gauge bosons as force carriers
 - Strong Interaction (8 gluons)
 - Electromagnetic Interaction (γ)
 - Weak Interaction (W and Z)
 - Gravitation (Graviton?)
- Neutrinos are massive particles-beyond the Standard Model

Pauli's Neutrino Hypothesis



Niels Bohr
(1885-1962)
Nobel Prize 1922


Energy not
conserved in
the quantum
domain?

Wolfgang Pauli
(1900–1958)
Nobel Prize 1945

there exist in the
nuclei electrically
neutral particles,
that I call neutrons



Detecting Neutrinos from Nuclear Explosions?



Bethe and Peierls (1934)


$$\bar{\nu}_e + p \rightarrow n + e^+ \quad \text{Cross section } \sigma < 10^{-44} \text{ cm}^2$$

“It is therefore absolutely impossible to observe the processes of this kind with the neutrinos created in nuclear transformations.”




Hans A. Bethe
(1906–2005)
Nobel Prize 1967

The combination of four protons and two electrons can occur essentially only in two ways. The first mechanism starts with the combination of two protons to form a deuteron with positron emission, *viz.*



The deuteron is then transformed into He^4 by further capture of protons; these captures occur very rapidly compared with process (1). The second mechanism uses carbon and nitrogen as catalysts, according to the chain reaction




No neutrinos in Bethe’s classic paper on nuclear reactions in stars (1939)




Reines to Fermi (1951): neutrino detector near ‘A-bombs’?

Discovery of Reactor Neutrinos 1954-1956



Gave up
'A-bombs'

Hanford
Nuclear
Reactor







three Gammas


5 μ s delay of γ
from n capture
on Cadmium

time coincidence


Discovery of Solar Neutrinos (1968)





Super-Kamiokande Neutrino Detector (since 1996)



Cherenkov Detectors



Super-Kamiokande : Image of the Sun in Neutrino Light



Neutrino Sources in Nature




Particle Accelerators


Nuclear Reactors




Cosmic rays
@ the Earth
Atmosphere




Natural radioactivity
in the Earth Crust





Nuclear Fusion
in the Sun



Supernova Explosion
SN 1987A




Cosmic Accelerators ?
PeV neutrino events
@ IceCube




Big Bang
(Indirect
Evidence)


Discovery of Neutrino Oscillations



L/E Significance



Angular Distribution of Events




↑ ~13000 km

← ~500 km

↓ ~15 km




Super-Kamiokande 1998, 2005

Discovery of Neutrino Oscillations



Super-Kamiokande Experiment

$$\nu_{\alpha} + e^{-} \rightarrow \nu_{\alpha} + e^{-}$$





SNO Experiment





$$\text{CC: } \nu_e + d \rightarrow p + p + e^{-}$$

$$\text{NC: } \nu_{\alpha} + d \rightarrow p + n + \nu_{\alpha}$$

$$\text{ES: } \nu_{\alpha} + e^{-} \rightarrow \nu_{\alpha} + e^{-}$$




Solar Neutrino Experiments






Reactor Neutrino Experiments



$$\bar{\nu}_e + p \rightarrow e^+ + n$$



REACTOR EXPERIMENTS




Accelerator Neutrino Experiments



From KEK to Super-Kamiokande

$E \sim 1.3 \text{ GeV}$, $L \sim 250 \text{ km}$





Accelerator Neutrino Experiments




MINOS Experiment

$E \sim 3 \text{ GeV}$, $L \sim 735 \text{ km}$



Story of the Smallest Mixing Angle

Global Analysis of Neutrino Oscillation Experiments



Schwetz et al., NJP 2008

No direct evidence for a nonzero θ_{13}

parameter	best fit	2σ	3σ
Δm_{21}^2 [10^{-5} eV^2]	$7.65^{+0.23}_{-0.20}$	7.25–8.11	7.05–8.34
$ \Delta m_{31}^2 $ [10^{-3} eV^2]	$2.40^{+0.12}_{-0.11}$	2.18–2.64	2.07–2.75
$\sin^2 \theta_{12}$	$0.304^{+0.022}_{-0.016}$	0.27–0.35	0.25–0.37
$\sin^2 \theta_{23}$	$0.50^{+0.07}_{-0.06}$	0.39–0.63	0.36–0.67
$\sin^2 \theta_{13}$	$0.01^{+0.016}_{-0.011}$	≤ 0.040	≤ 0.056


$\theta_{12}=33.5^\circ$, $\theta_{23}=45^\circ$, $\theta_{13}=5.7^\circ$

0.9 σ

Story of the Smallest Mixing Angle


June 2011 breakthrough: Appearance results from T2K and MINOS

T2K





2.5 σ

MINOS



1.7 σ

Story of the Smallest Mixing Angle



Dec. 2011

Double Chooz (far detector):

$$\sin^2 \theta_{13} = 0.022 \pm 0.013$$

1.7 σ

Mar. 2012

Daya Bay (near + far detectors):

$$\sin^2 \theta_{13} = 0.024 \pm 0.004$$

5.2 σ


Apr. 2012

RENO (near + far detectors):

$$\sin^2 \theta_{13} = 0.029 \pm 0.006$$

4.9 σ

Neutrino Oscillations



Pontecorvo-Maki-Nakagawa-Sakata Matrix

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$


Standard Parametrization

3-flavor
Mixing

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_{23} & \sin \theta_{23} \\ 0 & -\sin \theta_{23} & \cos \theta_{23} \end{pmatrix} \begin{pmatrix} \cos \theta_{13} & 0 & \sin \theta_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -\sin \theta_{13} e^{i\delta} & 0 & \cos \theta_{13} \end{pmatrix} \begin{pmatrix} \cos \theta_{12} & \sin \theta_{12} & 0 \\ -\sin \theta_{12} & \cos \theta_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Atmospheric Accelerator $\theta_{23} \sim 45^\circ$
 Reactor Accelerator $\theta_{13} \sim 9^\circ$
 Solar Reactor $\theta_{12} \sim 34^\circ$
 $0\nu\beta\beta?$


Neutrino Mass Ordering: JUNO



© JUNO, JPG, 2016



- ◆ 20 kton LS detector
- ◆ $3\%/\sqrt{E}$ resolution



Oscillation Probability:

$$P_{ee} = 1 - \sin^2 2\theta_{12} \cos^4 \theta_{13} \sin^2 \Delta_{21} - \frac{1}{2} \sin^2 2\theta_{13} [1 - \cos \Delta_* \cos \Delta_{21} + \cos 2\theta_{12} \sin \Delta_* \sin \Delta_{21}]$$

$$\Delta_* = \Delta_{31} + \Delta_{32} \quad \Delta_{ji} = \frac{(m_j^2 - m_i^2)L}{4E} \quad \begin{matrix} m_3 > m_2 > m_1 \\ m_2 > m_1 > m_3 \end{matrix}$$


Cores	YJ-C1	YJ-C2	YJ-C3	YJ-C4	YJ-C5	YJ-C6
Power (GW)	2.9	2.9	2.9	2.9	2.9	2.9
Baseline(km)	52.75	52.84	52.42	52.51	52.12	52.21
Cores	TS-C1	TS-C2	TS-C3	TS-C4	DYB	HZ
Power (GW)	4.6	4.6	4.6	4.6	17.4	17.4
Baseline(km)	52.76	52.63	52.32	52.20	215	265

JUNO: Jiangmen, Guangdong

3σ ~4σ in 6 years

Li et al., 1303.6733


Current Status of Absolute Neutrino Masses



$m_1 < m_2 < m_3$ (NO) or $m_3 < m_1 < m_2$ (IO)


Constraints on absolute neutrino masses

- Tritium β decays (95% C.L.)
 $m_\beta < 0.45$ eV (KATRIN 2025)
- Neutrinoless double- β decays (90% C.L.)
 $m_{\beta\beta} < (0.028 \sim 0.122)$ eV (KamLAND-Zen)
 $(0.078 \sim 0.239)$ eV (EXO-200)
 $(0.079 \sim 0.180)$ eV (GERDA-II)
 $(0.075 \sim 0.150)$ eV (CUORE)
- Cosmological observations (95% probability)
 $\Sigma < 0.12$ eV (Planck)



Part B: Neutrino Oscillation in Vacuum


Neutrino Production



Neutrino (flavor) oscillations:

well-established quantum phenomena of massive neutrinos at a macroscopic distance (from 1 to 1000 km or more)

Neutrino Propagation





Neutrino Detection

History of Neutrino Oscillations

23


- 1930** neutrino hypothesis by Pauli
- 1934** an effective theory for weak interactions by Fermi
- 1956** electron antineutrinos $\bar{\nu}_e$ discovered by Cowan and Reines
- 1957** neutrino-antineutrino transitions proposed by Pontecorvo
- 1962** The 2nd family of neutrinos $\nu_\mu/\bar{\nu}_\mu$ by Lederman, Schwarz & Steinberger
- 1962** neutrino flavor transitions proposed by Maki, Nakagawa & Sakata
- 1968** solar neutrinos ν_e detected by Davis & solar neutrino problem
- 1998** deficit of atmospheric neutrinos $\nu_\mu/\bar{\nu}_\mu$ in Super-Kamiokande (**Kajita**)
- 2000** The 3rd family of neutrinos $\nu_\tau/\bar{\nu}_\tau$ discovered in DONUT
- 2001** solar ν_e and ν_μ/ν_τ neutrinos found in SNO (**McDonald**)
- 2002** KamLAND selects the LMA-MSW solution to solar neutrino problem
- 2004** K2K & MINOS confirms the disappearance of atmospheric neutrinos
- 2012** reactor antineutrino disappearance discovered by Daya Bay





Neutrino **flavor eigenstates** are defined with respect to corresponding charged leptons, and they may not coincide with **mass eigenstates**

$$\begin{aligned}\nu_e &= +\cos\theta \nu_1 + \sin\theta \nu_2 \\ \nu_\mu &= -\sin\theta \nu_1 + \cos\theta \nu_2\end{aligned}$$



$$\begin{aligned}\nu_1 &= +\cos\theta \nu_e - \sin\theta \nu_\mu \\ \nu_2 &= +\sin\theta \nu_e + \cos\theta \nu_\mu\end{aligned}$$

The oscillation probability in the plane-wave approximation: **simple & useful**

$$|\nu_e(0)\rangle = |\nu_e\rangle = \cos\theta |\nu_1\rangle + \sin\theta |\nu_2\rangle$$

$$|\nu_e(t)\rangle = \cos\theta e^{-iE_1 t} |\nu_1\rangle + \sin\theta e^{-iE_2 t} |\nu_2\rangle$$

$$= e^{-iE_1 t} (\cos\theta |\nu_1\rangle + \sin\theta e^{-i\Delta E t} |\nu_2\rangle)$$

$$\begin{aligned}\Delta E &= E_2 - E_1 = \sqrt{p^2 + m_2^2} - \sqrt{p^2 + m_1^2} \\ &\approx \left(p + \frac{m_2^2}{2p}\right) - \left(p + \frac{m_1^2}{2p}\right) \equiv \frac{\Delta m^2}{2E}\end{aligned}$$

$$\Delta m^2 \equiv m_2^2 - m_1^2 \quad E \simeq p \gg m_1, m_2$$

The transitional amplitude

$$A = \langle \nu_\mu | \nu_e(t) \rangle = e^{-iE_1 t} \sin\theta \cos\theta (e^{-i\Delta E t} - 1) \quad |\nu_\mu\rangle = -\sin\theta |\nu_1\rangle + \cos\theta |\nu_2\rangle$$

The oscillation probability for the **appearance** channel


$$P(\nu_e \rightarrow \nu_\mu) \equiv |\langle \nu_\mu | \nu_e(t) \rangle|^2 = |\sin \theta \cos \theta (e^{-i\Delta Et} - 1)|^2$$

$$= 2 \sin^2 \theta \cos^2 \theta \left(1 - \cos \frac{\Delta m^2 t}{2E} \right)$$

$$= \sin^2 2\theta \sin^2 \frac{\Delta m^2 L}{4E}$$

$L \approx t$ for neutrinos in the relativistic limit

The survival probability for the **disappearance** channel



Natural units: $\hbar = c = 1$

Natural units

Realistic units

Phase factors

$$\exp(-iE_{1,2}t)$$

$$\exp\left(-i\frac{E_{1,2}}{\hbar}t\right)$$

Energies and momentum

$$E_{1,2} = \sqrt{p^2 + m_{1,2}^2}$$

$$E_{1,2} = \sqrt{p^2c^2 + m_{1,2}^2c^4}$$

Energy difference

$$\Delta E = \frac{\Delta m^2}{2E}$$

$$\Delta E = \frac{\Delta m^2c^3}{2p} = \frac{\Delta m^2c^4}{2E}$$

Time and distance

$$t = L$$

$$t = \frac{L}{c}$$


Oscillation argument

$$\frac{1}{2}\Delta Et = \frac{\Delta m^2L}{4E}$$

$$\frac{1}{2}\frac{\Delta E}{\hbar}t = \frac{c^3}{\hbar} \cdot \frac{\Delta m^2L}{4E}$$

- (1) Derive the factor 1.27 in the formula of oscillation probabilities
- (2) Daya Bay experiment is designed to observe the disappearance of reactor neutrinos, so it is better to put the far detector at the oscillation maximum for $\Delta m^2 = 2.5 \times 10^{-3} \text{ eV}^2$. Take $E = 4 \text{ MeV}$ & estimate the optimal baseline length for Daya Bay.

Neutrino Production



Charged-current Weak Interaction


$$\mathcal{L}_{\text{cc}} = \frac{g}{\sqrt{2}} (\bar{e} \quad \bar{\mu} \quad \bar{\tau})_L \gamma^{\mu} \mathbf{V} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}_L W_{\mu}^{-} + \text{h.c.}$$

Flavor Mixing Matrix
or the PMNS Matrix

$$|\nu_{\alpha}(0)\rangle = |\nu_{\alpha}\rangle = \sum_i V_{\alpha i}^* |\nu_i\rangle$$

$$|\nu_{\alpha}(t)\rangle = \sum_i V_{\alpha i}^* e^{-iE_i t} |\nu_i\rangle \quad |\nu_{\beta}\rangle = \sum_j V_{\beta j}^* |\nu_j\rangle$$

Neutrino Propagation



Neutrino Detection

$$A(\nu_{\alpha} \rightarrow \nu_{\beta}) = \langle \nu_{\beta} | \nu_{\alpha}(t) \rangle = \sum_i V_{\alpha i}^* V_{\beta i} e^{-iE_i t}$$

$$P(\nu_{\alpha} \rightarrow \nu_{\beta}) = |\langle \nu_{\beta} | \nu_{\alpha}(t) \rangle|^2 = \left| \sum_i V_{\alpha i}^* V_{\beta i} e^{-iE_i t} \right|^2$$

Derivation of the general formula of oscillation probabilities

Fritzsch & Minkowski, 76



$$\begin{aligned}
 P(\nu_\alpha \rightarrow \nu_\beta) &= |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_i V_{\alpha i}^* V_{\beta i} e^{-iE_i t} \right|^2 \\
 &= \sum_{i=1}^3 |V_{\alpha i}^* V_{\beta i}|^2 + 2 \sum_{i < j} \text{Re} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* e^{i(E_j - E_i)t} \right] \\
 &= \sum_{i=1}^3 |V_{\alpha i}^* V_{\beta i}|^2 + 2 \sum_{i < j} \text{Re} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* \right] \cos \frac{\Delta m_{ji}^2 L}{2E} \\
 &\quad - 2 \sum_{i < j} \text{Im} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* \right] \sin \frac{\Delta m_{ji}^2 L}{2E} \\
 &= \boxed{\sum_{i=1}^3 |V_{\alpha i}^* V_{\beta i}|^2 + 2 \sum_{i < j} \text{Re} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* \right]} \longrightarrow \left| \sum_{i=1}^3 V_{\alpha i}^* V_{\beta i} \right|^2 = \delta_{\alpha\beta} \\
 &\quad - 4 \sum_{i < j} \text{Re} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* \right] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} - 2 \sum_{i < j} \text{Im} \left[V_{\alpha i}^* V_{\beta i} V_{\alpha j} V_{\beta j}^* \right] \sin \frac{\Delta m_{ji}^2 L}{2E}
 \end{aligned}$$

$$\begin{aligned}
 \Delta m_{ji}^2 &\equiv m_j^2 - m_i^2 \\
 E &\simeq p \gg m_i, m_j
 \end{aligned}$$

The general formula of oscillation probabilities



Jarlskog Invariant

$$\text{Im} \left[V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^* \right]$$

$$\equiv \mathcal{J} \sum_{\gamma, k} \varepsilon_{\alpha \beta \gamma} \varepsilon_{i j k}$$

$$\begin{aligned} P(\nu_{\alpha} \rightarrow \nu_{\beta}) &= \delta_{\alpha \beta} - 4 \sum_{i < j}^3 \text{Re} \left[V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^* \right] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} \\ &\quad + 2 \sum_{i < j}^3 \text{Im} \left[V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^* \right] \sin \frac{\Delta m_{ji}^2 L}{2E} \\ &= \delta_{\alpha \beta} - 4 \sum_{i < j}^3 \text{Re} \left[V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^* \right] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} \end{aligned}$$

$$+ 8 \mathcal{J} \sum_{\gamma} \varepsilon_{\alpha \beta \gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$

CP Violation

Exercise: try to demonstrate that the imaginary part can be expressed in terms of the Jarlskog invariant and derive the CP-violating term

$$+ 2 \sum_{i < j}^3 \text{Im} \left[V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^* \right] \sin \frac{\Delta m_{ji}^2 L}{2E}$$

$$= + 2J \sum_{\gamma} \varepsilon_{\alpha\beta\gamma} \left(\sin \frac{\Delta m_{21}^2 L}{2E} - \sin \frac{\Delta m_{31}^2 L}{2E} + \sin \frac{\Delta m_{32}^2 L}{2E} \right)$$

$$= - 2J \sum_{\gamma} \varepsilon_{\alpha\beta\gamma} \left(\sin \frac{\Delta m_{12}^2 L}{2E} + \sin \frac{\Delta m_{31}^2 L}{2E} + \sin \frac{\Delta m_{23}^2 L}{2E} \right)$$

$$= + 8J \sum_{\gamma} \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{12}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E} \sin \frac{\Delta m_{23}^2 L}{4E}$$



Jarlskog Invariant

$$\begin{pmatrix} V_{e1} & V_{e2} & V_{e3} \\ V_{\mu1} & V_{\mu2} & V_{\mu3} \\ V_{\tau1} & V_{\tau2} & V_{\tau3} \end{pmatrix}$$

$$J = \text{Im} [V_{e1} V_{\mu2} V_{e2}^* V_{\mu1}^*]$$

Exercise: J is unique

Unitarity triangle and Jarlskog invariant



$$a = |V_{e1} V_{\mu1}^*| e^{i\alpha}$$

$$b = |V_{e2} V_{\mu2}^*| e^{i\beta}$$

$$c = |V_{e3} V_{\mu3}^*| e^{i\gamma}$$

$$V_{e1} V_{\mu1}^* + V_{e2} V_{\mu2}^* + V_{e3} V_{\mu3}^* = 0 \quad \text{Exercise: prove } |J| = 2 \times \text{area of triangle}$$

$$\mathcal{L}_{\text{cc}} = \frac{g}{\sqrt{2}} (\bar{e} \quad \bar{\mu} \quad \bar{\tau})_L \gamma^\mu \mathbf{V} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}_L W_\mu^- + \text{h.c.}$$


$$|\bar{\nu}_\alpha\rangle = \sum_i V_{\alpha i} |\bar{\nu}_i\rangle$$

$$\mathbf{V} \Rightarrow \mathbf{V}^*$$

The oscillation probabilities for antineutrinos

$$P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) = \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \text{Re} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin^2 \frac{\Delta m_{ji}^2 L}{4E}$$

$$- 8J \sum_\gamma \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$



$$J \Rightarrow -J$$

CP violation for neutrino oscillations in vacuum

$$A_{\text{CP}} = P(\nu_\alpha \rightarrow \nu_\beta) - P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) = 16J \sum_\gamma \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$

No CP asymmetries in the **disappearance** channels (via **CPT theorem**)

Basic formula:

$$P(v_\alpha \rightarrow v_\beta) = \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \operatorname{Re} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} + 8J \sum_{\gamma}^3 \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$

CP transformation:

$$V \Rightarrow V^*$$

$$J \Rightarrow -J$$

$$P(\bar{v}_\alpha \rightarrow \bar{v}_\beta) = \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \operatorname{Re} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} - 8J \sum_{\gamma}^3 \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$

T transformation:


$$\alpha \Leftrightarrow \beta$$

$$J \Rightarrow -J$$

$$P(v_\beta \rightarrow v_\alpha) = \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \operatorname{Re} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} - 8J \sum_{\gamma}^3 \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta m_{21}^2 L}{4E} \sin \frac{\Delta m_{32}^2 L}{4E} \sin \frac{\Delta m_{31}^2 L}{4E}$$

CPT theorem: $P(v_\alpha \rightarrow v_\beta) = P(\bar{v}_\beta \rightarrow \bar{v}_\alpha)$ **Invalid if matter effects exist**

Neutrino Production



- Neutrino-antineutrino oscillations can take place only if neutrinos are Majorana particles, as the lepton number is violated in this case.

Quantization of neutrino fields:

$$\nu(x) = \int \frac{d^3p}{(2\pi)^3 2E} \sum_{h=\pm} \left[a_h(p) u_h(p) e^{-ipx} + \underbrace{a_h^{\dagger}(p)}_{\text{creation operator}} v_h(p) e^{ipx} \right]$$

$$\begin{aligned} |v(p)\rangle &= |v(p, -)\rangle \\ |\bar{v}(p)\rangle &= |v(p, +)\rangle \end{aligned}$$

$$\nu^c(x) = \nu(x)$$


creation operator
for both ν and $\bar{\nu}$

Neutrino Propagation

$$A(\nu_{\alpha} \rightarrow \bar{\nu}_{\beta}) = \sum_i V_{\alpha i}^* V_{\beta i}^* \frac{m_i}{E_i} e^{-iE_i t}$$

$$A(\bar{\nu}_{\alpha} \rightarrow \nu_{\beta}) = \sum_i V_{\alpha i} V_{\beta i} \frac{m_i}{E_i} e^{+iE_i t}$$

Highly suppressed by a factor of m_i/E



Neutrino Detection

Oscillation probabilities

Pontecorvo, 57, 67; Bahcall & Primakoff, 78;
Schechter & Valle, 81; ...

$$P(\nu_\alpha \rightarrow \bar{\nu}_\beta) = \frac{1}{E^2} \left[|\langle m \rangle_{\alpha\beta}|^2 - 4 \sum_{i < j}^3 m_i m_j \operatorname{Re} \left[\underline{V_{\alpha i} V_{\beta i} V_{\alpha j}^* V_{\beta j}^*} \right] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} \right. \\ \left. + 2 \sum_{i < j}^3 m_i m_j \operatorname{Im} \left[\underline{V_{\alpha i} V_{\beta i} V_{\alpha j}^* V_{\beta j}^*} \right] \sin \frac{\Delta m_{ji}^2 L}{2E} \right]$$

$$P(\bar{\nu}_\alpha \rightarrow \nu_\beta) = \frac{1}{E^2} \left[|\langle m \rangle_{\alpha\beta}|^2 - 4 \sum_{i < j}^3 m_i m_j \operatorname{Re} \left[V_{\alpha i} V_{\beta i} V_{\alpha j}^* V_{\beta j}^* \right] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} \right. \\ \left. - 2 \sum_{i < j}^3 m_i m_j \operatorname{Im} \left[V_{\alpha i} V_{\beta i} V_{\alpha j}^* V_{\beta j}^* \right] \sin \frac{\Delta m_{ji}^2 L}{2E} \right]$$

$$10^{-12} \left(\frac{m_i}{\text{eV}} \right)^2 \left(\frac{\text{MeV}}{E} \right)^2$$

$$\langle m \rangle_{\alpha\beta} \equiv \sum_{i=1}^3 V_{\alpha i} V_{\alpha i} m_i$$

Dependence on all three
CP-violating phases

The oscillation probabilities are extremely small, but enhanced for $\bar{\nu}_e$ from ^3H decays.

Dirac neutrinos

$$c_{ij} \equiv \cos \theta_{ij}, \quad s_{ij} \equiv \sin \theta_{ij}$$

$$V = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \cdot \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \cdot \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Majorana neutrinos

$$V = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \cdot \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \cdot \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Exercise: prove that a general 2 x 2 unitary matrix contains four real parameters, and can be written as follows

$$V = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{i\beta} \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \cdot \begin{pmatrix} e^{i\gamma} & 0 \\ 0 & 1 \end{pmatrix}$$

then all the phases can be removed by redefining the charged-lepton and neutrino fields in the Dirac case. So, no CP violation if there are 2 generations of fermions. Consider also the Majorana case.

Standard Parametrization

$$V = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & +c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & c_{13}s_{23} \\ +s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{13}c_{23} \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Extraction of mixing angles and CP-violating phases from the PMNS matrix

$$\sin^2 \theta_{13} = |V_{e3}|^2$$

$$\mathcal{J} = \text{Im}[V_{e2}V_{\mu 3}V_{e3}^*V_{\mu 2}^*]$$

$$\sin^2 \theta_{12} = \frac{|V_{e2}|^2}{1 - |V_{e3}|^2}$$

$$= \text{Im}[(c_{13}s_{12}) \cdot (c_{13}s_{23}) \cdot (s_{13}e^{i\delta}) \cdot (+c_{12}c_{23})]$$

$$= s_{12}c_{12}s_{23}c_{23}s_{13}c_{13}^2 \sin \delta$$

$$\sin^2 \theta_{23} = \frac{|V_{\mu 3}|^2}{1 - |V_{e3}|^2}$$

$$\sin \delta = \frac{\text{Im}[V_{e2}V_{\mu 3}V_{e3}^*V_{\mu 2}^*]}{s_{12}c_{12}s_{23}c_{23}s_{13}c_{13}^2}$$



**three mixing angles
are restricted to the
first quadrant**

Exercise: prove that a general 3 x 3 unitary matrix contains nine real parameters, and there will be one non-factorizable phase.

Most oscillation experiments are measuring the disappearance of neutrinos

$$P(\nu_\alpha \rightarrow \nu_\beta) = \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \operatorname{Re} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin^2 \frac{\Delta m_{ji}^2 L}{4E} + 2 \sum_{i < j}^3 \operatorname{Im} [V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*] \sin \frac{\Delta m_{ji}^2 L}{2E}$$

$$\operatorname{Re} [V_{\alpha i} V_{\alpha j} V_{\alpha j}^* V_{\alpha i}^*] = |V_{\alpha i}|^2 |V_{\alpha j}|^2$$

$$\operatorname{Im} [V_{\alpha i} V_{\alpha j} V_{\alpha j}^* V_{\alpha i}^*] = 0$$

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{i < j}^3 |V_{\alpha i}|^2 |V_{\alpha j}|^2 \sin^2 \frac{\Delta m_{ji}^2 L}{4E}$$

$$= 1 - 4 |V_{\alpha 1}|^2 |V_{\alpha 2}|^2 \sin^2 \frac{\Delta m_{21}^2 L}{4E} - 4 |V_{\alpha 1}|^2 |V_{\alpha 3}|^2 \sin^2 \frac{\Delta m_{31}^2 L}{4E} - 4 |V_{\alpha 2}|^2 |V_{\alpha 3}|^2 \sin^2 \frac{\Delta m_{32}^2 L}{4E}$$

Comments:

- No CP asymmetry, so valid for both ν and $\bar{\nu}$ oscillations
- In a specific experiment with L and E fixed, so sensitive to a particular range of Δm_{ji}^2
- 2-flavor oscillations if θ_{13} is small and $|\Delta m_{21}^2| \ll |\Delta m_{31}^2| \simeq |\Delta m_{32}^2|$

One can determine the **mixing angles** and neutrino **mass-squared differences**

$$P(\nu_\mu \rightarrow \nu_e) = \sin^2 \theta_{23} \sin^2 2\theta_{13} \sin^2 \frac{\Delta m_{31}^2 L}{4E}$$

+ **CPV** + **Matter Effects**

Comments:

- Sensitive to θ_{13} , depending on other mixing parameters

For accelerator neutrinos (T2K)

$$E \simeq 0.6 \text{ GeV} \quad L \simeq 295 \text{ km}$$



$$\frac{L}{E} \simeq 500 \text{ m/MeV}$$



$$\Delta m^2 \simeq 2.5 \times 10^{-3} \text{ eV}^2$$

3-flavor results without matter effects

$$P(\nu_\mu \rightarrow \nu_e) = \underbrace{(\sin^2 \theta_{23} \sin^2 2\theta_{13} - 8J \sin \frac{\Delta m_{21}^2 L}{2E}) \sin^2 \frac{\Delta m_{31}^2 L}{4E}}_{\text{leading terms}} + \underbrace{c_{13}^2 \left[\sin^2 2\theta_{12} (c_{23}^2 - s_{23}^2 s_{13}^2) + \frac{1}{4} \sin 4\theta_{12} \sin 2\theta_{23} \cos \delta \right] \sin^2 \frac{\Delta m_{21}^2 L}{4E}}_{\text{sub-leading terms}}$$

Main features of LBL accelerator neutrino experiments: sensitive to δ , octant of θ_{23} , and also sign of Δm_{31}^2 when including matter effects


Neutrino oscillations in matter

L. Wolfenstein


Carnegie-Mellon University, Pittsburgh, Pennsylvania 15213

(Received 6 October 1977; revised manuscript received 5 December 1977)

The effect of coherent forward scattering must be taken into account when considering the oscillations of neutrinos traveling through matter. In particular, for the case of massless neutrinos for which vacuum oscillations cannot occur, oscillations can occur in matter if the neutral current has an off-diagonal piece connecting different neutrino types. Applications discussed are solar neutrinos and a proposed experiment involving transmission of neutrinos through 1000 km of rock.



Lincoln Wolfenstein
(1923-2015)




Refraction of light in media

$$\begin{aligned} \nu_e &: \exp[ipx(n_{nc} + n_{cc} - 1)] \\ \nu_\mu &: \exp[ipx(n_{nc} - 1)] \\ \nu_\tau &: \exp[ipx(n_{nc} - 1)] \end{aligned}$$

Refraction of neutrinos in media, where both CC and NC interactions contribute to refractive indices (not far from 1)

When neutrinos are traveling in matter, the effect of coherent forward scattering with background particles leads to a modification of their energies. Such a modification can be described by an potential energy. The difference between the potentials of distinct neutrino flavors is relevant for neutrino oscillations.

Ordinary matter contains only electrons, neutrons and protons:



Incoherent scattering

$$\ell_{\text{matter}} \sim \frac{10^{14} \text{ cm}}{(E/\text{GeV})}$$

We take the number density of particles in normal matter to be about 10^{24} cm^{-3} .

First of all, we look at the Hamiltonian of free neutrinos in vacuum

in mass basis

$$\begin{aligned} \hat{\mathcal{H}} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} &= \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} \\ &= \left[E \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2E} \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} \right] \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} \end{aligned}$$


in flavor basis

$$\begin{aligned} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} &= \begin{pmatrix} c_\theta & s_\theta \\ -s_\theta & c_\theta \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} \\ &\equiv U \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} \end{aligned}$$

$$\hat{\mathcal{H}} \begin{pmatrix} c_\theta & -s_\theta \\ s_\theta & c_\theta \end{pmatrix} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} \begin{pmatrix} c_\theta & -s_\theta \\ s_\theta & c_\theta \end{pmatrix} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} \rightarrow$$

$$\mathcal{H} = \frac{1}{2E} U \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} U^\dagger$$

Effective Hamiltonian density for weak interactions



$$\mathcal{H}_{cc}(x) = \frac{G_F}{\sqrt{2}} [\bar{\nu}_e(x) \gamma^\mu (1 - \gamma_5) e(x)] [\bar{e}(x) \gamma_\mu (1 - \gamma_5) \nu_e(x)]$$

Fierz transformation

$$\mathcal{H}_{cc}(x) = \frac{G_F}{\sqrt{2}} [\bar{\nu}_e(x) \gamma^\mu (1 - \gamma_5) \nu_e(x)] [\bar{e}(x) \gamma_\mu (1 - \gamma_5) e(x)]$$

Averaged Hamiltonian density over the electron background

$$\bar{\mathcal{H}}_{cc}(x) = \frac{G_F}{\sqrt{2}} \bar{\nu}_e(x) \gamma^\mu (1 - \gamma_5) \nu_e(x) \int d^3 p_e \mathbf{f}(\mathbf{p}_e) \text{ Distribution function of electrons}$$


Averaged over electron helicities $\times \frac{1}{2} \sum_{h_e=\pm} \langle e(p_e, h_e) | \bar{e}(x) \gamma_\mu (1 - \gamma_5) e(x) | e(p_e, h_e) \rangle$

$$|e(p_e, h_e)\rangle = \frac{1}{2EV} a_e^{(h_e)\dagger} |0\rangle$$

$$\int d^3 p_e f(p_e) = N_e V$$

$$\bar{\mathcal{H}}_{cc}(x) = \frac{G_F}{\sqrt{2}} \int d^3 p_e f(p_e) \bar{\nu}_e(x) \frac{\mathbf{p}_{e\mu}}{E_e V} \gamma^\mu (1 - \gamma_5) \nu_e(x)$$

$$\begin{aligned}
 \bar{\mathcal{H}}_{cc}(x) &= \frac{G_F}{\sqrt{2}} \int d^3 p_e f(p_e) \bar{\nu}_e(x) \frac{\mathbf{p}_{e\mu}}{E_e V} \boldsymbol{\gamma}^\mu (1 - \gamma_5) \nu_e(x) \\
 &= \frac{G_F}{\sqrt{2} V} \bar{\nu}_e(x) \int d^3 p_e f(p_e) \left(\boldsymbol{\gamma}^0 - \frac{\vec{\mathbf{p}}_e \cdot \vec{\boldsymbol{\gamma}}}{E_e} \right) (1 - \gamma_5) \nu_e(x) \\
 &= \frac{G_F}{\sqrt{2} V} \bar{\nu}_e(x) \boldsymbol{\gamma}^0 (1 - \gamma_5) \nu_e(x) N_e V \\
 &= \sqrt{2} G_F N_e \nu_{eL}^\dagger(x) \nu_{eL}(x)
 \end{aligned}$$



NC interactions contribute equally to all flavors, which is irrelevant for oscillations



$$V_e = \sqrt{2} G_F N_e$$

Matter Potential (minus sign for $\bar{\nu}$)

Then, we obtain the Hamiltonian for neutrinos travelling in matter

$$A = 2\sqrt{2} G_F N_e E$$

$$\mathcal{H}_m = \frac{1}{2E} U \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} U^\dagger + \begin{pmatrix} V_e & 0 \\ 0 & 0 \end{pmatrix} = \frac{1}{2E} \left[U \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} U^\dagger + \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix} \right]$$

Exercise: evaluate the electron number density in Earth matter of a mass density $\rho = 4 \text{ g/cm}^3$. Note that the normal matter is electrically neutral, so the number fraction of electrons is $Y_e = 0.5$.

The Hamiltonian in a more compact form:

$$\mathcal{H}_m = \frac{1}{4E} \left[U \begin{pmatrix} -\Delta m_{21}^2 & 0 \\ 0 & +\Delta m_{21}^2 \end{pmatrix} U^\dagger + \begin{pmatrix} A & 0 \\ 0 & -A \end{pmatrix} \right] + \frac{m_1^2 + m_2^2 + A}{4E}$$



$$\mathcal{H}_m = \frac{1}{4E} \begin{pmatrix} A - \Delta m_{21}^2 c_{2\theta} & \Delta m_{21}^2 s_{2\theta} \\ \Delta m_{21}^2 s_{2\theta} & \Delta m_{21}^2 c_{2\theta} - A \end{pmatrix} \quad \text{in flavor basis}$$

Converted into the mass basis

$$\mathcal{H}_m = \frac{1}{4E} \tilde{U} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & 0 \\ 0 & +\Delta \tilde{m}_{21}^2 \end{pmatrix} \tilde{U}^\dagger$$

Mixing matrix & mass states in matter

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} c_{\tilde{\theta}} & s_{\tilde{\theta}} \\ -s_{\tilde{\theta}} & c_{\tilde{\theta}} \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1\rangle \\ |\tilde{\nu}_2\rangle \end{pmatrix} \equiv \tilde{U} \begin{pmatrix} |\tilde{\nu}_1\rangle \\ |\tilde{\nu}_2\rangle \end{pmatrix}$$

$$\Delta \tilde{m}_{21}^2 = \sqrt{(\Delta m_{21}^2 c_{2\theta} - A)^2 + (\Delta m_{21}^2 s_{2\theta})^2} \quad \tan 2\tilde{\theta} = \frac{\Delta m_{21}^2 s_{2\theta}}{\Delta m_{21}^2 c_{2\theta} - A}$$

Relationship between the mixing angle (mass difference) in vacuum and that in matter

Mixing angle in matter

$$\tan 2\tilde{\theta} = \frac{\Delta m_{21}^2 s_{2\theta}}{\Delta m_{21}^2 c_{2\theta} - A}$$


MSW resonance $\tilde{\theta} = 45^\circ$

Resonance condition:

$$\Delta m_{21}^2 c_{2\theta} = 2\sqrt{2}G_F N_e E$$



Stanislav Mikheyev (1940-2011) Alexei Smirnov (1951-)



Lincoln Wolfenstein (1923-2015)

Exercise: assume that the energy of solar ${}^8\text{B}$ neutrinos is $E = 10 \text{ MeV}$, and take $N_e = 100 N_A/\text{cm}^3$ for $\rho = 150 \text{ g/cm}^3$ in the solar center. The density decreases from the center to the surface. Check if the MSW resonance can be reached, given $\Delta m_{21}^2 = 7.5 \times 10^{-5} \text{ eV}^2$.

Solution:

$$A = 2\sqrt{2}G_F N_e E \approx 2\sqrt{2} \cdot (1.17 \times 10^{-5} \text{ GeV}^{-2}) \cdot 10 \text{ MeV} \cdot (6 \times 10^{25} \text{ cm}^{-3}) \\ \approx 1.5 \times 10^{-4} \text{ eV}^2$$

$$V_e = \sqrt{2}G_F N_e \approx 7.5 \times 10^{-6} \text{ eV}^2/\text{MeV}$$

A useful relation: $197 \text{ MeV} \cdot 1 \text{ fm} = 1$

$$P(\nu_e \rightarrow \nu_\mu) = \sin^2 2\theta \sin^2 \frac{\Delta m^2 L}{4E} \\ \tilde{P}(\nu_e \rightarrow \nu_\mu) = \sin^2 2\tilde{\theta} \sin^2 \frac{\Delta \tilde{m}^2 L}{4E}$$

The matter density $\rho(r)$ is varying in astrophysical environments, like the Sun and SNe

$$\tan 2\tilde{\theta}(r) = \frac{\Delta m_{21}^2 s_{2\theta}}{\Delta m_{21}^2 c_{2\theta} - 2\sqrt{2}G_F N_e(r)E} \quad \mathcal{H}_m = \frac{1}{4E} \tilde{U} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & 0 \\ 0 & +\Delta \tilde{m}_{21}^2 \end{pmatrix} \tilde{U}^\dagger$$

Recall how to calculate neutrino oscillation probabilities in matter of **a constant density**

$$i \frac{d}{dr} \begin{pmatrix} |\nu_e(r)\rangle \\ |\nu_\mu(r)\rangle \end{pmatrix} = \frac{1}{4E} \tilde{U} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & 0 \\ 0 & +\Delta \tilde{m}_{21}^2 \end{pmatrix} \tilde{U}^\dagger \begin{pmatrix} |\nu_e(r)\rangle \\ |\nu_\mu(r)\rangle \end{pmatrix}$$

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \tilde{U} \begin{pmatrix} |\tilde{\nu}_1\rangle \\ |\tilde{\nu}_2\rangle \end{pmatrix}$$



$$i \frac{d}{dr} \left[\tilde{U} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix} \right] = \frac{1}{4E} \tilde{U} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & 0 \\ 0 & +\Delta \tilde{m}_{21}^2 \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix}$$



$$i \frac{d}{dr} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix} = \left[\frac{1}{4E} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & 0 \\ 0 & +\Delta \tilde{m}_{21}^2 \end{pmatrix} - i \tilde{U}^\dagger \dot{\tilde{U}} \right] \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix}$$

$$\tilde{U}^\dagger \dot{\tilde{U}} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{d\tilde{\theta}}{dr}$$

$$i \frac{d}{dr} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix} = \frac{1}{4E} \begin{pmatrix} -\Delta \tilde{m}_{21}^2 & -4iEd\tilde{\theta}/dr \\ 4iEd\tilde{\theta}/dr & +\Delta \tilde{m}_{21}^2 \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix}$$

If $\rho(r)$ or $\tilde{\theta}(r)$ changes slowly, no transition between mass states

Evolution of mass states in matter

$$i \frac{d}{dr} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix} = \frac{1}{4E} \begin{pmatrix} -\Delta\tilde{m}_{21}^2 & -4iEd\tilde{\theta}/dr \\ 4iEd\tilde{\theta}/dr & +\Delta\tilde{m}_{21}^2 \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1(r)\rangle \\ |\tilde{\nu}_2(r)\rangle \end{pmatrix}$$

Exercise: calculate $\frac{d\tilde{\theta}}{dr}$

Adiabaticity parameter: the ratio between the diagonal and off-diagonal elements

$$\gamma = \frac{\Delta\tilde{m}_{21}^2}{4E|d\tilde{\theta}/dr|} = \frac{(\Delta\tilde{m}_{21}^2)^2}{2E \sin 2\tilde{\theta} |dA/dr|}$$

Mass ordering: sign of Δm^2

$$\mathcal{H} = U(\theta) \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} U^\dagger(\theta)$$


invariant under

$$\theta \Rightarrow \frac{\pi}{2} - \theta \quad m_1^2 \Leftrightarrow m_2^2$$

$$|\nu_e\rangle \Rightarrow +|\nu_e\rangle \quad |\nu_\mu\rangle \Rightarrow -|\nu_\mu\rangle$$

$$\theta \in [0, \frac{\pi}{4}] \quad \Delta m_{21}^2 > 0$$

$$\theta \in [\frac{\pi}{4}, \frac{\pi}{2}] \quad \Delta m_{21}^2 < 0$$



The effective Hamiltonian in matter

$$\mathcal{H}_m = \frac{1}{2E} \left[V \begin{pmatrix} m_1^2 & 0 & 0 \\ 0 & m_2^2 & 0 \\ 0 & 0 & m_3^2 \end{pmatrix} V^\dagger + \begin{pmatrix} A & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right]$$


Diagonalize the effective Hamiltonian in matter

$$\tilde{V}^\dagger \mathcal{H}_m \tilde{V} = \frac{1}{2E} \begin{pmatrix} \tilde{m}_1^2 & 0 & 0 \\ 0 & \tilde{m}_2^2 & 0 \\ 0 & 0 & \tilde{m}_3^2 \end{pmatrix} \quad \text{Parametrize } \tilde{V} \text{ in the standard way by three mixing angles } \tilde{\theta}_{ij} \text{ and one CP-violating phase } \tilde{\delta} \text{ in matter}$$


Oscillation probabilities in matter of a constant density

$$\begin{aligned} \tilde{P}(v_\alpha \rightarrow v_\beta) = & \delta_{\alpha\beta} - 4 \sum_{i < j}^3 \text{Re} \left[\tilde{V}_{\alpha i} \tilde{V}_{\beta j} \tilde{V}_{\alpha j}^* \tilde{V}_{\beta i}^* \right] \sin^2 \frac{\Delta \tilde{m}_{ji}^2 L}{4E} \\ & + 8\tilde{J} \sum_{\gamma} \varepsilon_{\alpha\beta\gamma} \sin \frac{\Delta \tilde{m}_{21}^2 L}{4E} \sin \frac{\Delta \tilde{m}_{32}^2 L}{4E} \sin \frac{\Delta \tilde{m}_{31}^2 L}{4E} \end{aligned}$$


The oscillation probabilities for antineutrinos can be obtained by $V \Rightarrow V^*$ and $A \Rightarrow -A$




Total Rates: Standard Model vs. Experiment
Bahcall–Serenelli 2005 [BS05(OP)]



Solar ν_e converted to ν_μ or ν_τ





For high-energy ^8B neutrinos
at production $r = 0$

$$\begin{pmatrix} |\tilde{\nu}_1(0)\rangle \\ |\tilde{\nu}_2(0)\rangle \end{pmatrix} = \begin{pmatrix} c_{\hat{\theta}} & -s_{\hat{\theta}} \\ s_{\hat{\theta}} & c_{\hat{\theta}} \end{pmatrix} \begin{pmatrix} |\nu_e(0)\rangle \\ |\nu_{\mu}(0)\rangle \end{pmatrix}$$

adiabatic evolution

$$\begin{pmatrix} |\tilde{\nu}_1(R)\rangle \\ |\tilde{\nu}_2(R)\rangle \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1(0)\rangle \\ |\tilde{\nu}_2(0)\rangle \end{pmatrix}$$

on the solar surface $r = R$

$$\begin{pmatrix} |\nu_e(R)\rangle \\ |\nu_{\mu}(R)\rangle \end{pmatrix} = \begin{pmatrix} c_{\theta} & s_{\theta} \\ -s_{\theta} & c_{\theta} \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_1(R)\rangle \\ |\tilde{\nu}_2(R)\rangle \end{pmatrix}$$

survival probability

$$P_{ee} = c_{\hat{\theta}}^2 c_{\theta}^2 + s_{\hat{\theta}}^2 s_{\theta}^2 = \sin^2 \theta$$

$$\hat{\theta} \rightarrow \pi/2 \quad \text{as } A \gg \Delta m^2$$

For the MSW resonance to happen

$$\theta \in [0, \frac{\pi}{4}]$$

$$\Delta m_{21}^2 > 0$$



$$\theta \in [\frac{\pi}{4}, \frac{\pi}{2}]$$

$$\Delta m_{21}^2 < 0$$

$$\theta_{12} = 34^\circ$$

Normal neutrino
mass ordering

For low-energy ^7Be neutrinos

$$P_{ee} \approx 1 - \frac{1}{2} \sin^2 2\theta_{12}$$

Oscillations in vacuum

nu-fit.org

		Normal Ordering (best fit)		Inverted Ordering ($\Delta\chi^2 = 1.5$)	
		bfp $\pm 1\sigma$	3σ range	bfp $\pm 1\sigma$	3σ range
IC23 without SK atmospheric data	$\sin^2 \theta_{12}$	$0.3088^{+0.0067}_{-0.0066}$	$0.2893 \rightarrow 0.3295$	$0.3088^{+0.0067}_{-0.0066}$	$0.2893 \rightarrow 0.3295$
	$\theta_{12}/^\circ$	$33.76^{+0.42}_{-0.41}$	$32.54 \rightarrow 35.03$	$33.76^{+0.42}_{-0.41}$	$32.54 \rightarrow 35.03$
	$\sin^2 \theta_{23}$	$0.470^{+0.017}_{-0.014}$	$0.432 \rightarrow 0.587$	$0.555^{+0.013}_{-0.016}$	$0.437 \rightarrow 0.590$
	$\theta_{23}/^\circ$	$43.27^{+1.0}_{-0.82}$	$41.11 \rightarrow 50.02$	$48.15^{+0.75}_{-0.92}$	$41.40 \rightarrow 50.21$
	$\sin^2 \theta_{13}$	$0.02249^{+0.00057}_{-0.00057}$	$0.02070 \rightarrow 0.02420$	$0.02261^{+0.00056}_{-0.00056}$	$0.02091 \rightarrow 0.02433$
	$\theta_{13}/^\circ$	$8.62^{+0.11}_{-0.11}$	$8.27 \rightarrow 8.95$	$8.65^{+0.11}_{-0.11}$	$8.31 \rightarrow 8.97$
	$\delta_{\text{CP}}/^\circ$	207^{+23}_{-20}	$114 \rightarrow 405$	283^{+24}_{-28}	$202 \rightarrow 347$
	$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.537^{+0.094}_{-0.10}$	$7.236 \rightarrow 7.823$	$7.537^{+0.094}_{-0.10}$	$7.236 \rightarrow 7.822$
	$\frac{\Delta m_{3\ell}^2}{10^{-3} \text{ eV}^2}$	$+2.521^{+0.026}_{-0.018}$	$+2.454 \rightarrow +2.592$	$-2.500^{+0.024}_{-0.023}$	$-2.569 \rightarrow -2.430$

Neutrino Mass Ordering

- Reactor: **JUNO**
- LBL Acc.: T2K, NOvA, DUNE
- Atm: PINGU, ORCA, Hyper-K

Leptonic CP Violation

- LBL Acc.: **T2HK**, DUNE
- Super-B: ESSvSB, MOMENT
- NF & Beta-Beams

Absolute Masses: KATRIN, $0\nu\beta\beta$ (^{136}Xe , ^{76}Ge), cosmology, ...