

Further Discussions on Massive Neutrinos

Shun Zhou

(IHEP, Beijing)

2026/9/18

1 Quantum Mechanics of Neutrino Oscillations

- A Simple Derivation
- Wave-packet Formalism

2 Density Matrix Formalism

- General Comments
- Neutrino Oscillations in Vacuum
- Neutrino Oscillations in Matter

3 Massive Majorana Neutrinos

- Fermion Masses
- Majorana Fermions
- Right-handed Neutrinos in Seesaw

A Simple Derivation

Equal Momenta

As we have shown before, the neutrino oscillation probabilities can be computed as

$$P(\nu_\alpha \rightarrow \nu_\beta; t) = \left| \sum_i V_{\alpha i}^* e^{-iE_i t} V_{\beta i} \right|^2. \quad (1.1)$$

Then we take the following approximation ($E_i \approx p \gg m_i$)

$$E_i = \sqrt{p^2 + m_i^2} = p \sqrt{1 + m_i^2/p^2} \approx p + m_i^2/2p \quad (1.2)$$

and sometimes $E_i \approx p \approx E$ as well. The energy difference reads

$$E_i - E_j \approx (m_i^2 - m_j^2)/(2p) \equiv \Delta m_{ij}^2/(2p), \quad (1.3)$$

and finally the time-to-distance conversion $t \approx L$ is performed.

A Simple Derivation

Equal Energies

We assume that the energies of neutrino mass eigenstates are the same, and they are propagating in space, leading to the oscillation probabilities

$$P(\nu_\alpha \rightarrow \nu_\beta; x) = \left| \sum_i V_{\alpha i}^* e^{+ip_i x} V_{\beta i} \right|^2. \quad (1.4)$$

Then we take the following approximation ($p_i \approx E \gg m_i$)

$$p_i = \sqrt{E^2 - m_i^2} = E \sqrt{1 - m_i^2/E^2} \approx E - m_i^2/2E \quad (1.5)$$

and sometimes $p_i \approx p \approx E$ as well. The energy difference reads

$$p_i - p_j \approx -(m_i^2 - m_j^2)/(2E) \equiv -\Delta m_{ij}^2/(2E), \quad (1.6)$$

and no time-to-distance conversion is needed.

Equal Energies or Momenta

Which is Correct?

The answer is that both are wrong. Consider the production of neutrino mass eigenstates from $\pi^+ \rightarrow \mu^+ + \nu_i$. If the energies and momenta of π^+ and μ^+ are given, then we have

$$E_{\pi^+} = E_{\mu^+} + E_i, \quad \mathbf{p}_{\pi^+} = \mathbf{p}_{\mu^+} + \mathbf{p}_i, \quad (1.7)$$

where the energies and momenta of ν_i (for $i = 1, 2, 3$) fulfill the relation

$$E_i = \sqrt{\mathbf{p}_i^2 + m_i^2}. \quad (1.8)$$

Hence both energies and momenta of different neutrino mass eigenstates are different. In the rest frame of π^+ , we have $E_{\pi^+} = m_\pi$ and $\mathbf{p}_{\pi^+} = 0$:

$$|\mathbf{p}_i| = \sqrt{[(m_\pi - m_\mu)^2 - m_i^2] [(m_\pi + m_\mu)^2 - m_i^2]} / (2m_\pi). \quad (1.9)$$

Wave-packet Formalism

A Realistic Description

The correct description should be the wave-packet formalism:

$$|\nu_\alpha(t, \mathbf{x})\rangle = \sum_i V_{\alpha i}^* \psi_i(t, \mathbf{x}) |\nu_i\rangle, \quad (1.10)$$

where the wave packet for a free neutrino mass eigenstate is

$$\psi_i(t, \mathbf{x}) = \int \frac{d^3 p}{(2\pi)^{3/2}} f_i^S(\mathbf{p} - \mathbf{p}_i) e^{i[\mathbf{p} \cdot \mathbf{x} - E_i(p)t]}. \quad (1.11)$$

Some comments are helpful:

- $f_i^S(\mathbf{p} - \mathbf{p}_i)$ is the momentum distribution sharply peaked at $\mathbf{p}_i$ for the neutrino eigenstates at the source with the width $\sigma_p \ll p_i$.
- $E_i(p) = \sqrt{\mathbf{p}^2 + m_i^2}$ is the energy corresponding to the momentum $\mathbf{p}$.

A Realistic Description

Expand the energy $E_i(p)$ around the mean momentum p_i

$$\begin{aligned} E_i(p) = & E_i(p_i) + \left. \frac{\partial E_i(p)}{\partial p^j} \right|_{\mathbf{p}_i} (p - p_i)^j \\ & + \frac{1}{2} \left. \frac{\partial^2 E_i(p)}{\partial p^j \partial p^k} \right|_{\mathbf{p}_i} (p - p_i)^j (p - p_i)^k + \cdots \end{aligned} \quad (1.12)$$

We can rewrite $\psi_i(t, \mathbf{x})$ as below

$$\begin{aligned} \psi_i(t, \mathbf{x}) & \approx \int \frac{d^3 p}{(2\pi)^{3/2}} f_i^S(\mathbf{p} - \mathbf{p}_i) e^{i \left\{ \mathbf{p} \cdot \mathbf{x} - \left[E_i(p_i) + \left. \frac{\partial E_i(p)}{\partial p^j} \right|_{\mathbf{p}_i} (p - p_i)^j \right] t \right\}} \\ & = e^{i[\mathbf{p}_i \cdot \mathbf{x} - E_i(p_i)t]} \cdot g_i^S(\mathbf{x} - \mathbf{v}_{ig} t) . \end{aligned} \quad (1.13)$$

A Realistic Description

The shape factor is defined as

$$g_i^S(\mathbf{x} - \mathbf{v}_{ig}t) = \int \frac{d^3p}{(2\pi)^{3/2}} f_i^S(\mathbf{p}) e^{i[\mathbf{p} \cdot (\mathbf{x} - \mathbf{v}_{ig}t)]}, \quad (1.14)$$

where the group velocity of the wave packet is

$$\mathbf{v}_{ig} = \left. \frac{\partial E_i(p)}{\partial \mathbf{p}} \right|_{\mathbf{p}_i} = \frac{\mathbf{p}}{E_i(p)} \Big|_{\mathbf{p}_i}. \quad (1.15)$$

- The shape factor $g_i^S(\mathbf{x} - \mathbf{v}_g t)$ stands for the distribution of the wave packet in the coordinate space.
- The wave packet is travelling at the speed of $\mathbf{v}_{ig}$ without changing its shape. The width is $\sigma_x \gtrsim \sigma_p^{-1}$ according to the uncertainty principle.

Wave-packet Formalism

A Realistic Description

Next, we have to specify the neutrino state at the detector, which is taken to be another wave packet located at $\mathbf{x} = \mathbf{L}$, i.e.,

$$\begin{aligned} |\nu_\beta(\mathbf{x} - \mathbf{L})\rangle &= \sum_i V_{\beta i}^* \psi_i^D(\mathbf{x} - \mathbf{L}) |\nu_i\rangle \\ &= \sum_i V_{\beta i}^* e^{i\mathbf{p}_i \cdot (\mathbf{x} - \mathbf{L})} g_i^D(\mathbf{x} - \mathbf{L}) |\nu_i\rangle \end{aligned} \quad (1.16)$$

where g_i^D stands for the shape of the wave packet of $|\nu_i\rangle$ at the detector.

Oscillation Amplitudes

The amplitudes are given by

$$\mathcal{A}_{\alpha\beta}(t, \mathbf{L}) = \int d^3x \langle \nu_\beta(\mathbf{x} - \mathbf{L}) | \nu_\alpha(t, \mathbf{x}) \rangle . \quad (1.17)$$

Wave-packet Formalism

Oscillation Amplitudes

The amplitudes are given by

$$\begin{aligned}\mathcal{A}_{\alpha\beta}(t, \mathbf{L}) &= \int d^3x \langle \nu_\beta(\mathbf{x} - \mathbf{L}) | \nu_\alpha(t, \mathbf{x}) \rangle \\ &= \sum_i V_{\alpha i}^* V_{\beta i} G(\mathbf{L} - \mathbf{v}_{ig} t) e^{-iE_i(p_i)t + i\mathbf{p}_i \cdot \mathbf{L}},\end{aligned}\quad (1.18)$$

where

$$G(\mathbf{L} - \mathbf{v}_{ig} t) \equiv \int d^3x g_i^S(\mathbf{x} - \mathbf{v}_{ig} t) \cdot g_i^{D*}(\mathbf{x} - \mathbf{L}). \quad (1.19)$$

Note that $g_i^S(\mathbf{x} - \mathbf{v}_{ig} t)$ decreases rapidly when $\mathbf{x}$ deviates far from $\mathbf{v}_{ig} t$, so does $g_i^D(\mathbf{x} - \mathbf{L})$ for $\mathbf{x}$ from $\mathbf{L}$. Hence $G(\mathbf{L} - \mathbf{v}_{ig} t)$ peaks at $\mathbf{L} = \mathbf{v}_{ig} t$, i.e., the amplitudes are not small only when the wave packets overlap.

Density Matrix Formalism: General Comments

Pure States

Given a pure quantum state $|\psi(t)\rangle$, one can construct the density matrix operator via

$$\hat{\rho}(t) \equiv |\psi(t)\rangle\langle\psi(t)|, \quad (2.1)$$

where $\langle\psi(t)| \equiv |\psi(t)\rangle^\dagger$ has been defined such that the density matrix operator is Hermitian $\hat{\rho}^\dagger = \hat{\rho}$.

Matrix Representation

Expanding the state $|\psi(t)\rangle$ in terms of a complete set of orthonormal eigenstates $|\phi_i\rangle$ (for $i = 1, 2, \dots, n$) of the Hamiltonian, we have

$$|\psi(t)\rangle = \sum_{i=1}^n a_i(t) |\phi_i\rangle. \quad (2.2)$$

Density Matrix Formalism: General Comments

Matrix Representation

Then, the density matrix can be written as

$$\hat{\rho}(t) = |\psi(t)\rangle\langle\psi(t)| = \sum_{i,j=1}^n a_i(t)a_j^*(t)|\phi_i\rangle\langle\phi_j|, \quad (2.3)$$

and its element is given by

$$\hat{\rho}_{kl}(t) \equiv \langle\phi_k|\hat{\rho}(t)|\phi_l\rangle = \sum_{i,j=1}^n a_i(t)a_j^*(t)\langle\phi_k|\phi_i\rangle\langle\phi_j|\phi_l\rangle = a_k(t)a_l^*(t). \quad (2.4)$$

The trace of the density matrix for a pure state is

$$\text{Tr}[\hat{\rho}(t)] = \sum_{i=1}^n |a_i(t)|^2 = 1, \quad (2.5)$$

where the state $|\psi(t)\rangle$ is normalized to one.

Density Matrix Formalism: General Comments

Liouville-von Neumann Equation

Starting with the Schrödinger equation

$$i\frac{\partial}{\partial t}|\psi(t)\rangle = \hat{H}|\psi(t)\rangle, \quad (2.6)$$

and its Hermitian conjugate

$$-i\frac{\partial}{\partial t}\langle\psi(t)| = \langle\psi(t)|\hat{H}, \quad (2.7)$$

we can find the time evolution of the density matrix

$$i\frac{\partial\hat{\rho}(t)}{\partial t} = i\frac{\partial|\psi(t)\rangle}{\partial t}\langle\psi(t)| + i|\psi(t)\rangle\frac{\partial\langle\psi(t)|}{\partial t} = [\hat{H}, \hat{\rho}]. \quad (2.8)$$

Here $\hat{H}$ is the Hamiltonian operator. One can figure out the matrix representations of any operators in the chosen basis of state vectors.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum

We apply the density matrix formalism to two-flavor neutrino oscillations in vacuum, for which the Hamiltonian is given by

$$\begin{aligned}\hat{H} &= \frac{1}{2E} \begin{pmatrix} m_1^2 c_\theta^2 + m_2^2 s_\theta^2 & \Delta m_{21}^2 s_\theta c_\theta \\ \Delta m_{21}^2 s_\theta c_\theta & m_1^2 s_\theta^2 + m_2^2 c_\theta^2 \end{pmatrix} \\ &= \frac{m_1^2 + m_2^2}{4E} + \frac{\Delta m_{21}^2}{4E} \begin{pmatrix} -c_{2\theta} & s_{2\theta} \\ s_{2\theta} & +c_{2\theta} \end{pmatrix} \\ &\equiv \omega_0 \sigma^0 + \frac{1}{2} \omega \mathbf{B} \cdot \boldsymbol{\sigma},\end{aligned}\tag{2.9}$$

where $\omega \equiv \Delta m_{21}^2/(2E)$ is the oscillation frequency and

$$\sigma^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.\tag{2.10}$$

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum

The term $\omega_0 \sigma^0$ with $\omega_0 = (m_1^2 + m_2^2)/(4E)$ contributes a common phase factor and is irrelevant. The “magnetic field” vector can be identified as

$$\mathbf{B} = (B_1, B_2, B_3) \equiv (s_{2\theta}, 0, -c_{2\theta}) . \quad (2.11)$$

On the other hand, we can expand the density matrix in terms of Pauli matrices and obtain the polarization vector $\mathbf{P}(t)$:

$$\hat{\rho}(t) = P_0 \sigma^0 + \frac{1}{2} \mathbf{P} \cdot \boldsymbol{\sigma} . \quad (2.12)$$

Notice that the Hamiltonian $\hat{H}$ has been given in the flavor basis, for which the basis vectors are $|\nu_e\rangle$ and $|\nu_\mu\rangle$, namely,

$$\hat{H} = H_{\alpha\beta} |\nu_\alpha\rangle \langle \nu_\beta| , \quad (2.13)$$

for $\alpha, \beta = e, \mu$. What are $\hat{\rho}(0)$ and $\hat{\rho}(t)$ in the same basis?

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum

For a general state $|\psi(t)\rangle = a_e(t)|\nu_e\rangle + a_\mu(t)|\nu_\mu\rangle$ in the flavor basis, we can get the expression of the density matrix

$$\hat{\rho}(t) = \begin{pmatrix} |a_e(t)|^2 & a_e(t)a_\mu^*(t) \\ a_e^*(t)a_\mu(t) & |a_\mu(t)|^2 \end{pmatrix}, \quad (2.14)$$

and its value at $t = 0$

$$\hat{\rho}(0) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (2.15)$$

where $a_e(0) = 1$ and $a_\mu(0) = 0$ imply that the initial state is $|\psi(0)\rangle = |\nu_e\rangle$.

One can observe the physical meaning of diagonal elements of $\hat{\rho}(t)$, i.e., $\hat{\rho}_{ee}(t) = |a_e(t)|^2$ for probability in the state $|\nu_e\rangle$, and $\hat{\rho}_{\mu\mu}(t) = |a_\mu(t)|^2$ for probability in the state $|\nu_\mu\rangle$.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum

It is now straightforward to find out the polarization vector:

$$\begin{aligned}\hat{\rho}(t) &= \frac{|a_e(t)|^2 + |a_\mu(t)|^2}{2} + \frac{1}{2} \begin{pmatrix} |a_e(t)|^2 - |a_\mu(t)|^2 & 2a_e(t)a_\mu^*(t) \\ 2a_e^*(t)a_\mu(t) & |a_\mu(t)|^2 - |a_e(t)|^2 \end{pmatrix} \\ &\equiv P_0\sigma^0 + \frac{1}{2}\mathbf{P}(t) \cdot \boldsymbol{\sigma} ,\end{aligned}$$

where $|a_e(t)|^2 + |a_\mu(t)|^2 = 1$ holds, i.e., $P_0 = 1/2$ and

$$P_1 = 2\text{Re} [a_e^*(t)a_\mu(t)] , P_2 = 2\text{Im} [a_e^*(t)a_\mu(t)] , P_3 = |a_e(t)|^2 - |a_\mu(t)|^2 .$$

Note that the probability to be in $|\nu_e\rangle$ is given by $|a_e(t)|^2$, or equivalently

$$|a_e(t)|^2 = \frac{1}{2} [1 + P_3(t)] . \quad (2.16)$$

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum

The initial value of the polarization vector can be extracted from $\hat{\rho}(0)$, namely,

$$P_1(0) = 0, \quad P_2(0) = 0, \quad P_3(0) = 1. \quad (2.17)$$

Equation of Motion (EoM)

The Liouville-von Neumann equation can be recast into

$$\begin{aligned} i \frac{\partial}{\partial t} \left[P_0 \sigma^0 + \frac{1}{2} \mathbf{P}(t) \cdot \boldsymbol{\sigma} \right] &= \left[\omega_0 \sigma^0 + \frac{1}{2} \omega \mathbf{B} \cdot \boldsymbol{\sigma}, P_0 \sigma^0 + \frac{1}{2} \mathbf{P}(t) \cdot \boldsymbol{\sigma} \right] \\ \Rightarrow i \frac{d}{dt} \mathbf{P}(t) \cdot \boldsymbol{\sigma} &= \frac{1}{2} \omega B_i P_j(t) [\sigma^i, \sigma^j] = i \epsilon^{ijk} \omega B_i P_j \sigma^k \\ \Rightarrow \frac{d}{dt} \mathbf{P}(t) &= \omega \mathbf{B} \times \mathbf{P}(t). \end{aligned} \quad (2.18)$$

The EoM should be solved with the initial conditions in Eq. (2.17).

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum: Geometrical Description




Figure: The polarization vector $\mathbf{P}(t)$ is revolving around the magnetic field $\mathbf{B}$.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum: Geometrical Description

Recall that $\mathbf{B} = (s_{2\theta}, 0, -c_{2\theta})$ and write down the EoM of $\mathbf{P}(t)$ in its three components

$$\dot{P}_1 = \omega (B_2 P_3 - B_3 P_2) = \omega c_{2\theta} P_2, \quad (2.19)$$

$$\dot{P}_2 = \omega (B_3 P_1 - B_1 P_3) = -\omega (c_{2\theta} P_1 + s_{2\theta} P_3), \quad (2.20)$$

$$\dot{P}_3 = \omega (B_1 P_2 - B_2 P_1) = \omega s_{2\theta} P_2. \quad (2.21)$$

The initial conditions are $\mathbf{P}(0) = (0, 0, 1)$ and $\dot{\mathbf{P}}(0) = (0, -\omega s_{2\theta}, 0)$. Then differentiate Eq. (2.20) and insert the other two equations into it. We get

$$\ddot{P}_2 = -\omega^2 P_2, \quad (2.22)$$

whose general solution can be easily found $P_2(t) = A_0 \sin(\omega t + \varphi_0)$. The constants A_0 and φ_0 will be determined by the initial conditions.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum: Geometrical Description

- For $P_2(0) = 0$, we have

$$P_2(0) = A_0 \sin \varphi_0 = 0 ; \quad (2.23)$$

- For $\dot{P}_2(0) = -\omega s_{2\theta}$, we obtain

$$\dot{P}_2(0) = \omega A_0 \cos \varphi_0 = -\omega s_{2\theta} . \quad (2.24)$$

Hence $A_0 = -s_{2\theta}$ and $\varphi_0 = 0$, leading to $P_2(t) = -s_{2\theta} \sin(\omega t)$, which can be inserted back into Eq. (2.22) to get

$$\dot{P}_3(t) = -\omega s_{2\theta}^2 \sin(\omega t) , \quad (2.25)$$

with the initial condition $P_3(0) = 1$. The solution is

$$P_3(t) = s_{2\theta}^2 \cos(\omega t) + c_{2\theta}^2 . \quad (2.26)$$

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Vacuum: Geometrical Description

Finally, we arrive at the survival probability

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= |a_e(t)|^2 = \frac{1}{2} [1 + P_3(t)] \\ &= \frac{1}{2} [1 + c_{2\theta}^2 + s_{2\theta}^2 \cos(\omega t)] \\ &= 1 - s_{2\theta}^2 \frac{1 - \cos(\omega t)}{2} \\ &= 1 - s_{2\theta}^2 \sin^2 \frac{\omega t}{2} = 1 - \sin^2 2\theta \sin^2 \frac{\Delta m_{21}^2 L}{4E}, \end{aligned}$$

where $\omega = \Delta m_{21}^2/(2E)$ and $t \approx L$ have been used. The ordinary oscillation probability is reproduced in the geometrical language.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Matter

We turn to two-flavor neutrino oscillations in matter for which the effective Hamiltonian is given by

$$\begin{aligned}\hat{H}_m &= \frac{1}{2E} \left[\begin{pmatrix} m_1^2 c_\theta^2 + m_2^2 s_\theta^2 & \Delta m_{21}^2 s_\theta c_\theta \\ \Delta m_{21}^2 s_\theta c_\theta & m_1^2 s_\theta^2 + m_2^2 c_\theta^2 \end{pmatrix} + \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix} \right] \\ &= \frac{1}{4E} \left[\begin{pmatrix} -\Delta m_{21}^2 c_{2\theta} & \Delta m_{21}^2 s_{2\theta} \\ \Delta m_{21}^2 s_{2\theta} & \Delta m_{21}^2 c_{2\theta} \end{pmatrix} + A \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right] \\ &\equiv \frac{1}{2} \omega \mathbf{B} \cdot \boldsymbol{\sigma} + \frac{1}{2} \mu \boldsymbol{\lambda} \cdot \boldsymbol{\sigma},\end{aligned}\tag{2.27}$$

where $\omega \equiv \Delta m_{21}^2/(2E)$ is the oscillation frequency in vacuum, the matter potential is $\mu \equiv \sqrt{2}G_F N_e = A/(2E)$, and $\boldsymbol{\lambda} = (0, 0, 1)$ contributes as an extra magnetic field.

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Matter

The effective “magnetic field” can be computed as

$$\begin{aligned}\hat{H}_m &= \frac{1}{2}\omega\mathbf{B}\cdot\boldsymbol{\sigma} + \frac{1}{2}\mu\boldsymbol{\lambda}\cdot\boldsymbol{\sigma} \\ &= \frac{1}{2}\omega_m\mathbf{B}_m\cdot\boldsymbol{\sigma},\end{aligned}\tag{2.28}$$

where the effective mass-squared difference $\Delta\tilde{m}_{21}^2$ and the effective mixing angle $\tilde{\theta}$ can be identified as below

$$\omega\mathbf{B} + \mu\boldsymbol{\lambda} = (\omega s_{2\theta}, 0, \mu - \omega c_{2\theta}) = \frac{\Delta\tilde{m}_{21}^2}{2E}(s_{2\tilde{\theta}}, 0, c_{2\tilde{\theta}}) \equiv \omega_m\mathbf{B}_m\tag{2.29}$$

with

$$\omega_m = \frac{\Delta\tilde{m}_{21}^2}{2E} = \sqrt{(\mu - \omega c_{2\theta})^2 + \omega^2 s_{2\theta}^2}.\tag{2.30}$$

Density Matrix Formalism: Neutrino Oscillations

Neutrino Oscillations in Matter: Geometrical Description




Figure: The polarization vector $\mathbf{P}(t)$ is revolving around the magnetic field $\mathbf{B}_m$.

Why Yukawa interactions?

Main features of the standard model include (1) minimal particle content; (2) gauge symmetries & spontaneous breaking; (3) renormalizability.

Q: Why not mass terms? Take quarks for example. We have left-handed quark doublets and right-handed singlets

$$Q_i = \begin{pmatrix} u_{iL} \\ d_{iL} \end{pmatrix}, \quad u_{iR}, \quad d_{iR}. \quad (3.1)$$

The mass terms are not gauge invariant

$$(M_u)_{ij} \overline{u_{iL}} u_{jR} + (M_d)_{ij} \overline{d_{iL}} d_{jR} + \text{h.c.}, \quad (3.2)$$

but Yukawa interactions are gauge invariant

$$(Y_u)_{ij} \overline{Q}_i \tilde{H} u_{jR} + (Y_d)_{ij} \overline{Q}_i H d_{jR} + \text{h.c.} \quad (3.3)$$

Majorana Fermions

Why Majorana Condition?

The Dirac equation for a massive fermion reads

$$i\gamma^\mu \partial_\mu \psi_L = m\psi_R , \quad (3.4)$$

$$i\gamma^\mu \partial_\mu \psi_R = m\psi_L . \quad (3.5)$$

with $\psi = \psi_L + \psi_R$. Hence ψ_L is already related to ψ_R . However, the Majorana equation is

$$i\gamma^\mu \partial_\mu \psi_L = m\psi_L^C , \quad (3.6)$$

$$i\gamma^\mu \partial_\mu \psi_L^C = m\psi_L . \quad (3.7)$$

with $\psi = \psi_L + \psi_L^C$. The number of independent equations should be equal to the number of degrees, namely, two for Majorana fermions and four for Dirac fermions.

Right-handed Neutrinos

Dirac Neutrinos

For massive Dirac neutrinos, we have

$$\hat{\nu} = \hat{\nu}_L + \hat{\nu}_R, \quad (3.8)$$

where there should be no confusion at all as massive Dirac fermions should have independent left- and right-handed chiral components.

Majorana Neutrinos

For massive Majorana neutrinos in seesaw models, we adopt the same notation ν_R . For clarity, one can use N_R instead of ν_R . In this case,

$$\hat{\nu} = \hat{\nu}_L + \hat{\nu}_L^C \quad (3.9)$$

for light Majorana neutrinos $\hat{\nu}^C = \hat{\nu}$; but for heavy Majorana neutrinos

$$\hat{N} = \hat{N}_R^C + \hat{N}_R \quad (3.10)$$

Right-handed Neutrinos

Majorana Neutrinos

Let us recapitulate the one-generation version of seesaw model. Adding one right-handed neutrino N_R into the SM, we can obtain the neutrino mass terms

$$\mathcal{L}_{\text{mass}}^\nu = -\frac{1}{2} \overline{(\nu_L \quad N_R^C)} \begin{pmatrix} 0 & m_D \\ m_D & m_N \end{pmatrix} \begin{pmatrix} \nu_L^C \\ N_R \end{pmatrix} + \text{h.c.} , \quad (3.11)$$

where $\nu_L^C \equiv C\overline{\nu_L}^T$ and $N_R^C \equiv C\overline{N_R}^T$, and $\{m_D, m_N\}$ are positive numbers.

Diagonalization

The overall mass matrix

$$\mathcal{M} = \begin{pmatrix} 0 & m_D \\ m_D & m_N \end{pmatrix} \quad (3.12)$$

is real and symmetric, so it can be exactly diagonalized.

Right-handed Neutrinos

Diagonalization

First, let us diagonalize $\mathcal{M}$ by an orthogonal matrix $\mathcal{O}$, i.e.,

$$\mathcal{O} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}, \quad (3.13)$$

$$\mathcal{O}^T \mathcal{M} \mathcal{O} = \begin{pmatrix} x_1 & 0 \\ 0 & x_2 \end{pmatrix}, \quad (3.14)$$

where the mixing angle and the eigenvalues can be exactly calculated:

$$x_1 = \frac{1}{2} \left[m_N - \sqrt{m_N^2 + 4m_D^2} \right], \quad (3.15)$$

$$x_2 = \frac{1}{2} \left[m_N + \sqrt{m_N^2 + 4m_D^2} \right], \quad (3.16)$$

$$\tan 2\theta = \frac{2m_D}{m_N}. \quad (3.17)$$

Right-handed Neutrinos

Diagonalization

Assuming $m_N \gg m_D$, we have the excellent approximations

$$x_1 = \frac{1}{2} \left[m_N - \sqrt{m_N^2 + 4m_D^2} \right] \approx -m_D^2/m_N, \quad (3.18)$$

$$x_2 = \frac{1}{2} \left[m_N + \sqrt{m_N^2 + 4m_D^2} \right] \approx m_N, \quad (3.19)$$

$$\theta \approx m_D/m_N. \quad (3.20)$$

Since x_1 is negative, it cannot be the mass of the light Majorana neutrino. We must perform another transformation

$$\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}^T \begin{pmatrix} -m_D^2/m_N & 0 \\ 0 & m_N \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} m_L & 0 \\ 0 & m_H \end{pmatrix}, \quad (3.21)$$

where $m_L = m_D^2/m_N$ and $m_H = m_N$ are light and heavy neutrino masses.

Right-handed Neutrinos

Mass Eigenstates

Since the overall mass matrix $\mathcal{M}$ is diagonalized via

$$V^\dagger \mathcal{M} V^* = \begin{pmatrix} m_L & 0 \\ 0 & m_H \end{pmatrix}, \quad V = \mathcal{O} \begin{pmatrix} -i & 0 \\ 0 & 1 \end{pmatrix}, \quad (3.22)$$

we rewrite the neutrino mass terms as

$$\mathcal{L}_{\text{mass}}^\nu = -\frac{1}{2} \overline{(\nu_L \quad N_R^C)} V \begin{pmatrix} m_L & 0 \\ 0 & m_H \end{pmatrix} V^T \begin{pmatrix} \nu_L^C \\ N_R \end{pmatrix} + \text{h.c.} . \quad (3.23)$$

Therefore, the mass eigenstates of Majorana neutrinos can be found

$$\hat{\nu}_L = V_{11}^* \nu_L + V_{21}^* N_R^C, \quad (3.24)$$

$$\hat{N}_R = V_{12} \nu_L^C + V_{22} N_R, \quad (3.25)$$

where $V_{11} = i c_\theta$, $V_{12} = s_\theta$, $V_{21} = i s_\theta$ and $V_{22} = c_\theta$.