

Spin Alignment, Phase Transition and Spectral Features of QGP in the Presences of Magnetic and Vorticity Fields


- 1. The Spin Alignment and Phase Structure of Thermal QGP under Rotation**
- 2. The Anomalous Magnetic Moment and Phase Transition of the Magnetized QCD Background**
- 3. Spectral features of heavy quarkonia in a rotating and anisotropic QGP:
A holographic study**
- 4. Summary and Conclusions**

Sheng-Qin Feng (冯笙琴) (Three Gorges Univ.)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

Extremal Vorticity and Magnetic Fields in Relativistic HIC

For non-central high-energy collisions, the dense matter produced in the overlapped region of the collision will carry a global angular momentum along the direction opposite to the reaction plane ($-\hat{y}$). Assuming that a partonic system is formed immediately following the initial collision, interactions among produced partons will lead to formation of a quark – gluon plasma (QGP).



Vorticity fields $\omega \sim 10^{21} s^{-1}$

STAR, Nature 548 (2017) 62-65;

Becattini, Liao, Lisa,

Lect.Notes Phys. 987 (2021).

Magnetic fields $B \sim 10^{18} \text{ Gauss}$

W. T. Deng, X. G. Huang,

PRC 85, 044907 (2012).

QCD

Phase structure

Spin polarization

Jet quenching

Meson/baryon Spectra

$\vdots$

NJL model

Quark meson model

Lattice QCD

Holographic model

$\vdots$

1. The Spin Alignment and Phase Structure of Thermal QGP under Rotation

(1) Y. Hua and S.-Q. Feng, Phys. Rev. D **111**, 036012 (2025);

(2) Y.-R. Bao and S.-Q. Feng, Phys. Rev. D **109**, 096033 (2024).

第二届旋转与磁场下的QCD物质研讨会, 淮南 2026. 9. 12-14

From Gluon Topology to Quark Chirality

$$Q_w = \frac{1}{32\pi^2} \int d^4x F^{a\mu\nu} \tilde{F}^a_{\mu\nu}$$

In the quark–gluon plasma (QGP), once these gluon configurations are excited at a certain spacetime point, the topological charge of the vacuum around that point will be altered by these gluon configurations. The chirality imbalance $N_5 = N_R - N_L = -2Q_w$ is induced by the nonzero topological charge through the axial anomaly of QCD.

A non-zero Q_w will induce a corresponding fluctuation in the quark chiral imbalance N_5 . This process will result in chiral imbalance between right and left quarks, leading to the violation of parity (P) and charge-parity (CP) symmetry in the thermal plasma.

The effects of a chiral imbalance in a medium can be implemented in the grand canonical ensemble by introducing a chiral chemical potential μ_5 .

The Lagrangian of NJL model with the chiral chemical potential μ_5

The QCD matter produced in heavy-ion non-central collisions can rotate rapidly with local angular velocities ranging from 0.01 to 0.1 GeV. The Lagrangian of two flavors under the mean field approximation of the NJL model is given

chiral imbalance

Orbital motion

Spin motion

$$\mathcal{L}_{MFA} = \bar{\psi} \left[i\gamma^\mu \partial_\mu - M + \mu\gamma^0 + \mu_5 \gamma^0 \gamma^5 + (\gamma^0)^{-1} \left((\vec{\omega} \times \vec{x}) \cdot (-i\vec{\partial}) + \vec{\omega} \cdot \vec{S}_{4 \times 4} \right) \right] - G_s \sigma^2$$

In cylindrical coordinates, the general positive-energy solutions for the quark field from the Dirac equation corresponding to the above Lagrangian is given as

Reference:

- (1). Y. Jiang and J. Liao, Phys. Rev. Lett. 117, 192302 (2016);
- (2). Y. Hua and S.-Q. Feng, Phys. Rev. D 111, 036012 (2025)

$$\psi(\theta, r) = e^{-iEt + iP_z z} \begin{pmatrix} ce^{in\theta} J_n(p_t r) \\ ide^{i(n+1)\theta} J_{n+1}(p_t r) \\ c'e^{in\theta} J_n(p_t r) \\ id'e^{i(n+1)\theta} J_{n+1}(p_t r) \end{pmatrix}$$

The energy level, thermodynamic potential and chiral charge density⁷

Through the calculation of the finite temperature field, the energy level and thermodynamic potential by rotation are obtained as follows:

$$E_{n,s} = \sqrt{\left(\sqrt{p_t^2 + p_z^2} - s\mu_5\right)^2 + M^2} - \left(n + \frac{1}{2}\right)\omega$$


$$\Omega = \frac{(M-m)^2}{4G_s} - \frac{N_f N_c}{8\pi^2} \sum_{n=-\infty}^{+\infty} \sum_{s=\pm 1} \int dp_t^2 dp_z W_{n,s} \left\{ E_{n,s} + T \ln \left[1 + e^{-\beta(E_{n,s}-u)} \right] + T \ln \left[1 + e^{-\beta(E_{n,s}+u)} \right] \right\}$$

$$W_{n,s} = \left[J_n^2(p_t r) + \lambda^2 J_{n+1}^2(p_t r) \right] / (1 + \lambda^2)$$

Gap equation: $\frac{\partial \Omega}{\partial M} = 0, \quad \frac{\partial^2 \Omega}{\partial M^2} > 0$

The chiral charge density n_5 is defined by : $n_5 = -\frac{\partial \Omega}{\partial \mu_5}$

The phase structure of $T - \omega$ plane with chiral chemical potential



This figure illustrates the chiral phase diagram in the $T_{pc} - \omega$ plane for varying chiral chemical potentials μ_5 . The critical temperature T_{pc} decreases with increasing μ_5 , but the critical end point (CEP) shifts toward higher T and lower ω .

It resolves how **chiral imbalance** alters the order and criticality of the chiral phase transition under rotation, providing insights into the **interplay of vorticity and parity violation in quark-gluon plasma (QGP)**.

Y. Hua and S.-Q. Feng, Phys. Rev. D 111, 036012 (2025)

Spin Alignment of ρ Mesons

The ρ meson is a vector meson composed of a quark and an antiquark with parallel spins (total spin $s = 1$). In extreme environments such as the rotating quark-gluon plasma (QGP) produced in non-central heavy-ion collisions, ρ mesons may exhibit spin alignment — a phenomenon where their spin states correlate with the direction of system rotation.

The key observable for studying spin alignment is the spin density matrix element ρ_{00} , which describes the probability of the meson being in the spin states $s_z = 0$. In the absence of polarization, $\rho_{00} = 1/3$ indicating isotropic spin distribution. A deviation from 1/3 signals spin polarization along a specific direction.

Spin Alignment of ρ Mesons

The spin alignment ρ_{00} is derived using a quark recombination model, where polarized quarks and antiquarks combine to form ρ mesons:

$$\rho_{00} = \frac{1 - P_q P_{\bar{q}}}{3 + P_q P_{\bar{q}}} \approx \frac{1}{3} - \frac{4}{9} P_q P_{\bar{q}}$$

P_q and $P_{\bar{q}}$ are the polarization of quarks and antiquarks, respectively.

[1] Z.-T. Liang and X.-N. Wang, Phys. Rev. Lett. **94**, 102301 (2005);

[2] Z.-T. Liang and X.-N. Wang, Phys. Lett. **B 629**, 20 (2005).

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14



Spin Alignment of ρ Mesons

According to the quark recombination model [1, 2], the spin alignment of fermions is composed of the particle number densities of quarks and antiquarks, which is given by the partial derivative of the thermodynamic potential with respect to the chemical potential:

$$N_{\uparrow}^{+} = \frac{N_f N_c}{4\pi^2} \sum_{n=-\infty}^{+\infty} \sum_{s=\pm 1} \int dp_t dp_z p_t \frac{J_n^2(p_t r)}{1 + \lambda^2} \frac{e^{-\beta(E_{n,s} - \mu)}}{1 + e^{-\beta(E_{n,s} - \mu)}}$$

$$N_{\downarrow}^{+} = \frac{N_f N_c}{4\pi^2} \sum_{n=-\infty}^{+\infty} \sum_{s=\pm 1} \int dp_t dp_z p_t \frac{\lambda^2 J_{n+1}^2(p_t r)}{1 + \lambda^2} \frac{e^{-\beta(E_{n,s} - \mu)}}{1 + e^{-\beta(E_{n,s} - \mu)}}$$

$$N_{\uparrow}^{-} = -\frac{N_f N_c}{4\pi^2} \sum_{n=-\infty}^{+\infty} \sum_{s=\pm 1} \int dp_t dp_z p_t \frac{J_n^2(p_t r)}{1 + \lambda^2} \frac{e^{-\beta(E_{n,s} + \mu)}}{1 + e^{-\beta(E_{n,s} + \mu)}}$$

$$N_{\downarrow}^{-} = -\frac{N_f N_c}{4\pi^2} \sum_{n=-\infty}^{+\infty} \sum_{s=\pm 1} \int dp_t dp_z p_t \frac{\lambda^2 J_{n+1}^2(p_t r)}{1 + \lambda^2} \frac{e^{-\beta(E_{n,s} + \mu)}}{1 + e^{-\beta(E_{n,s} + \mu)}}$$

[1] Z.-T. Liang and X.-N. Wang, Phys. Rev. Lett. **94**, 102301 (2005);

[2] Z.-T. Liang and X.-N. Wang, Phys. Lett. **B 629**, 20 (2005).



Comparison with other models

11

When studying the spin alignment of vector mesons ρ , besides our quark recombination model, there is also the quark condensation model [1]. The spin alignment corresponding to vector mesons ρ can take the following form


$$\rho_{00} = \frac{1}{3} - \frac{4}{9}(\beta\omega)^2$$

Another spin polarization model is self-consistent NJL model [2], the alignment of vector mesons (including ρ and ϕ) ($T = 150$ MeV), exhibits the following relationship:


$$\rho_{00} = \frac{1}{3} - 5.10\omega^2 + 39.62\omega^4$$

[1] Y.-G. Yang, R.-H. Fang, Q. Wang, and X.-N. Wang, Phys. Rev. C 97, 034917(2018);

[2] M. Wei and M. Huang, Chin. Phys. C 47, 104105(2023).



The relationship between spin alignment ρ_{00} and temperature T



Spin alignment ρ_{00} of ρ mesons as a function of temperature T

ρ_{00} approaches $1/3$ (isotropic spin alignment) at high T , while deviating $1/3$ near the phase transition temperature.

Increasing μ_5 enhances ρ_{00} , reducing spin polarization.


Innovation & Impact: This is the first work to connect chiral imbalance (μ_5) with vector meson spin alignment, **showing that μ_5 counteracts rotation-induced polarization near T_{pc} .**

Y. Hua and S.-Q. Feng, Phys. Rev. D 111, 036012 (2025)

Key Problem Solved: **It explains how chiral imbalance moderates spin polarization in a temperature-dependent manner**, crucial for interpreting experimental ρ_{00} data in QGP.

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

The relationship between spin alignment ρ_{00} and rotational velocity ω



Research Content: ρ_{00} decreases with ω , deviating from $1/3$ (indicating polarization). large ω makes more polarization, but larger μ_5 reduces this deviation, particularly at high ω , makes less polarization.

Innovation & Impact: It demonstrates that chiral imbalance suppresses rotation-induced spin polarization, offering a mechanism to tune meson spin states via μ_5


Spin alignment ρ_{00} of ρ mesons as a function of ω

Key Problem Solved: It resolves how competing effects of rotation and chiral imbalance determine the net polarization of vector mesons, vital for probing spin-orbit coupling in QCD matter.

Y. Hua and S.-Q. Feng, Phys. Rev. D 111, 036012 (2025)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

The relationship between spin alignment ρ_{00} and rational radius



Research Content: ρ_{00} increases with r , approaching $1/3$, showing reduced polarization away from the rotation center. μ_5 enhances ρ_{00} at all r , especially near T_{pc} .

Innovation & Impact: This study pioneers the spatial analysis of spin alignment in rotating systems, revealing radial gradients in polarization and the moderating role of μ_5 .

Spin alignment ρ_{00} of ρ mesons as a function of radius r

Y. Hua and S.-Q. Feng, Phys. Rev. D 111, 036012 (2025)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

Key Problem Solved: It identifies the spatial inhomogeneity of spin polarization in vortical QGP, essential for understanding local spin dynamics in finite-size systems like heavy-ion collisions.

2. Anomalous Magnetic Moment (AMM) and Phase transition in Magnetized QCD Matter

- (1) C.-Y. Yang and S.-Q. Feng, Phys. Rev. D **112**, 036008 (2025);
- (2) X.-Q. Zhu and S.-Q. Feng, Phys. Rev. D **107**, 016018 (2023);
- (3) Y.-W. Qiu and S.-Q. Feng, X.-Q. Zhu, Phys. Rev. D **108**, 116022 (2023)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

The SU(3) NJL model with quark anomalous magnetic moment (AMM)

Lagrangian density (**three flavors, broken isospin symmetry**)

Landau gauge $A_\mu = (0, 0, xB, 0)$

$$\mathcal{L}_{NJL} = \sum_{f=u,d,s} \bar{\psi}_f (i\gamma^\mu D_\mu^{(f)} - m_f - \frac{1}{2} e_f \kappa_f \sigma^{\mu\nu} F_{\mu\nu}) \psi_f + G \sum_{a=0}^8 [(\bar{\psi} \lambda_a \psi)^2 + (\bar{\psi} i\gamma^5 \lambda_a \psi)^2] - K(\det[\bar{\psi}(1 + \gamma^5)\psi] + \det[\bar{\psi}(1 - \gamma^5)\psi]).$$

M. Strickland, V. Dexheimer, and D. P. Menezes, Phys. Rev.D 86, 125032 (2012).

The contribution of the anomalous magnetic moment (AMM) to the Lagrangian density does arise from **the interaction between the AMM and the quark spin.**

In the NJL Lagrangian used , the AMM term is:

$$-\frac{1}{2} e_f \kappa_f \sigma^{\mu\nu} F_{\mu\nu}$$

- $\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$ is the **spin tensor operator**, which is directly related to the quark's spin structure.
- $F_{\mu\nu}$ is the electromagnetic field strength tensor.
- κ_f is the flavor-dependent AMM of the quark.

The SU(3) NJL model with quark anomalous magnetic moment (AMM)

Mean-field approximation:

Thermodynamic potential: $\Omega_{MF} = \Omega_q + 2G(\sigma_u^2 + \sigma_d^2 + \sigma_s^2) - 4K\sigma_u\sigma_d\sigma_s$

$$\Omega_q = -3 \sum_{f=u,d,s} \frac{|q_f B|}{2\pi} \sum_n \sum_{s=\pm 1} \int \frac{dp_z}{2\pi} \left[E_{fns} + T \ln \left(1 + e^{-\frac{E_{fns} + \mu}{T}} \right) + T \ln \left(1 + e^{-\frac{E_{fns} - \mu}{T}} \right) \right]$$

Gap eqs: $\partial \Omega_{MF} / \partial \sigma_f = 0 \implies M_f = m_f - 4G\sigma_f + 2K \prod_{f'=f} \sigma_{f'}$

Ω_{MF} nizing:

Chiral condensates: $\sigma_f = \langle \bar{\psi}_f \psi_f \rangle = -\frac{|e_f B|}{(2\pi)^2} \int dp_z \sum_{n,s} \frac{M_f}{E_{fns}} \left(1 - \frac{s\kappa_f e_f B}{M_{nf}} \right) \left(1 - \frac{1}{e^{(E_{fns} - \mu)/T} + 1} - \frac{1}{e^{(E_{fns} + \mu)/T} + 1} \right)$

Energy eigenvalue:

$$E_{fns} = \sqrt{p_z^2 + (M_{nf} - \underline{s\kappa_f e_f B})^2}, \quad M_{nf} = \sqrt{M_f^2 + 2n|e_f B|}$$

AMM

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

The SU(3) NJL model with quark AMM

Transforming the six-fermion interaction into an effective four-fermion interaction,

➤ one obtains:

$$\begin{aligned}\mathcal{L}_{NJL} = & \sum_{f=u,d,s} \bar{\psi}_f (i\gamma^\mu D_\mu^{(f)} - m_f - \frac{1}{2} e_f \kappa_f \sigma^{\mu\nu} F_{\mu\nu}) \psi_f \\ & + \sum_{a=0}^8 [K_a^- (\bar{\psi} \lambda_a \psi)^2 + K_a^+ (\bar{\psi} i\gamma^5 \lambda_a \psi)^2] \\ & + K_{30}^- (\bar{\psi} \lambda_3 \psi) (\bar{\psi} \lambda_0 \psi) + K_{30}^+ (\bar{\psi} i\gamma^5 \lambda_3 \psi) (\bar{\psi} i\gamma^5 \lambda_0 \psi) \\ & + K_{80}^- (\bar{\psi} \lambda_8 \psi) (\bar{\psi} \lambda_0 \psi) + K_{80}^+ (\bar{\psi} i\gamma^5 \lambda_8 \psi) (\bar{\psi} i\gamma^5 \lambda_0 \psi) \\ & + K_{83}^- (\bar{\psi} \lambda_8 \psi) (\bar{\psi} \lambda_3 \psi) + K_{83}^+ (\bar{\psi} i\gamma^5 \lambda_8 \psi) (\bar{\psi} i\gamma^5 \lambda_3 \psi) \\ & + K_{03}^- (\bar{\psi} \lambda_0 \psi) (\bar{\psi} \lambda_3 \psi) + K_{03}^+ (\bar{\psi} i\gamma^5 \lambda_0 \psi) (\bar{\psi} i\gamma^5 \lambda_3 \psi) \\ & + K_{08}^- (\bar{\psi} \lambda_0 \psi) (\bar{\psi} \lambda_8 \psi) + K_{08}^+ (\bar{\psi} i\gamma^5 \lambda_0 \psi) (\bar{\psi} i\gamma^5 \lambda_8 \psi) \\ & + K_{38}^- (\bar{\psi} \lambda_3 \psi) (\bar{\psi} \lambda_8 \psi) + K_{38}^+ (\bar{\psi} i\gamma^5 \lambda_3 \psi) (\bar{\psi} i\gamma^5 \lambda_8 \psi)\end{aligned}$$

Effective coupling constants:

$$K_0^\pm = G \pm \frac{1}{3} K (\sigma_u + \sigma_d + \sigma_s),$$

$$K_1^\pm = K_2^\pm = K_3^\pm = G \pm \frac{1}{2} K \sigma_s,$$

$$K_4^\pm = K_5^\pm = G \pm \frac{1}{2} K \sigma_d,$$

$$K_6^\pm = K_7^\pm = G \pm \frac{1}{2} K \sigma_u,$$

$$K_8^\pm = G \pm \frac{1}{6} K (2\sigma_u + 2\sigma_d - \sigma_s),$$

$$K_{30}^\pm = K_{03}^\pm = \mp \frac{1}{2\sqrt{6}} K (\sigma_u - \sigma_d),$$


$$K_{80}^\pm = K_{08}^\pm = \pm \frac{\sqrt{2}}{12} K (\sigma_u + \sigma_d - 2\sigma_s),$$

$$K_{83}^\pm = K_{38}^\pm = \pm \frac{1}{2\sqrt{3}} K (\sigma_u - \sigma_d),$$


➤ **Chiral condensates:** $\sigma_u = \langle \bar{\psi}_u \psi_u \rangle, \sigma_d = \langle \bar{\psi}_d \psi_d \rangle, \sigma_s = \langle \bar{\psi}_s \psi_s \rangle$

➤ Ideas:

(1) Quarks: mean-field



(2) Mesons: RPA resummation (quantum fluctuation)



S. Klevansky, Rev. Mod. Phys. 64, 649 (1992).
M. Buballa, Phys. Rept. 407, 205 (2005).

➤ Presenting the calculation results directly:

K meson:
$$M = \frac{2K_6^+}{1 - 2K_6^+ \Pi_{K^0}^{ps}(p)}$$

Where


$$\Pi_{K^0}^{ps}(p_0 = m_{K^0}) = \frac{\beta N_c}{2(2\pi)^2} \sum_{n,s,l} \left(1 + sl \frac{M_d M_s + 2\beta n}{M_{nd} M_{ns}} \right)$$

$$\times \left\{ \int dk_z \frac{1}{E_{dns}} \left[1 - \frac{1}{e^{(E_{dns}-\mu)/T} + 1} - \frac{1}{e^{(E_{dns}+\mu)/T} + 1} \right] \right. \\ \left. + \int dk_z \frac{1}{E_{snl}} \left[1 - \frac{1}{e^{(E_{snl}-\mu)/T} + 1} - \frac{1}{e^{(E_{snl}+\mu)/T} + 1} \right] \right.$$

$$\left. + \left\{ [(M_{nd} - s\kappa_d e_d B) - sl(M_{ns} - l\kappa_s e_s B)]^2 - p_0^2 \right\} B(m_d, m_s) \right\}$$

$$B(m_d, m_s) = \int dk_z \left\{ \frac{1}{E_{dns}} \left[\frac{1}{(E_{dns} + p_0)^2 - E_{snl}^2} \frac{1}{e^{-(E_{dns}+\mu)/T} + 1} \right. \right. \\ \left. \left. - \frac{1}{(E_{dns} - p_0)^2 - E_{snl}^2} \frac{1}{e^{(E_{dns}-\mu)/T} + 1} \right] \right. \\ \left. + \frac{1}{E_{snl}} \left[\frac{1}{(E_{snl} - p_0)^2 - E_{dns}^2} \frac{1}{e^{-(E_{snl}+\mu)/T} + 1} \right. \right. \\ \left. \left. - \frac{1}{(E_{snl} + p_0)^2 - E_{dns}^2} \frac{1}{e^{(E_{snl}-\mu)/T} + 1} \right] \right\}.$$

QCD Phase Diagram Restructuring



The critical temperature (T_c) of chiral phase transitions as a function of eB at different chemical potentials (μ), with and without AMM.



- Magnetic catalysis without AMM
- Inverse magnetic catalysis with AMM
- Non-contradiction with LQCD calculation results.

(1) Phase Diagram Reshaping and CEP Shift: AMM suppresses T_c and shifts critical endpoints (CEP) toward lower eB and higher T , contrasting conventional NJL predictions dominated by magnetic catalysis (MC).

(2) Crossover-to-First-Order Transition: AMM replaces crossover transitions with first-order transitions under strong eB , validating AMM's role in altering phase transition sequences in multi-flavor systems.

- (1) C.-Y. Yang and S.-Q. Feng, Phys. Rev. D **112**, 036008 (2025);
- (2) X.-Q. Zhu and S.-Q. Feng, Phys. Rev. D **107**, 016018 (2023);
- (3) Y.-W. Qiu and S.-Q. Feng, X.-Q. Zhu, Phys. Rev. D **108**, 116022 (2023).

QCD Phase Diagram Restructuring




The critical temperature (T_c) for chiral phase transition as a function of chemical potential for different magnetic fields

- (1) C.-Y. Yang and S.-Q. Feng, Phys. Rev. D **112**, 036008 (2025);
- (2) X.-Q. Zhu and S.-Q. Feng, Phys. Rev. D **107**, 016018 (2023);
- (3) Y.-W. Qiu and S.-Q. Feng, X.-Q. Zhu, Phys. Rev. D **108**, 116022 (2023).

- Crossover transitions into first-order transitions without AMM
- Uniform manifestation as first-order phase transitions with AMM
- T_c is a decreasing function of μ

The magnetic dependences of neutral pseudoscalar meson mass spectra



- (1) C.-Y. Yang and S.-Q. Feng, *Phys. Rev. D* **112**, 036008 (2025);
(2) X.-Q. Zhu and S.-Q. Feng, *Phys. Rev. D* **107**, 016018 (2023).

(1) Meson Mass Collapse and Chiral Restoration: AMM triggers abrupt mass collapses (e.g., π at $eB \approx 0.1 \text{ GeV}^2$), directly linking meson stability to chiral symmetry restoration.

(2) η' Resonance Limitations: η' mass diverges under strong eB due to non-perturbative decay width effects, exposing NJL's limitations in handling resonance states.

Key Issues Addressed:

Systematically reveals AMM's role in regulating multi-flavor meson masses and flavor mixing (e.g., $\pi^0 - \eta - \eta'$), addressing gaps in two-flavor models' predictions for magnetic sensitivity.

3. Spectral Signatures of Heavy Quarkonia in a Rotating and Anisotropic QGP: A Holographic Study

- (1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D 113, 096006 (2026)
- (2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D 110, 086018 (2024)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

Global Rotation and Spatial Anisotropy

Non-central heavy-ion collisions produce a strongly coupled QGP with:

Global rotation (angular velocity $\omega \sim 0.01 - 0.1 \text{ GeV}$)

Spatial anisotropy (expansion along collision axis)

Heavy quarkonia (J/Ψ , $\Upsilon(1S)$) are sensitive probes of the QGP medium.

Key Questions:

How do rotation and anisotropy jointly modify in-medium spectral functions and effective masses?

Do they exhibit directional selectivity in dissociation?

Is their effect additive or competitive?

Theoretical Framework – Holographic Setup

Used Gauge/Gravity Duality (AdS/CFT) as a non-perturbative tool

5D Einstein-Dilaton-Two-Maxwell action

First Maxwell field $\rightarrow$ finite chemical potential μ

Second Maxwell field $\rightarrow$ spatial anisotropy

Dilaton field $\phi(z)$ with parameters fitted to charmonium/bottomonium spectra.

Anisotropic metric (stationary frame):

$$d\tilde{s}^2 = \frac{b(z)}{z^2} \left[-g(z)dt^2 + dx_1^2 + z^{2-2/\nu}(dx_2^2 + dx_3^2) + \frac{dz^2}{g(z)} \right]$$

x_1 : longitudinal (anisotropy axis);

x_2, x_3 : transverse;

ν : anisotropy parameter ($1 < \nu \leq 4.5$);

$b(z) = e^{\frac{1}{2}\alpha z^2}$: warp factor.

Rotation introduced via Lorentz boost along angular direction $\rightarrow$ rotating metric with ωl .

(1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D 113, 096006 (2026)

(2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D 110, 086018 (2024)

The key steps for computing the spectral functions

1. Set up the holographic model

(1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D **113**, 096006 (2026)

2. Introduce rotation

(2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D **110**, 086018 (2024)

Apply a Lorentz boost along the angular coordinate to the static metric, obtaining a rotating frame metric with angular velocity ω . The Hawking temperature and chemical potential are rescaled by $\sqrt{1-\omega^2 l^2}$

3. Derive the flow equations for conductivities

For a vector field (dual to heavy quarkonia), decompose fluctuations into longitudinal (along the anisotropy direction x_1) and transverse (x_2, x_3) channels. Using the membrane paradigm, derive first-order differential equations for the longitudinal/transverse conductivities $\sigma_{L,T}(\varepsilon, z)$: $\partial_z \sigma = i\varepsilon \Delta (\sigma^2 - \Sigma^2)$, with Δ and Σ given by the metric coefficients and dilaton background.


4. Solve the flow equations

Impose regularity at the horizon ($\partial_z \sigma = 0$ at $z = z_h$) and integrate numerically from the horizon to the boundary ($z \rightarrow 0$) to obtain the boundary conductivities $\sigma_{L,T}(\varepsilon, 0)$.

5. Extract the spectral function

The spectral function is obtained via the retarded Green's function: $\rho(\varepsilon) \equiv -\text{Im } G_R(\varepsilon) = \varepsilon \text{Re } \sigma(\varepsilon, 0)$.

Temperature dependence of heavy quarkonium spectral functions in a rotating and anisotropic QGP




Temperature dependence: Raising the temperature systematically suppresses the spectral peaks and broadens the widths for both polarizations.

Crucially, the suppression is slightly stronger in the longitudinal (anisotropy-parallel) channel than in the transverse one, revealing that thermal dissociation acquires a subtle directional imprint even when rotation is present.

(1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D **113**, 096006 (2026)

(2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D **110**, 086018 (2024)

Chemical potential dependence of heavy quarkonium spectral functions in a rotating and anisotropic QGP




Chemical-potential dependence: At fixed temperature, increasing the baryon chemical potential μ also suppresses peaks and broadens widths in both longitudinal and transverse polarizations.

The key novelty is that **density-driven dissociation persists in the rotating anisotropic medium, acting as an independent melting mechanism via enhanced color screening.**

- (1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D **113**, 096006 (2026)
- (2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D **110**, 086018 (2024)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

Competitive modulation of heavy quarkonium spectral functions and polarization-dependent dissociation by anisotropy and rotation




(1) Global rotation and spatial anisotropy act not additively, but competitively.

(2) The directional suppression pattern of quarkonia in noncentral heavy-ion collisions reflects a dynamical battle between initial-state spatial deformation (anisotropy, which preferentially disrupts longitudinal polarizations) and later-stage collective vorticity (rotation, which more strongly affects transverse polarizations).

(1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D **113**, 096006 (2026)


(2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D **110**, 086018 (2024)

Modulation of heavy-quarkonium effective masses with temperature by rotation and anisotropy



The in-medium mass shift of heavy quarkonia is not merely thermal but is modulated by distinct directional geometric and dynamical distortions – rotation generates centrifugal effects in the transverse direction, whereas anisotropy stretches the medium along its axis.

Moreover, they uncover polarization-dependent nonmonotonic behavior that is state-sensitive: only the lighter J/ψ exhibits such complexity, while the heavier Υ (1S) remains largely monotonic.



(1) X.-W. Shi and S.-Q. Feng, Phys. Rev. D **113**, 096006 (2026)

(2) X.-L. Wang and S.-Q. Feng, Phys. Rev. D **110**, 086018 (2024)

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

4. Summary and Conclusions

第二届旋转与磁场下的QCD物质研讨会, 淮南2026. 9. 12-14

Summary and conclusions

- 1. The spin alignment of η_{00} and phase structure of thermal QGP under rotation are investigated. It is found that chiral imbalance have some effects on phase structure of QGP medium. We also study some dependences of spin alignment of η_{00} with temperature, angular velocity, rotational radius on chiral imbalance.**
- 2. The research investigates the impact of the anomalous magnetic moment (AMM) of quarks on the mass spectra of neutral pseudoscalar mesons (π , K , η , η') under conditions of strong magnetic fields, finite temperatures, and chemical potentials, based on the three-flavor Nambu-Jona-Lasinio (NJL) model.**
- 3. This holographic study examines spectral functions and effective masses of J/ψ and $\Upsilon(1S)$ in a rotating and anisotropic QGP, using a five-dimensional Einstein-dilaton-two-Maxwell model. We find that both rotation and anisotropy individually enhance quarkonium dissociation (peak suppression and broadening), but with opposite directional biases: anisotropy preferentially suppresses longitudinal polarizations, while rotation more strongly affects transverse ones.**

Thanks !

第二届旋转与磁场下的QCD物质研讨会 , 淮南2026. 9. 12-14