

CPV in $B^+ \rightarrow K^+ \pi^- \pi^+$ decays

Nanyang Wang
on behalf of LHCb collaboration

CLHCP2026, 2026/10/30

- Introduction
- Amplitude analysis
- Physics results and implications
 - CP violation
 - Strong interaction modelling implications
- Conclusion

Introduction: CP violation

- **Direct CPV requires interfering amplitudes with different weak and strong phases**


- Weak phases ϕ from CKM matrix
- Strong phases δ from final-state interactions

$$A_{CP} = \frac{2|T||P| \sin(\delta_T - \delta_P) \sin(\phi_T - \phi_P)}{|T|^2 + |P|^2 + 2|T||P| \cos(\delta_T - \delta_P) \cos(\phi_T - \phi_P)}$$

$$A = |T|e^{i(\delta_T - \phi_T)} + |P|e^{i(\delta_P - \phi_P)}$$

$$\bar{A} = |T|e^{i(\delta_T + \phi_T)} + |P|e^{i(\delta_P + \phi_P)}$$


- Major milestones in CPV studies, **with no conclusive evidence of physics beyond the Standard Model**



- **Multi-body decays with amplitude analysis**
 - Rich resonant structure and interference
 - Access to additional observables: BFs, A_{CP} and relative phases for resonances

Introduction: Previous LHCb results

- Large localised CPV seen model-independently in $B^+ \rightarrow hh'h''$ ($h = K, \pi$) [PRD.108.012008\(2023\), Run2](#)



- Run 1 amplitude analysis: $B^+ \rightarrow \pi^+ \pi^- \pi^+$ and $B^+ \rightarrow \pi^+ K^- K^+$
[PRL.124.031801\(2020\)](#) [PRL.123.231802\(2019\)](#)
[PRD.101.012006\(2020\)](#)
- No previous LHCb amplitude analysis of $B^+ \rightarrow K^+ \pi^- \pi^+$ and $B^+ \rightarrow K^+ K^- K^+$
 - More challenging due to higher statistics
 - Need to model accurately subtle effects

Introduction: Previous results and puzzles

Puzzle 1: $\rho(770)^0$ CP asymmetry

- Evidence for CP violation in $B^+ \rightarrow \rho(770)^0 K^+$ seen in amplitude analysis by BaBar and Belle
- BaBar amplitude analysis: $A_{CP}^{\rho(770)^0} = (+0.44 \pm 0.11)$
PRD.78.012004(2008)


- Model-independent determination of CPV in $\pi\pi$ P-Wave at the $\rho(770)$ peak, by LHCb PRD.108.012013(2023)

$$A_{CP}^{\rho \text{ window}} = (+0.150 \pm 0.022)$$

Puzzle 2: $K_0^*(1430)\pi$ Branching fraction

PRD.87.114001(2013)

- QCdf describes $B \rightarrow K_0^*(1430)^0[\eta', \rho, \omega, \phi]$, but predicts a much smaller $B^+ \rightarrow K_0^*(1430)^0 \pi^+$ BF than measured in BaBar and Belle Eur.Phys.J.C.50.877(2007)



Puzzle 3: The mysterious $f_X(1300)$

- Belle and BaBar introduced a scalar $f_X(1300)$
- Freely parameterised $f_X(1300)$ gives **better fit quality** than $f_0(1370) + f_0(1500)$
- Fitted parameters consistent with $f_0(1370)(\text{Belle})/f_0(1500)(\text{BaBar})$
- If $f_X(1300) = f_0(1500)$
 - $B^+ \rightarrow f_0(1500)K^+$ BF from $B^+ \rightarrow K^+K^-K^+$ is $\sim 10 \times$ larger than implied by $B^+ \rightarrow K^+\pi^-\pi^+$
 - $f_0(1500)$ cannot dominate the $\pi^+\pi^-$ channel

PRD.85.112010(2012)

- Introduction
- **Amplitude analysis**
- Physics results and implications
 - CP violation
 - Strong interaction modelling implications
- Conclusion

Amplitude analysis: Isobar

- 3 alternative S-wave models, following strategy used in $B^+ \rightarrow \pi^+ \pi^- \pi^+$
- 2 different fitters Laura++ and Mint2-devel in use, allows cross-checks
- Isobar**
 - $K\pi$: effective-range-like (ER)+ $K_0^*(1430)^0$, $K_0^*(1950)^0$
 - $\pi\pi$: $f_0(500)$, $f_0(980)$, $f_0(1370)$, $f_0(1500)$, $f_0(1710)$, rescattering

$$\mathcal{A}^\pm(\Phi_3) = \sum_j c_j^\pm F_j(\Phi_3)$$

Contribution	Approach	Mass lineshape	Fixed parameters
$(K^+ \pi^-)_{\text{SVP}}$	Isobar/K-matrix	GLASS (PRD78 034023)	—
$K_0^*(1430)^0$	Isobar/K-matrix	Breit-Wigner	$m_0 = 1.945 \pm 0.022 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.201 \pm 0.086 \text{ GeV}$ (PDG2020)
$K_0^*(1950)^0$			
$f_0(500)$	Isobar	Bugg	fit (iii) (J. Phys. G34 151)
$f_0(980)$	Isobar	Flatté (PLB 63 224)	—
$f_0(1370)$	Isobar	T-matrix pole	$m_0 = 1.38 \pm 0.07 \text{ GeV}/c^2$ (PRL130 051902) $\Gamma_0 = 0.22 \pm 0.08 \text{ GeV}$ (PRL130 051902)
$f_0(1500)$	Isobar	Breit-Wigner	$m_0 = 1.506 \pm 0.006 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.112 \pm 0.009 \text{ GeV}$ (PDG2020)
$f_0(1710)$	Isobar	Breit-Wigner	$m_0 = 1.704 \pm 0.012 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.123 \pm 0.018 \text{ GeV}$ (PDG2020)
$\pi\pi\text{-}\pi\pi$	Isobar	Rescattering	CFD (PRD83 074004)
$\pi\pi\text{-}KK$	Isobar	Rescattering	CFD _B (Phys. Rept. 969 1)
$\pi^+ \pi^-$ S-wave	K-matrix	K-matrix	(PRD78 034023, EPJA16 229)
$K^+ \pi^-$ and $\pi^+ \pi^-$ S-waves	QMI	Constant per bin	—

Amplitude analysis: K-matrix

- K-matrix**



- $K\pi$: same as in Isobar approach
- $\pi\pi$: 5 poles and slow-varying parts, 5 channels
 - Interface to scattering results, approach with unitary condition applied

Contribution	Approach	Mass lineshape	Fixed parameters
$(K^+\pi^-)_{\text{SVP}}$	Isobar/K-matrix	GLASS (PRD78 034023)	—
$K_0^*(1430)^0$	Isobar/K-matrix	Breit–Wigner	$m_0 = 1.945 \pm 0.022 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.201 \pm 0.086 \text{ GeV}$ (PDG2020)
$K_0^*(1950)^0$			
$f_0(500)$	Isobar	Bugg	fit (iii) (J. Phys. G34 151)
$f_0(980)$	Isobar	Flatté (PLB 63 224)	—
$f_0(1370)$	Isobar	T-matrix pole	$m_0 = 1.38 \pm 0.07 \text{ GeV}/c^2$ (PRL130 051902) $\Gamma_0 = 0.22 \pm 0.08 \text{ GeV}$ (PRL130 051902)
$f_0(1500)$	Isobar	Breit–Wigner	$m_0 = 1.506 \pm 0.006 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.112 \pm 0.009 \text{ GeV}$ (PDG2020)
$f_0(1710)$	Isobar	Breit–Wigner	$m_0 = 1.704 \pm 0.012 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.123 \pm 0.018 \text{ GeV}$ (PDG2020)
$\pi\pi\text{-}\pi\pi$	Isobar	Rescattering	CFD (PRD83 074004)
$\pi\pi\text{-}KK$	Isobar	Rescattering	CFD_B (Phys. Rept. 969 1)
$\pi^+\pi^-$ S-wave	K-matrix	K-matrix	(PRD78 034023, EPJA16 229)
$K^+\pi^-$ and $\pi^+\pi^-$ S-waves	QMI	Constant per bin	—

Amplitude analysis: Quasi-Model-Independent


- Quasi-Model-Independent (QMI)

- Simultaneous QMI treatment of the two crossing S-waves
- $K\pi/\pi\pi$: 35/40 bins below charm, 15/10 above
- Bin boundaries aligned with hadronic thresholds to capture expected cusps from rescattering effects



Contribution	Approach	Mass lineshape	Fixed parameters
$(K^+\pi^-)_{\text{SVP}}$	Isobar/K-matrix	GLASS (PRD78 034023)	—
$K_0^*(1430)^0$	Isobar/K-matrix	Breit-Wigner	$m_0 = 1.945 \pm 0.022 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.201 \pm 0.086 \text{ GeV}$ (PDG2020)
$K_0^*(1950)^0$			
$f_0(500)$	Isobar	Bugg	fit (iii) (J. Phys. G34 151)
$f_0(980)$	Isobar	Flatté (PLB 63 224)	—
$f_0(1370)$	Isobar	T-matrix pole	$m_0 = 1.38 \pm 0.07 \text{ GeV}/c^2$ (PRL130 051902) $\Gamma_0 = 0.22 \pm 0.08 \text{ GeV}$ (PRL130 051902)
$f_0(1500)$	Isobar	Breit-Wigner	$m_0 = 1.506 \pm 0.006 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.112 \pm 0.009 \text{ GeV}$ (PDG2020)
$f_0(1710)$	Isobar	Breit-Wigner	$m_0 = 1.704 \pm 0.012 \text{ GeV}/c^2$ (PDG2020) $\Gamma_0 = 0.123 \pm 0.018 \text{ GeV}$ (PDG2020)
$\pi\pi\text{-}\pi\pi$	Isobar	Rescattering	CFD (PRD83 074004)
$\pi\pi\text{-}KK$	Isobar	Rescattering	CFD _B (Phys. Rept. 969 1)
$\pi^+\pi^-$ S-wave	K-matrix	K-matrix	(PRD78 034023, EPJA16 229)
$K^+\pi^-$ and $\pi^+\pi^-$ S-waves	QMI	Constant per bin	—

- Introduction
- Amplitude analysis
- Physics results and implications
 - CP violation
 - Strong interaction modelling implications
- Conclusion



- Reasonable agreement between data and models: all three S-wave models with $\chi^2/\text{ndf} \sim 1.6$ without considering systematic uncertainties

- CP violation of individual component: $A_{CP}^i = \frac{F_i^- - F_i^+}{F_i^- + F_i^+}$
- CP violation of interference between two components: $A_I^{ij} = (I_{ij}^- - I_{ij}^+)/F_{ij}$

Parameter	Isobar	K-matrix	QMI
$\mathcal{A}_{CP}(K^*(892)^0)$	$-0.038 \pm 0.010 \pm 0.018$ (1.9 σ)	$-0.035 \pm 0.012 \pm 0.017$ (1.7 σ)	$-0.027 \pm 0.014 \pm 0.023$ (1.0 σ)
$\mathcal{A}_{CP}(K_0^*(1430)^0)$	$-0.094 \pm 0.020 \pm 0.046$ (1.9 σ)	$-0.121 \pm 0.015 \pm 0.050$ (2.3 σ)	—
$\mathcal{A}_{CP}(\rho(770)^0)$	$+0.280 \pm 0.022 \pm 0.029$ (7.6 σ)	$+0.271 \pm 0.022 \pm 0.015$ (10.2 σ)	$+0.359 \pm 0.037 \pm 0.048$ (5.9 σ)
$\mathcal{A}_{CP}^S(m_{\pi^+\pi^-} < 4m_\pi)$	$+0.665 \pm 0.032 \pm 0.053$ (>10 σ)	$+0.638 \pm 0.008 \pm 0.016$ (>10 σ)	$+0.754 \pm 0.115 \pm 0.113$ (4.7 σ)
$\mathcal{A}_I^{\text{SD}} (\times 10^{-1})$	$+0.058 \pm 0.007 \pm 0.008$ (5.2 σ)	$+0.055 \pm 0.007 \pm 0.007$ (5.3 σ)	$+0.071 \pm 0.014 \pm 0.009$ (4.4 σ)
$\mathcal{A}_I^{\text{SP}} (\times 10^{-1})$	$-0.237 \pm 0.018 \pm 0.040$ (5.4 σ)	$-0.249 \pm 0.021 \pm 0.040$ (5.5 σ)	$-0.253 \pm 0.025 \pm 0.028$ (6.7 σ)
$\mathcal{A}_{CP}(\rho_3(1690)^0)$	$+0.603 \pm 0.090 \pm 0.121$ (4.0 σ)	$+0.477 \pm 0.100 \pm 0.096$ (3.4 σ)	$+0.698 \pm 0.135 \pm 0.163$ (3.3 σ)

- Angular distribution: a sum of Legendre polynomials


helicity angle $\frac{dN}{d \cos \theta} \propto 1 + \sum_{n=1}^N \text{Legendre moments } a_n P_n(\cos \theta)$

- Sensitivity of Legendre moments to partial-wave interference

$$\langle P_1 \rangle \propto SP \cos(\theta_S - \theta_P) + 2\sqrt{\frac{2}{5}} PD \cos(\theta_P - \theta_D)$$



$$\langle P_2 \rangle \propto \sqrt{\frac{2}{5}} P^2 + \sqrt{\frac{10}{7}} D^2 + \sqrt{2} SD \cos(\theta_S - \theta_D)$$

$$\Delta \langle P_n \rangle \equiv \langle P_n \rangle_{B^+} - \langle P_n \rangle_{B^-}$$




CP violation results: CPV in $\rho(770)^0 K^+$

Probably associated with known phase variation





Explains puzzle 1: $\rho(770)^0$ CP asymmetry




- Clear positive CP asymmetry for S-wave in $m_{\pi^+\pi^-} < 4m_\pi$
- Clear positive CP asymmetry for $\rho(770)^0$
- Observed CP violation in S-P interference, sign change around the $\rho(770)^0$ pole, very similar to $B^+ \rightarrow \pi^+\pi^-\pi^+$
- No significant CPV in ω and $\rho - \omega$ mixing contribution, diluting the overall asymmetry in the $\pi\pi$ P-wave


CP violation results: $f_2(1270)$ CPV





- Raw asymmetry in the $\pi\pi$ helicity angle in $f_2(1270)$ mass range
- $\cos \theta_{\pi^+\pi^-} < -0.5$: Negative CP asymmetry for $f_2(1270)$ ✓
- In other regions S-wave CP asymmetry cancels or dominates D-wave contribution
- S-D interference contributes to raw asymmetry
- 2D test shows CP conservation is rejected at minimum 8.2σ for three approaches
 - $\Delta x/\Delta y$: CP -violating Cartesian coefficients



CP violation results: CPV in $\rho_3(1690)^0 K^+$




- Sizeable $\rho_3(1690)^0$ contribute above 1.5 GeV
- $\cos \theta_{\pi^+\pi^-} \sim -0.5$: $\rho_3(1690)^0$ responsible for the asymmetry
- S-D interference contribution near $\cos \theta_{\pi^+\pi^-} \sim 0$




The $K_0^*(1430)\pi$ BF and $f_X(1300)$ Puzzles

- **Previous:**
 - Larger $B^+ \rightarrow K_0^*(1430)^0 \pi^+$ BF favour pQCD
 - QCDf describes $B \rightarrow K_0^*(1430)[\eta', \rho, \omega, \phi]$
 - $f_X(1300)$
- **Belle and BaBar:** used BW or LASS shapes, relying on unitary-based arguments in the charmless $K\pi$ S-wave
- **LHCb:** GLASS allows independent coupling between the resonant and effective-range components, relaxing the LASS constraints.
- **Resolution:**
 - The smaller $K_0^*(1430)$ contribution obtained with GLASS leads to a BF consistent with QCDf, removing the previous preference for pQCD
 - **No $f_X(1300)$ required:** Improved $K\pi$ S-wave model describes data with the established $f_0(1370)$ and $f_0(1500)$



New intermediate processes

- Measure significance of each resonance
- Two approaches:
 - Fit fraction z-score
 - 2D deviation of amplitude coupling from (0,0)
- Ten intermediate processes are observed for the first time
- $\pi\pi$ - $\pi\pi$ rescattering help Isobar achieve better fit quality than K-matrix



Component	HFLAV (10^{-6})	Branching fraction (10^{-6})			
$B^+ \rightarrow K_2^*(1430)^0 \pi^+$	$5.6^{+2.2}_{-1.4}$	3.90 ± 0.57	± 0.31	± 0.13	
$B^+ \rightarrow f_2(1270) K^+$	1.07 ± 0.31	2.21 ± 0.30	± 0.30	± 0.09	
$B^+ \rightarrow K^*(1410)^0 \pi^+$	< 45 (90% CL)	9.17 ± 2.39	± 2.49	± 1.82	
$B^+ \rightarrow \rho_3(1690)^0 K^+$	—	1.00 ± 0.26	± 0.37	± 0.06	
$B^+ \rightarrow K_0^*(1950)^0 \pi^+$	—	1.32 ± 0.27	± 0.08	± 0.36	
$B^+ \rightarrow f_0(1370) K^{*+}$	< 11 (90% CL)	9.67 ± 0.70		± 0.23	
$B^+ \rightarrow f_0(1500) K^+$	—	4.65 ± 0.74		± 0.32	
$B^+ \rightarrow f_0(500) K^{*+}$	—	1.06 ± 0.22		± 0.03	
$B^+ \rightarrow f_0(1710) K^{*+}$	—	0.41 ± 0.34		± 0.01	
$B^+ \rightarrow (\pi\pi - \pi\pi) K^{*+}$	—	3.09 ± 1.03		± 0.08	

- Introduction
- Amplitude analysis
- Physics results and implications
 - CP violation
 - Strong interaction modelling implications
- Conclusion

Conclusion

- **An amplitude analysis** of $B^+ \rightarrow K^+ \pi^- \pi^+$ with CP violation has been performed **LHCb-PAPER-2025-067**
- **CP violation** **LHCb-PAPER-2025-069**
 - Observed in $B^+ \rightarrow \rho(770)^0 K^+$ and $B^+ \rightarrow f_2(1270)^0 K^+$ decays
 - Observed in $B^+ \rightarrow (S - P)K^+$ interference
 - Evidence for CP violation in $B^+ \rightarrow \rho_3(1690)^0 K^+$ decay
 - Evidence for CP violation in low-mass $\pi\pi$ S-wave region
 - Evidence for CP violation in $B^+ \rightarrow (S - D)K^+$ interference
- **Implications for strong-interaction modelling** **LHCb-PAPER-2025-068**
 - $B^+ \rightarrow K_0^*(1430)^0 \pi^+$ BF now agrees with QCDf, resolving a longstanding puzzle
 - No additional $f_X(1300)$ contribution required
 - Ten intermediate processes observed for the first time, with improved BF precision
- **Outlook**
 - Larger datasets now available at LHCb, but need reductions of both experimental and model unc.

Thanks for your attention

Backups→