

Measurements of the effective weak mixing angle in dimuon events at D0

Friday, 1 September 2017 16:30 (25 minutes)

We present the measurements of forward-backward charge asymmetry A_{FB} in $p\bar{p} \rightarrow Z/\gamma^* \rightarrow \mu^+\mu^-$ events using 9.7 fb^{-1} of $p\bar{p}$ data collected at $\sqrt{s} = 1.96 \text{ TeV}$ by the D0 detector at the Fermilab Tevatron collider. A_{FB} is measured as a function of the invariant mass of the dimuon system to extract the effective weak mixing angle $\sin^2 \theta_{eff}^{lep}$. In the context of the standard model, using the on-shell renormalization scheme where $\sin^2 \theta_W = 1 - M_W^2/M_Z^2$, measurements of $\sin^2 \theta_{eff}^{lep}$ yield indirect extractions of the W mass. The result will contribute to the final Tevatron combination of the weak mixing angle and indirect m_W measurements.

Primary authors: COLLABORATION, D0 (Fermilab); Dr HAN, Liang (USTC)

Presenters: COLLABORATION, D0 (Fermilab); Dr HAN, Liang (USTC)

Session Classification: Standard model physics at the TeV scale

Track Classification: 6) Standard model physics at the TeV scale