

The NEMO-3 and SuperNEMO experiments

M. Cascella
(for the NEMO-3 and SuperNEMO collaborations)



Introduction



- Double beta and neutrinoless double beta decay
- NEMO-3 results
- SuperNEMO demonstrator module



Double β decay

Analogue to β decay, for cases where the $Z \pm 1$ is forbidden (heavier nucleus) but $Z \pm 2$ is allowed.



Neutrinoless double β decay


- If neutrinos are Majorana fermions $0\nu 2\beta$ is allowed
- electrons $\sim$ few MeV



Why is $0\nu2\beta$ interesting?

- Lepton number violation
- New physics
- Neutrino mass and mass ordering



$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu} |\mathcal{M}_{0\nu}|^2 |m_{\beta\beta}|^2$$



NEMO-3


NEMO design principles

- Source independent from detector
- Compromise between calorimeter resolution / tracking
- Topological information, timing and particle tagging
- Background suppression and BSM decays




NEMO-3

- Active from 2003 to 2011
- Hosted in Frejus tunnel (1700m, 4800 w.e.m. 10^{-6} x cosmics)
- 7x 2β and several calibration sources




NEMO-3 structure



NEMO-3 structure


1. Source strips



NEMO-3 structure


1. Source strips

2. 3D tracking multi-wire Geiger cells




NEMO-3 structure

1. Source strips
2. 3D tracking multi-wire Geiger cells
3. PS scintillator with low activity PMTs






NEMO-3 structure

1. Source strips
2. 3D tracking multi-wire Geiger cells
3. PS scintillator with low activity PMTs
4. Cu/Fe frame



Bismuth and Radon

- Tracker essential in rejecting background
- Internal contaminations
- External radiation
- ^{214}Bi ($Q_\beta = 3.3 \text{ MeV}$)
 - α tagging





^{150}Nd

- Large $Q_{\beta\beta} = 3.4$ MeV and phase space (36.6 g of isotope)
- BDT using 10 kinematic variables
- Quadruple beta decay search

$$T_{1/2}^{0\nu\beta\beta} > 2.0 \times 10^{22} \text{ yr (90\% C.L.)}$$

$$T_{1/2}^{2\nu} = [9.34 \pm 0.22 \text{ (stat.) } {}^{+0.62}_{-0.60} \text{ (syst.)}] \times 10^{18} \text{ yr}$$

- Phys. Rev. D 94 072003




^{48}Ca


- highest $Q_{\beta\beta} = 4.3 \text{ MeV}$
- 7g of isotope
(~ 300 $2\nu 2\beta$ events)

$$T_{1/2}^{2\nu} = [6.4^{+0.7}_{-0.6}(\text{stat.})^{+1.2}_{-0.9}(\text{syst.})] \times 10^{19} \text{ yr}$$

- Phys. Rev. D 93, 112008



MO-3 - ^{48}Ca - 7 g, 5.25 y




SuperNEMO

SuperNEMO



NEMO-3		SuperNEMO
^{100}Mo	isotope	^{82}Se (or ^{150}Nd)
7 kg	isotope mass	7 (demonstrator) $\rightarrow$ 100 kg
5 mBq/m ³	radon	0.15 mBq/m ³
^{208}Tl : 100 $\mu\text{Bq/kg}$ ^{214}Bi : 300 $\mu\text{Bq/kg}$	internal contamination	$^{208}\text{Tl} \leq 2 \mu\text{Bq/kg}$ $^{214}\text{Bi} \leq 10 \mu\text{Bq/kg}$
14% @ 1 MeV	Calorimeter	8% @ 1 MeV

- Next gen demonstrator
- Modular design
- Better hermeticity




The SuperNEMO tracker





- Restricted materials: copper, steel, duracon. HPGe and Rn tested
- Robotic construction of 2034 cells (~45Km of wires)
- Controlled construction, assembly and testing conditions.

Radon measurements





1. Purge for several $T^{1/2}$

Radon measurements




1. Purge for several $T_{1/2}$
2. Flow through cooled carbon trap



Radon measurements




1. Purge for several $T_{1/2}$
2. Flow through cooled carbon trap
3. Release into electrostatic detector



Radon measurements



1. Purge for several $T_{1/2}$
2. Flow through cooled carbon trap
3. Release into electrostatic detector
4. Rn: $41.3 \pm 4.7 \text{ mBq}$ ($\rightarrow 150 \mu\text{Bq/m}^3$ with flushing)



SuperNEMO calorimeter



- 520 x 8" + 192 x 5" high q.e. PM
- 8% FWHM,
 $\sigma_t = 400$ ps @ 1 MeV
(1% stability)



SuperNEMO source foil

- Purification methods tested: distillation, chromatography, chemical precipitation
- All foils checked with “BiPo” to measure contamination at few $\mu\text{Bq/kg}$ level





arXiv:1702.07176




Current detector status

- Calorimeter and tracker delivered
- Source frame arriving next month
- Demonstrator running by end of year



Conclusions

- NEMO-3
 - World leading 0ν and $2\nu\beta\beta$ measurements
 - Unique new physics searches
- SuperNEMO demonstrator ($T_{1/2}^{0\nu} \sim 6.5 \cdot 10^{24} \text{y}$)
- Scales to 500 kg $\cdot$ yr ($T_{1/2}^{0\nu} \sim 10^{26} \text{y}$)
- Tracker-calorimeter technique powerful $\beta\beta$ physics probe



Thank you

Questions?



Majorana and Dirac models

Two models to give neutrinos their masses

Dirac

$$\begin{bmatrix} \nu_L \\ \nu_R \\ \bar{\nu}_L \\ \bar{\nu}_R \end{bmatrix}$$

ν_R and $\bar{\nu}_L$
exists, even if
we (currently)
have no way to
detect them



Majorana

$$\begin{bmatrix} \nu_L \\ \bar{\nu}_R \end{bmatrix}$$




ν are their own
antiparticle!


Why is $0\nu2\beta$ interesting

- Window into new physics
 - Supersymmetry
 - Right handed weak currents
 - Majorons
 - Doubly-charged Higgs bosons
 - Heavy neutral leptons
 - Light Majorana neutrino exchange



1S





$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu} |\mathcal{M}_{0\nu}|^2 |m_{\beta\beta}|^2$$

$$m_{\beta\beta} = \sum_k U_{ek}^2 m_k$$

