







# Higgs Physics at the CEPC

Manqi Ruan

On behalf of the CEPC Simulation Study Group

# Higgs @ CEPC





Process	Cross section	Events in 5 ab <sup>-1</sup>
Higgs boson production, cross section in fb		
$e^+e^- \rightarrow ZH$	212	$1.06 \times 10^6$
$e^+e^- \rightarrow \nu\bar{\nu}H$	6.72	$3.36 \times 10^4$
$e^+e^- \rightarrow e^+e^-H$	0.63	$3.15 \times 10^3$
Total	219	$1.10 \times 10^6$

Observables: Higgs mass, CP,  $\sigma(ZH)$ , event rates ( $\sigma(ZH, \nu\nu H) \cdot \text{Br}(H \rightarrow X)$ ), Diff. distributions

Derive: **Absolute** Higgs width, branching ratios, **couplings**

# CEPC Conceptual detector, developed from ILD






A detector reconstruct all the physics object (lepton, photon, tau, Jet, MET, ...) with high efficiency/precision

High Precision VTX located close to IP: b, c, tau tagging

High Precision Tracking system:  $\delta(1/Pt) \sim 2 \cdot 10^{-5} (\text{GeV}^{-1})$

PFA oriented Calorimeter System ( $\sim 10^8$  channels): Tagging, ID, Jet energy resolution, ect

# Detector performance



Acceptance	$ \cos(\theta)  < 0.995$ (from the inner radius of the outmost tracking disk)
Tracking Efficiency	For isolated charged particle with energy $> 1\text{ GeV}$ : $\sim 100\%$
Photon Reconstruction Efficiency	For isolated photon with energy $> 0.5\text{ GeV}$ : $\sim 100\%$
Tracker resolution	$\delta(1/P_T) = 2 \cdot 10^{-5} (\text{GeV}^{-1})$
ECAL intrinsic resolution	$\delta E/E = 16\%/\sqrt{E/\text{GeV}} \oplus 0.5\%$
HCAL intrinsic resolution	$\delta E/E = 60\%/\sqrt{E/\text{GeV}} \oplus 1\%$
Jet energy resolution	$\delta E/E = 4\%$
<b>Typical Distance for shower separation</b>	$< 3\text{ cm}$
<b>Lepton identification</b>	For charged particle with Energy $> 2\text{ GeV}$ : Lepton identification efficiency $> 99.5\%$ , $P(\text{hadron} \rightarrow \text{muon}) \sim P(\text{hadron} \rightarrow \text{electron})$ : 1%
b-tagging	At Z pole samples & $\text{eff}(b \rightarrow b) = 80\%$ , $P(\text{uds} \rightarrow b) < 1\%$ , $P(c \rightarrow b) \sim 10\%$
c-tagging	At Z pole samples & $\text{eff}(c \rightarrow c) = 60\%$ , $P(\text{uds} \rightarrow c) = 7\%$ , $P(b \rightarrow c) = 12\%$



Performance at full reconstruction

*Benchmark separation distance  $< 3\text{ cm}$   
(Testing on 10 GeV Pion + 5 GeV Photon Sample)*


# Higgs analyses at Pre-CDR

$$\mathcal{L} = \mathcal{L}_{SM} + \sum_i \frac{c_i}{M^2} \mathcal{O}_{6,i} \quad \delta \sim c_i \frac{v^2}{M^2}$$

% precision  $\rightarrow$   $M \sim 1$  TeV  
 to new physics  $\rightarrow \sim \times 10$  over LHC




Kappa-framework  
 model-dependent measurements



Absolute measurements

# Status now






~50 independent analysis at Full Simulation level

	PreCDR (Jan 2015)	Now (Aug 2016)
$\sigma(\text{ZH})$	0.51%	0.50%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{bb})$	0.28%	0.21%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{cc})$	2.1%	2.5%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{gg})$	1.6%	1.2%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{WW})$	1.5%	1.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{ZZ})$	4.3%	4.3%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \pi\pi)$	1.2%	1.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \gamma\gamma)$	9.0%	9.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{Z}\gamma)$	-	$\sim 4 \sigma$
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \mu\mu)$	17%	12%
$\sigma(\text{vvH}) \cdot \text{Br}(\text{H} \rightarrow \text{bb})$	2.8%	2.8%
<b>Higgs Mass/MeV</b>	5.9	5.0
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{inv})$	95%. CL = 1.4e-3	1.4e-3
$\text{Br}(\text{H} \rightarrow \text{ee}/\text{emu})$	-	1.7e-4/1.2e-4
$\text{Br}(\text{H} \rightarrow \text{bb}\gamma\gamma)$	$< 10^{-3}$	3.0e-4

# Model-independent measurement of $\sigma(\text{ZH})$

Zhenxing Chen & Yacine Haddad



- Recoil mass method. Combined precision:  
 $\delta\sigma(\text{ZH})/\sigma(\text{ZH}) = 0.5\%$  -  
 $\delta g(\text{HZZ})/g(\text{HZZ}) = 0.25\%$
- Indirect Access to  $g(\text{HHH})$

$$\sigma_{Zh} = \left| \begin{array}{c} e \\ \text{---} \\ e \end{array} \right. \begin{array}{c} \text{---} \\ Z \\ \text{---} \\ h \end{array} \left. \right|^2 + 2 \text{Re} \left[ \begin{array}{c} e \\ \text{---} \\ e \end{array} \right. \begin{array}{c} \text{---} \\ z \\ \text{---} \\ h \end{array} \cdot \left( \begin{array}{c} e^+ \\ \text{---} \\ e^- \end{array} \right. \begin{array}{c} \text{---} \\ Z \\ \text{---} \\ h \end{array} + \begin{array}{c} e^+ \\ \text{---} \\ e^- \end{array} \left. \begin{array}{c} \text{---} \\ Z \\ \text{---} \\ h \end{array} \right) \right]$$

$$\delta_{\pi}^{240} = 100 (2\delta_Z + 0.014\delta_h) \%$$



- M. McCullough, 1312.3322

# Workflow for Br(H->bb, cc, gg) measurements general event selection + Template fit

## 2. Selection

Cut Definition	Sig.	qq	qqnn	qqln	xxh
FSClasser output	148955	25M	183687	3698817	63194
$N_{\text{PFO}(E>0.4\text{GeV})} > 20$	148808	23M	163088	3439927	58882
$110 < E_{\text{total}} < 150$	132561	10M	125878	705357	34215
$P_T > 19$	126006	34198	116314	627602	32300
Isolation lepton veto	123586	33775	115867	327206	23773
$100 < M_{\text{inv}} < 135$	117845	9506	10420	162511	21277
$70 < M_{\text{rec}} < 125$	111886	7521	10045	110426	20458
$0.15 < y_{12} < 1$	111353	7405	9702	101797	19983
$y_{23} < 0.06$	105078	6644	8456	69313	14495
$y_{34} < 0.008$	100117	6504	7878	58532	6899
$-0.98 < \cos(\theta_{\text{included}}^{2\text{jets}}) < -0.4$	97277	5178	5365	33293	6273
$BDT > -0.01$	76666	344	118	69	1594
<b>Significance</b>			<b>265.20</b>		
<b>Efficiency</b>			<b>51.5%</b>		

## 3. BDT & final results



## Flavor tagging





Figure 7: The B-C Likelihood characteristics for Signal and other Higgs Background. The Standard Model Background isn't included in because there is no B-C Likelihood.

## vvH events

## Template fit




Template: 2000 fb-1  
Data: 500 fb-1  
Fit Times: 5000

No SM bkg in template fit,  
since no full simulation



# Br(H→WW)

$H \rightarrow WW/ZZ$ : Portal to Higgs width & perfect test bed for detector/reconstruction performance...



Expected Number of events with different objects

	Z→ll	tautau	vv	qq
H→WW*→4q	6.91k	3.45k	19.74k	69.1k
μνqq	2.27k	1.14k	6.47k	22.7k
eνqq	2.27k	1.14k	6.47k	22.7k
eeνν	186	93	527	1.9k
μμνν	186	93	527	1.9k
eμνν	372	186	1154	3.7k
X + tau	3.2k	1.6k	9.14k	32.0k

	Extrapolated from ILC results
	Await for tau finder
	Await for the SM Background simulation
	Full Simulation
	Preliminary result acquired
	Unexplored

- Br(H→WW), Combined accuracy ~ 1.0% from 13 independent full simulation analyses
  - 1.45% at llH,  $H \rightarrow WW^* \rightarrow$  inc channels, 12 independent channels.
  - ~ 1.7% at ννH,  $H \rightarrow WW^* \rightarrow 4q$  channel (Preliminary. ILC extrapolation = 2.3%)
  - 2.3% at qqH,  $H \rightarrow WW^* \rightarrow 2ql\nu$  channel (extrapolated from ILC full simulation)
- High efficiency in event reconstruction


# Br(H → WW)

ZH, H → WW*	Yield	Object reconstructed	Isolation	Signal Efficiency	Main Background	Accuracy	Combined
Z(μμ)H(evev)	88	76(86.36%)	61(80.26%)	36(40.91%)	4(ZH)	17.57%	2.68%
Z(μμ)H(μνμν)	89	80(89.89%)	77(96.25%)	52(58.43%)	6(ZH&ZZ)	14.65%	
Z(μμ)H(evμν)	174	157(90.23%)	147(93.63%)	105(60.34%)	0	9.76%	
Z(μμ)H(evqq)	1105	1042(94.30%)	864(82.92%)	663(60.00%)	45(ZH)	4.02%	
Z(μμ)H(μνqq)	1110	1056(95.14%)	988(93.56%)	717(64.59%)	159(ZH&ZZ)	4.13%	
Z(μμ)H(qqqq)	Preliminary						3.0%
Z(ee)H(evev)	91	62(68.13%)	60(96.77%)	22(24.16%)	16(SZ)	28.02%	2.87%
Z(ee)H(μνμν)	82	63(76.83%)	63(100%)	44(53.66%)	24(SZ)	18.74%	
Z(ee)H(evμν)	178	132(74.16%)	124(93.94%)	82(46.07%)	25(ZH&SZ)	12.61%	
Z(ee)H(evqq)	1182	1041(88.07%)	916(87.99%)	621(51.78%)	188(SZ&ZH)	4.62%	
Z(ee)H(μνqq)	1221	1194(97.79%)	1048(87.77%)	684(56.02%)	49(ZH&SZ)	3.96%	
Z(ee)H(qqqq)	Preliminary estimation						3.2%

- Full Simulation on 12 independent channels
  - Very high object reconstruction efficiency
  - Combined result: 1.1%
- Extrapolation from other ILC channels: 2.2%
- Combined: 1.0%

	Z → ll	tautau	νν	qq
H → WW* → 4q	1.45%	3.45k	< 1.9%	69.1k
μνqq		1.14k	6.47k	2.2%
evqq		1.14k	6.47k	
eeνν		93	527	1.9k
μμνν		93	527	1.9k
eμνν		186	1154	3.7k
X + tau		3.2k	1.6k	9.14k

# Br(H→ZZ)



Expected Number of events with different objects

	Z→ll	tautau	vv	qq
H→ZZ*→4q	888	444	3.10k	9.24k
2v + 2q	508	254	1.77k	5.29k
2l + 2q	170	85	596	1.8k
4v	73	36	254	756
2l + 2v	49	24	170	508
4l	8	4	28	86
X + tau	120	60	418	1246

	More than 2 jets, Await for sophisticated Jet Clustering
	Await for tau finder
	limited accuracy ~ > 50%
	Explored by H->invisible analysis -> Accuracy ~ 40%
	Promising channels
	Unexplored

- Br(H→ZZ), explored at 18 different channels with full simulation (llvvqq, 4lqq, ll4q, 2l4v)
  - 8 Channels has individual accuracy better than 25%: Combined accuracy ~ **5.4%**
  - 8 with accuracy worse than 25 - 50%
  - 2 with accuracy worse than 50% (llH, H→ZZ→4q and vvH, H→ZZ→llvv)
- High efficiency in event reconstruction

# Br(H→ZZ)

ZZZ*	Yield	Object reconstructed	Signal Efficiency(%)	Main Background	Accuracy (%)	Comments
μμννqq	128	118	63.3	h->ww&zz_sl	<b>12.9</b>	Tau finder would be highly appreciated
μμqqνν	128	125	-	h->bb&zz_sl	>25	
eeννqq	132	91	53.8	h->ww&sze_sl	<b>15.8</b>	Reconstructed efficiency of electron need to be improved
eeqqνν	132	88	-	h->bb&zz_sl	>25	
ννμμqq	158	144	61.4	h->t,w&zz_sl	<b>11.0</b>	
ννqqμμ	158	149	51.9	h->w,b&zz_sl	<b>12.9</b>	
ννeeqq	151	118	43.1	h->w&sze_sl	<b>21.3</b>	
ννqqee	151	134	-	h->bb&sze_sl	>25	
qqμμνν	135	115	-	h->tt&zz_sl	>25	Compare to ll recoil, qq recoil mass has much worse distinguishing power to SM background
qqννμμ	135	122	-	h->t,w&zz_sl	>25	
qqeeνν	127	107	-	h->tt&sze_sl	>25	
qqννee	127	123	-	h->t,w&sze_sl	>25	
μμμμqq/qqμμ	43	39	69.8	h->tt&zz_sl	<b>19.9</b>	Tau finder & Electron Reconstruction
μμeeqq/qqee	43	39	60.5	h->tt&zz_sl	<b>21.2</b>	
eeeeqq/eeqqee	43	33	-	h->tt&sze_sl	>25	
eeμμqq/eeqqμμ	43	41	58.2	h->tt&sze_sl	<b>19.9</b>	

Full Simulation analysis performed on 16 independent channels.

8 Channels acquire accuracy better than 25%.

Combined accuracy: **5.4%**


*If electron id efficiency ~ muon id: **4.8%***

***If tau finder (used for veto) is mature: ??***

TLEP extrapolation: **4.3%**

# Higgs width measurement

- $g^2(\text{HXX}) \sim \Gamma_{\text{H} \rightarrow \text{XX}} = \Gamma_{\text{total}} * \text{Br}(\text{H} \rightarrow \text{XX})$
- Branching ratios: determined simply by
  - $\sigma(\text{ZH})$  and  $\sigma(\text{ZH}) * \text{Br}(\text{H} \rightarrow \text{XX})$
- $\Gamma_{\text{total}}$ : determined from:
  - From  $\sigma(\text{ZH})$  ( $\sim g^2(\text{HZZ})$ ) and  $\sigma(\text{ZH}) * \text{Br}(\text{H} \rightarrow \text{ZZ})$  ( $\sim g^4(\text{HZZ}) / \Gamma_{\text{total}}$ )
  - From  $\sigma(\text{ZH}) * \text{Br}(\text{H} \rightarrow \text{bb})$ ,  $\sigma(\text{vvH}) * \text{Br}(\text{H} \rightarrow \text{bb})$ ,  $\sigma(\text{ZH}) * \text{Br}(\text{H} \rightarrow \text{WW})$ ,  $\sigma(\text{ZH})$





$\text{Br}(\text{H} \rightarrow \text{ZZ})$ : relative error of 6.9% achieved with  $\text{ZH} \rightarrow \text{ZZZ}^* \rightarrow \text{vv}(\text{Z})\text{llqq}(\text{H})$  final states.  
 Extrapolation of TLEP result leads to 4.3% relative error

$\sigma(\text{vvH}) * \text{Br}(\text{H} \rightarrow \text{bb})$ : relative error of 2.8%

A combined accuracy of 2.8% for the Higgs total width measurements

# Br(H→ $\tau\tau$ ) & Br(H→Z $\gamma$ )









Br(H→ $\tau\tau$ ): 3% accuracy acquired by  $\mu\mu H$  channel; overall accuracy < 1% (Dan Yu)

H→Z $\gamma$  events: 4 $\sigma$  signal (Weiming Yao)

# Higgs rare decay

Feng Wang, Jianhuan Xiang, Yitian Li. etc

Binlong Wang, Zhenwei Cui






18/11/2016

$\text{Br}(H \rightarrow \gamma\gamma): \sim 10\%$

$\text{Br}(H \rightarrow \mu\mu): \sim 12\%$   
(Exclusive analysis)

# Exotic: Higgs invisible decays

Assuming  $\sigma(ZH) \cdot Br(H \rightarrow inv) = 200 \text{ fb}$





Invisible up limit at CEPC: 0.14% at 95% C.L



# Up limit of $\text{Br}(H \rightarrow ee)$ & $\text{Br}(H \rightarrow e\mu)$


Lei Wang

Assuming  $\sigma(\text{ZH}) \cdot \text{Br}(H \rightarrow ee/e\mu) = 200 \text{ fb}$



95% up limit:  $\text{Br}(H \rightarrow ee) = 1.7e-4$ ;  
 $\text{Br}(H \rightarrow e\mu) = 1.2e-4$ ;






# H → Exotic, Hadronic



- Typical processes in 2HDM & NMSSM
- Joint efforts by HK Cluster and IHEP
  - Study proposed by T. Liu
  - Main analyzer, Jiawei Wang, Kevin & Zhenxing Chen
- 95% CL up limit  $\sim \mathcal{O}(10^{-4})$ .

# H->Exotic, hadronic

Para:  $M(\text{LSP}) = 0$ ;  $M(h_0) = 15 \text{ GeV}$ ;  $M(\text{NLSP}) = 20 \text{ GeV}$




Object found	Cut 1 $m_{\mu}$	Cut 2 $m_{\text{rec}}$	Cut 3 b likeness	Cut 4 $m_j$
Signal	17	15	12	10
ZH BGs	34093	30732	16026	4
ZZ BGs	538790	281198	30825	20

Cut3:  $\text{sum}(\text{B-likeness}) > 0.9$

- 95% CL. Uplimit set to be  $5E-4$ ; will be significantly improved by including di-electron/tau channel...
- ISR effect not included in the Signal sample.  $\sigma(\text{ZH})$  referred to SM Xsec of 200 fb. Effect on uplimit setting could be ignored

# H → Exotic, Hadronic





## Benchmark Points

Scan over the parameter space for sensitivity:

- Fix  $m_{\tilde{\chi}_1^0} = 0$  GeV and make exclusion contours on the  $m_{h^0}$  and  $m_{\tilde{\chi}_2^0}$  plane with the range:
  - $10 \text{ GeV} < m_{h^0} < 60 \text{ GeV}$  (15,25,35,45,55 GeV)
  - $10 \text{ GeV} < m_{\tilde{\chi}_2^0} < 125 \text{ GeV}$  (20,40,60,80,100,120 GeV)
- Fix  $m_{h^0} = 30$  GeV and make exclusion contours on the  $m_{\tilde{\chi}_1^0}$  and  $m_{\tilde{\chi}_2^0}$  plane, with the range:
  - $0 \text{ GeV} < m_{\tilde{\chi}_1^0} < 60 \text{ GeV}$  (5,15,25,35,45,55 GeV)
  - $10 \text{ GeV} < m_{\tilde{\chi}_2^0} < 125 \text{ GeV}$  (20,40,60,80,100,120 GeV)

Suggested by prof. Liu



# Summary

- CEPC, an electron-positron Higgs factory & an precision EW machine, expecting 1M Higgs bosons
  - $\mathcal{O}(0.1-1\%)$  level accuracy in absolute measurement of Higgs Branching ratio and couplings
  - Higgs total width measured to 2.8%
  - Good access to SM Higgs rare decays ( $\mu\mu$ ,  $\gamma\gamma$ ,  $Z\gamma$ )
  - Higgs exotic decays, limited to better than 0.1% level
  - EW Program significantly enhance the access to the New Physics
- Complementary to LHC & Linear Colliders
  - + LHC, enhance the access to rare decays;
  - + Linear Colliders, enhance the measurement to the Higgs boson width;
  - Both provide direct access to  $g(H\tau\tau)$  &  $g(HHH)$

Back

IHEP-CEPC-DR-2015-01

IHEP-EP-2015-01

IHEP-TH-2015-01

IHEP-CEPC-DR-2015-01

IHEP-AC-2015-01

**Can be downloaded from**

<http://cepc.ihep.ac.cn/preCDR/volume.html>

# CEPC-SPPC

*Preliminary Conceptual Design Report*

Volume I - Physics & Detector

**403 pages, 480 authors**

The CEPC-SPPC Study Group

March 2015

# CEPC-SPPC

*Preliminary Conceptual Design Report*



Volume II - Accelerator

**328 pages, 300 authors**

The CEPC-SPPC Study Group

March 2015



# CEPC-SPPC



- Electron-positron collision phase
  - Higgs factory: collision at  $\sim 240 - 250$  GeV center-of-mass energy, Instant luminosity  $\sim 2 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ , 1M clean Higgs event at 2 IP over 10 years
  - Z pole operation for precise EW measurement
- Proton-Proton collision phase
  - center-of-mass energy constrained by tunnel circumference and high-field dipole
  - Peak luminosity  $\sim 1 \cdot 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$  (*ArXiv: 1504.06108, discussion on needed Luminosity*)
- Tunnel circumference: 54 km in the baseline design. Longer tunnel to be evaluated.




# EW Physics




From 10 Billion Z bosons + the data at Higgs runs

# New Physics Reach via dim-6 operators

## Sensitivities from Existing EWPO & Future HO



## Sensitivity from EWPO+HO+Z-Pole




1603.03385

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_{ij} \frac{y_{ij}}{\Lambda \sim 10^{14} \text{GeV}} (\bar{L}_i \tilde{\mathbf{H}}) (\tilde{\mathbf{H}}^\dagger L_j) + \sum_i \frac{c_i}{\Lambda^2} \mathcal{O}_i.$$


Higgs	EW Gauge Bosons	Fermions
$\mathcal{O}_H = \frac{1}{2}(\partial_\mu  \mathbf{H} ^2)^2$	$\mathcal{O}_{WW} = g^2  \mathbf{H} ^2 W_{\mu\nu}^a W^{a\mu\nu}$	$\mathcal{O}_L^{(3)} = (i\mathbf{H}^\dagger \overleftrightarrow{D}_\mu \mathbf{H})(\bar{\Psi}_L \gamma^\mu \sigma^a \Psi_L)$
$\mathcal{O}_T = \frac{1}{2}(\mathbf{H}^\dagger \overleftrightarrow{D}_\mu \mathbf{H})^2$	$\mathcal{O}_{BB} = g^2  \mathbf{H} ^2 B_{\mu\nu} B^{\mu\nu}$	$\mathcal{O}_{LL}^{(3)} = (\bar{\Psi}_L \gamma^\mu \sigma^a \Psi_L)(\bar{\Psi}_L \gamma^\mu \sigma^a \Psi_L)$
	$\mathcal{O}_{WB} = gg' \mathbf{H}^\dagger \sigma^a \mathbf{H} W_{\mu\nu}^a B^{\mu\nu}$	$\mathcal{O}_L = (i\mathbf{H}^\dagger \overleftrightarrow{D}_\mu \mathbf{H})(\bar{\Psi}_L \gamma^\mu \Psi_L)$
	$\mathcal{O}_{HW} = ig(D^\mu \mathbf{H})^\dagger \sigma^a (D^\nu \mathbf{H}) W_{\mu\nu}^a$	$\mathcal{O}_R = (i\mathbf{H}^\dagger \overleftrightarrow{D}_\mu \mathbf{H})(\bar{\psi}_R \gamma^\mu \psi_R)$
	$\mathcal{O}_g = g_s^2  \mathbf{H} ^2 G_{\mu\nu}^a G^{a\mu\nu}$	
	$\mathcal{O}_{HB} = ig'(D^\mu \mathbf{H})^\dagger (D^\nu \mathbf{H}) B_{\mu\nu}$	

# Status



Br(H→ττ) via Z to ll



• Initial study of  $zH \rightarrow z\gamma \rightarrow qq\nu\nu\gamma$  is promising to be  $4\sigma$  with  $5\text{ ab}^{-1}$ .




	PreCDR (Jan 2015)	Now (Aug 2016)
$\sigma(\text{ZH})$	0.51%	0.50%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{bb})$	0.28%	0.21%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{cc})$	2.1%	2.5%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{gg})$	1.6%	1.3%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{WW})$	1.5%	1.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{ZZ})$	4.3%	4.3%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \tau\tau)$	1.2%	1.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \gamma\gamma)$	9.0%	9.0%
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{Z}\gamma)$	-	$\sim 4\sigma$
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \mu\mu)$	17%	17%
$\sigma(\text{v}\nu\text{H}) \cdot \text{Br}(\text{H} \rightarrow \text{bb})$	2.8%	2.8%
Higgs Mass/MeV	5.9	5.0
$\sigma(\text{ZH}) \cdot \text{Br}(\text{H} \rightarrow \text{inv})$	95%. CL = $1.4\text{e-}3$	$1.4\text{e-}3$
$\text{Br}(\text{H} \rightarrow \text{ee}/\text{emu})$	-	$1.7\text{e-}4/1.2\text{e-}4$
$\text{Br}(\text{H} \rightarrow \text{bb}\chi\chi)$	$<10^{-3}$	$3.0\text{e-}4$




## *From Hits to Final State Particles*


*Goal: ... Access the origin of  
every detector hit ...*




$Z \rightarrow 2 \text{ muon},$   
 $H \rightarrow 2 \text{ b}$




$Z \rightarrow 2 \text{ jet},$   
 $H \rightarrow 2 \text{ tau}$



$ZH \rightarrow 4 \text{ jets}$




$Z \rightarrow 2 \text{ muon}$   
 $H \rightarrow WW^* \rightarrow eevv$





CMS Experiment at the LHC, CERN  
Data recorded: 2012-May-27 23:35:47.271030 GMT  
Run/Event: 195099 / 137440354



Specific Final State...  
Overlap with lots of PU events


# $e^+e^-$ and pp complementary

$\mathcal{O}(10^{9-10})$  Higgs at SPPC

Event rates measured at pp collision  $\sigma \cdot BR(X \rightarrow H \rightarrow Y) = \sigma_X \frac{\Gamma_Y}{\Gamma_{tot}}$

$e^+e^-$  collider: provide anchor for absolute measurement (total width, etc)


pp collision has direct access to  $g(HHH)$  &  $g(Htt)$  and better access to Higgs rare decays



**ttH/ttZ as a precision probe of the top Yukawa coupling**



MLM, Frederix



	HL-LHC	HE-LHC	VLHC
$\sqrt{s}$ (TeV)	14	33	100
$\int \mathcal{L} dt$ ( $\text{fb}^{-1}$ )	3000	3000	3000
$\sigma \cdot BR(pp \rightarrow HH \rightarrow bb\gamma\gamma)$ (fb)	0.089	0.545	3.73
$S/\sqrt{B}$	2.3	6.2	15.0
$\lambda$ (stat)	50%	20%	8%

	$\delta\sigma(ttH)$	$\delta\sigma(ttZ)$	$\delta[\sigma(ttH)/\sigma(ttZ)]$
14 TeV	$\pm 4.8\%$	$\pm 5.3\%$	$\pm 0.75\%$
100 TeV	$\pm 2.7\%$	$\pm 2.3\%$	$\pm 0.48\%$

ArXiv: 1310.8361 [hep-ex]




# Higgs measurements at electron/positron & proton colliders

	Productivity	Finding efficiency	Remarks
LHC	Run 1: $10^6$ Run 2/HL: $10^{7-8}$	$\sim \mathcal{O}(10^{-3})$	Lots of Pile Up; Large theoretical/systematic uncertainties. Access to signal strength in major decay channels; Access to $g(\text{HHH})/g(\text{Htt})$ .
CEPC	$10^6$	$\sim \mathcal{O}(1)$	Absolute measurements in very clean environment; $\sim 0.1\%$ accuracy on key observable ( $g(\text{HZZ})$ ); Excellent precision to total width, invisible/exotic decay ratios; Indirect constrain to $g(\text{HHH})/g(\text{Htt})$ ;
SPPC	$10^{9-10}$	?	Good access to Higgs rare decay/generation, $g(\text{HHH})/g(\text{Htt})$ ,

*High complementarity between electron-positron & pp colliders*



# EW@CEPC




- EW precision measurements with significantly reduced uncertainties:

$$R_b, A_{FB}^b, \sin \theta_W^{eff}, m_Z, m_W, N_\nu \dots$$

	Present data	CEPC fit
$\alpha_s(M_Z^2)$	$0.1185 \pm 0.0006$ [23]	$\pm 1.0 \times 10^{-4}$ [24]
$\Delta\alpha_{had}^{(5)}(M_Z^2)$	$(276.5 \pm 0.8) \times 10^{-4}$ [25]	$\pm 4.7 \times 10^{-5}$ [26]
$m_Z$ [GeV]	$91.1875 \pm 0.0021$ [27]	<b><math>\pm 0.0005</math></b>
$m_t$ [GeV] (pole)	$173.34 \pm 0.76_{exp} [28] \pm 0.5_{th} [26]$	$\pm 0.2_{exp} \pm 0.5_{th} [29, 30]$
$m_h$ [GeV]	$125.14 \pm 0.24$ [26]	$< \pm 0.1$ [26]
$m_W$ [GeV]	$80.385 \pm 0.015_{exp} [23] \pm 0.004_{th} [31]$	$(\pm 3_{exp} \pm 1_{th}) \times 10^{-3}$ [31]
$\sin^2 \theta_{eff}^\ell$	$(23153 \pm 16) \times 10^{-5}$ [27]	<b><math>(\pm 2.3_{exp} \pm 1.5_{th}) \times 10^{-5}</math> [32]</b>
$\Gamma_Z$ [GeV]	$2.4952 \pm 0.0023$ [27]	<b><math>(\pm 5_{exp} \pm 0.8_{th}) \times 10^{-4}</math> [33]</b>
$R_b \equiv \Gamma_b/\Gamma_{had}$	$0.21629 \pm 0.00066$ [27]	<b><math>\pm 1.7 \times 10^{-4}</math></b>
$R_\ell \equiv \Gamma_{had}/\Gamma_\ell$	$20.767 \pm 0.025$ [27]	<b><math>\pm 0.007</math></b>

# Higgs @ LHC



*PP collider: High productivity but low finding efficiency  
~already  $10^6$  Higgs in Run 1 data...*

*Higgs signal: found via the decay final states.*

$$\sigma(AA \rightarrow H \rightarrow BB) \sim g^2(HAA)g^2(HBB)/\Gamma_{total}$$

proton - (anti)proton cross sections

